CORRECTED VERSION

(19) World Intellectual Property Organization International Bureau



TO THE CONTROL OF STATE OF STATE AND STATE OF S

(43) International Publication Date 4 April 2002 (04.04.2002)

PCT

(10) International Publication Number WO 02/026968 A2

- (51) International Patent Classification7:
- -----

C12N 15/11

- (21) International Application Number: PCT/CA01/01379
- (22) International Filing Date:

27 September 2001 (27.09.2001)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09/672,717

28 September 2000 (28.09.2000) US

- (71) Applicants: UNIVERSITY OF OTTAWA [CA/CA]; Suite 2213, 451 Smyth Road, Ottawa, Ontario K1H 8M5 (CA). AEGERA THERAPEUTICS, INC. [CA/CA]; 810 Chemin du Golf, Verdun, Quebec H3E 1A8 (CA).
- (72) Inventors: KORNELUK, Robert, G.; 1901 Tweed Avenue, Ottawa, Ontario K1G 2L8 (CA). LACASSE, Eric; 1727 Featherston Drive, Ottawa, Ontario K1H 6P3 (CA). BAIRD, Stephen; 20 Julian Avenue, Ottawa, Ontario K1Y 0S5 (CA). HOLCIK, Martin; Apartment 9, 210 Stewart Street, Ottawa, Ontario K1N 6K2 (CA). YOUNG, Sean; 1903 West 14th Avenue, Vancouver, British Columbia V6J 2K1 (CA).
- (74) Agents: ROBINSON, J., Christopher et al.; Smart & Biggar, Box 11560, Suite 2200, 650 West Georgia Street, Vancouver, British Columbia V6B 4N8 (CA).

81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- (48) Date of publication of this corrected version:

15 August 2002

(15) Information about Correction:

see PCT Gazette No. 33/2002 of 15 August 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTISENSE IAP NUCLEIC ACIDS AND USES THEREOF

(57) Abstract: The present invention feature antisense IAP nucleic acids and other negative regulators of the IAP anti-apoptotic pathway, and methods for using them to enhance apoptosis.

ANTISENSE IAP NUCLEIC ACIDS AND USES THEREOF

Field of the Invention

The invention relates to antisense IAP nucleic acids and methods of using them to increase apoptosis.

10

15

20

30

Background of the Invention

One way by which cells die is referred to as apoptosis, or programmed cell death. Apoptosis often occurs as a normal part of the development and maintenance of healthy tissues. The process may occur so rapidly that it is difficult to detect.

The apoptosis pathway is now known to play a critical role in embryonic development, viral pathogenesis, cancer, autoimmune disorders, and neurodegenerative diseases, as well as other events. The failure of an apoptotic response has been implicated in the development of cancer, autoimmune disorders, such as lupus erythematosis and multiple sclerosis, and in viral infections, including those associated with herpes virus, poxvirus, and adenovirus.

Baculoviruses encode proteins that are termed inhibitors of apoptosis (IAPs) because they inhibit the apoptosis that would otherwise occur when insect cells are infected by the virus. These proteins are thought to work in a manner that is independent of other viral proteins. The baculovirus IAP genes include sequences encoding a ring zinc finger-like motif (RZF), which is presumed to be directly involved in DNA binding, and two N-terminal domains that consist of a 70 amino acid repeat motif termed a BIR domain (Baculovirus IAP Repeat).

37

5

10

15

20

25

The role of apoptosis in cancer has only recently been appreciated. The identification of growth promoting "oncogenes" in the late 1970's gave rise to an almost universal focus on cellular proliferation that dominated research in cancer biology for many years. Long-standing dogma held that anti-cancer therapies preferentially targeted rapidly dividing cancer cells relative to "normal" cells. This explanation was not entirely satisfactory, since some slow growing tumors are easily treated, while many rapidly dividing tumor types are extremely resistant to anticancer therapies. Progress in the cancer field has now led to a new paradigm in cancer biology wherein neoplasia is viewed as a failure to execute normal pathways of programmed cell death. Normal cells receive continuous feedback from their neighbors through various growth factors, and commit "suicide" if removed from this context. Cancer cells somehow ignore these commands and continue inappropriate proliferation. Cancer therapies, including radiation and many chemotherapies, have traditionally been viewed as causing overwhelming cellular injury. New evidence suggests that cancer therapies actually work by triggering apoptosis.

Both normal cell types and cancer cell types display a wide range of susceptibility to apoptotic triggers, although the determinants of this resistance are only now under investigation. Many normal cell types undergo temporary growth arrest in response to a sub-lethal dose of radiation or cytotoxic chemical, while cancer cells in the vicinity undergo apoptosis. This provides the crucial treatment "window" of appropriate toxicity that allows successful anti-cancer therapy. It is therefore not surprising that resistance of tumor cells to apoptosis is emerging as a major category of cancer treatment failure.

Compared to the numerous growth-promoting oncogenes identified to date (>100), relatively few genes have been isolated that regulate

apoptosis. The Bcl-2 gene was first identified as an oncogene associated with the development of follicular lymphomas. In contrast to all other oncogenes identified to date, Bcl-2 displays no ability to promote cell proliferation, and instead has been demonstrated to suppress apoptosis by a variety of triggers. Elevated Bcl-2 expression is associated with a poor prognosis in neuroblastoma, prostate and colon cancer, and can result in a multidrug resistant phenotype *in vitro*. Although the study of Bcl-2 has helped revolutionize cancer paradigms, the vast majority of human malignancies do not demonstrate aberrant Bcl-2 expression.

10

15

In contrast to the findings with Bcl-2, mutation of the p53 tumor suppresser gene has been estimated to occur in up to 50% of human cancers and is the most frequent genetic change associated with cancer to date. The p53 protein plays a crucial role in surveying the genome for DNA damage. The cell type and degree of damage determines whether the cell will undergo growth arrest and repair, or initiate apoptosis. Mutations in p53 interfere with this activity, rendering the cell resistant to apoptosis by a wide range of cellular insults. Some progress has been made in understanding the molecular biology of p53, but many questions remain. p53 is known to function as a transcription factor, with the ability to positively or negatively regulate the expression of a variety of genes involved in cell cycle control, DNA repair, and apoptosis (including the anti-apoptotic Bcl-2 gene described above and the related pro-apoptotic gene Bax). The drug resistant phenotype conferred by p53 alterations has been linked to Bcl-2/Bax regulation, but this correlation does not hold for most cancer types, leaving open the possibility that other critical genes regulated by p53 remain to be identified.

Summary of the Invention

We have discovered that inhibitor of apoptosis (IAP) protein overexpression is associated with a wide range of cancer types including ovarian cancer, adenocarcinoma, lymphoma, and pancreatic cancer. In addition, we have found that nuclear localization, fragmentation of the IAPs, and overexpression of the IAPs in the presence of p53 mutations correlate with a cancer diagnosis, a poor prognosis, and resistance to numerous chemotherapeutic cancer drugs. These discoveries provide diagnostic, prognostic, and therapeutic compounds and methods for the detection and treatment of proliferative diseases. One way in which the expression of an IAP in a cell can be decreased is by administering to the cell a negative regulator of the IAP apoptotic pathway, for example, an antisense nucleic acid.

10

15

20

25

In general, the invention features methods and reagents useful for inducing apoptosis in a cell. The methods and reagents of the invention are useful in treating cancers, and other proliferative diseases.

In a first aspect, the invention features an inhibitor of apoptosis (IAP) antisense nucleic acid that inhibits IAP biological activity, regardless of the length of the antisense nucleic acid. In preferred embodiments, the IAP is XIAP, HIAP1, or HIAP2. In other preferred embodiments, the antisense nucleic acid is mammalian, for example, mouse or human. In yet another embodiment, the antisense nucleic acid is between 8 and 30 nucleotides in length.

In still other further preferred embodiments, the XIAP antisense is chosen from any one of SEQ ID NOS: 1 through 96, and the HIAP1 antisense is chosen from any one of SEQ ID NOS: 97 through 194. Preferably the IAP biological activity is inhibition of apoptosis or inhibition of IAP RNA or polypeptide expression. The antisense nucleic acid may comprise at least one modified internucleoside linkage.

4

Preferably the modified internucleoside linkage is a phosphorothioate, a methylphosphonate, a phosphotriester, a phosphorodithioate, or a phosphoselenate linkage. In addition, the antisense nucleic acid may comprise at least one modified sugar moiety. Preferably this modified sugar moiety is a 2'-O methoxyethyl group or a 2'-O methyl group. In still another preferred embodiment, the antisense nucleic acid is a chimeric nucleic acid. Preferably the chimeric nucleic acid comprises DNA residues linked together by phosphorothioate linkages, and the DNA residues are flanked on each side by at least one 2'-O methyl RNA residue linked together by a phosphorothioate linkage. More preferably the DNA residues are flanked on each side by at least three 2'-O methyl RNA residues. In yet another embodiment, the antisense nucleic acid is a ribozyme.

5

10

15

25

In a second aspect, the invention features a method of enhancing apoptosis in a cell, involving administering to the cell a negative regulator of the IAP-dependent antiapoptotic pathway. In preferred embodiments the negative regulator is an antisense IAP nucleic acid, an antibody that specifically binds an IAP polypeptide, an IAP polypeptide comprising a ring zinc finger, said polypeptide having no more than two BIR domains, a nucleic acid encoding the ring zinc finger domain of an IAP polypeptide, or a compound that prevents cleavage of the IAP polypeptide.

In preferred embodiments of the second aspect of the invention, the cell is in a mammal diagnosed with a proliferative disease, for example, cancer. The cell may comprises a mucosa-associated lymphoid tissue (MALT), a tissue in which the IAP gene HIAP1 is frequently involved in a translocation, resulting in marginal zone cell lymphomas. The cell may also be a breast cancer cell, where increased HIAP1 expression is known to correlate with tumor progression. The cell may also be a cell in which NFkB expression or activity is increased, for example, cell of head and

 \mathcal{M}

10

15

20

neck carcinomas, adult T-cell lymphomas, nasopharyngeal carcinomas, and Hodgkin's disease. The cell may also be an acute myelogenous leukemia cell, where increased XIAP levels correlate with poor patient prognosis. In addition, the cell may be a small cell lung carcinoma cell, where increased levels of XIAP correlates with increased resistance to radiation treatment.

In preferred embodiments of the second aspect of the invention, the IAP is XIAP, HIAP1, or HIAP2. Preferably the antisense nucleic acid is mammalian, for example, mouse or human. In still other preferred embodiments, the XIAP antisense is chosen from any one of SEQ ID NOS: 1 through 96, and the HIAP1 antisense is chosen from any one of SEQ ID NOS: 97 through 194.

In still other embodiments of the second aspect of the invention, the antisense nucleic acid comprises at least one modified internucleoside linkage. Preferably the modified internucleoside linkage is a phosphorothioate, a methylphosphonate, a phosphotriester, a phosphorodithioate, or a phosphoselenate linkage. In addition, the antisense nucleic acid may comprise at least one modified sugar moiety. Preferably this modified sugar moiety is a 2'-O methoxyethyl group or a 2'-O methyl group. In still another preferred embodiment, the antisense nucleic acid is a chimeric nucleic acid. Preferably the chimeric nucleicacid comprises DNA residues linked together by phosphorothioate linkages, and the DNA residues are flanked on each side by at least one 2'-O methyl RNA residue linked together by a phosphorothioate linkage. More preferably the DNA residues are flanked on each side by at least three 2'-O methyl RNA residues. In still further embodiments, administration of the antisense nucleic acid sensitizes the cell to chemotherapy or radiotherapy. In addition, the cell may be in vitro or in vivo.

4

In a third aspect, the invention features a pharmaceutical composition comprising a mammalian IAP antisense nucleic acid. In one preferred embodiment, the mammalian antisense IAP nucleic acid is a human antisense nucleic acid. Preferably the antisense nucleic acid binds a target sequence of the human XIAP gene or mRNA, the human HIAP1 gene or mRNA, the human HIAP2 gene or mRNA, the murine XIAP gene or mRNA, the murine HIAP1 gene or mRNA, or the murine HIAP2 gene or mRNA. More preferably the composition comprises an antisense nucleic acid chosen from any one of SEQ ID NOS: 1 through 96 (XIAP) or SEQ ID NOS: 97 through 194 (HIAP1).

10

15

20

In another aspect, the invention features an IAP gene nucleic acid fragment or antisense RNA sequence for use in suppressing cell proliferation. Such nucleic acids of the invention and methods for using them may be identified according to a method involving: (a) providing a cell sample; (b) introducing by transformation into the cell sample a candidate IAP nucleic acid; (c) expressing the candidate IAP nucleic acid within the cell sample; and (d) determining whether the cell sample exhibits an altered apoptotic response, whereby decreased apoptosis identifies an anti-proliferative compound. Preferably the cell is a cancer cell.

In another aspect, the invention features a method of treating a patient diagnosed with a proliferative disease. In the method, apoptosis may be induced in a cell to control a proliferative disease either alone or in combination with other therapies by administering to the cell a negative regulator of the IAP-dependent or anti-apoptotic pathway. The negative regulator may be, but is not limited to, an IAP ring zinc finger, and an IAP polypeptide that includes a ring zinc finger and lacks at least one BIR domain. Alternatively, apoptosis may be induced in the cell by administering a nucleic acid encoding an IAP antisense RNA molecule

administered directly or via gene therapy (see U.S. Pat. No. 5,576,208 for general parameters that may be applicable in the selection of IAP antisense RNAs). In yet another method, the negative regulator may be a purified antibody, or a fragment thereof, that binds specifically to an IAP polypeptide. For example, in one preferred embodiment, the antibody may bind to an approximately 26 kDa cleavage product of an IAP polypeptide that includes at least one BIR domain but lacks a ring zinc finger domain.

In two additional aspects, the invention features a transgenic animal and methods of using the mammal for detection of anti-cancer therapeutics. Preferably the mammal overexpresses an IAP polypeptide and/or expresses an IAP antisense RNA or IAP fragment. In one embodiment, the animal also has a genetic predisposition to cancer or has cancer cells under conditions that provide for proliferation absent the transgenic construct encoding either the antisense RNA or fragment.

10

15

20

25

"Protein" or "polypeptide" or "polypeptide fragment" means any chain of more than two amino acids, regardless of post-translational modification (e.g., glycosylation or phosphorylation), constituting all or part of a naturally-occurring polypeptide or peptide, or constituting a non-naturally occurring polypeptide or peptide.

"Apoptosis" means the process of cell death wherein a dying cell displays a set of well-characterized biochemical hallmarks that include cell membrane blebbing, cell soma shrinkage, chromatin condensation, and DNA laddering. Cells that die by apoptosis include neurons (e.g., during the course of neurodegenerative diseases such as stroke, Parkinson's disease, and Alzheimer's disease), cardiomyocytes (e.g., after myocardial infarction or over the course of congestive heart failure), and cancer cells (e.g., after exposure to radiation or chemotherapeutic agents). Environmental stress (e.g., hypoxic stress) that is not alleviated may cause

a cell to enter the early phase of the apoptotic pathway, which is reversible (i.e., cells at the early stage of the apoptotic pathway can be rescued). At a later phase of apoptosis (the commitment phase), cells cannot be rescued, and, as a result, are committed to die.

5

10

15

20

25

Proteins and compounds known to stimulate and inhibit apoptosis in a diverse variety of cells are well known in the art. For example, intracellular expression and activation of the caspase (ICE) family induces or stimulates apoptotic cell death, whereas expression of the IAPs or some members of the Bcl-2 family inhibits apoptotic cell death. In addition, there are survival factors that inhibit cell death in specific cell types. For example, neurotrophic factors, such as NGF inhibit neuronal apoptosis.

In some situations it may be desirable to artificially stimulate or inhibit apoptotic cell death by gene therapy or by a compound that mimics a gene therapeutic effect. For example, a cell that is susceptible to apoptosis induced by disease or environmental stress may be made more resistant to apoptosis by introducing an expression vector encoding an anti-apoptotic protein (such as an IAP, a Bcl-2 family member, or a neurotrophin) into the cell. Conversely, a cancer cell may be made less resistant to apoptosis by introducing into it an expression vector encoding a pro-apoptotic protein (such as a caspase) or by introducing into it an antisense nucleic acid, for example, an IAP antisense nucleic acid, regardless of its length. In addition, placement of the encoded protein of interest under the translational regulation of a XIAP IRES ensures that copious quantities of the protein are produced, especially under cellular conditions during which most protein translation (i.e., cap-dependent protein translation) is down-regulated, e.g., when a cell is under environmental stress, and when a cell is at a threshold for entering the apoptotic pathway.

By "IAP gene" is meant a gene encoding a polypeptide having at least one BIR domain and a ring zinc finger domain that is capable of modulating (inhibiting or enhancing) apoptosis in a cell or tissue when provided by other intracellular or extracellular delivery methods (see, e.g., U.S. Patent No. 5,919,912, U.S.S.N. 08/576,965, and PCT/IB96/01022). In preferred embodiments, the IAP gene is a gene having about 50% or greater nucleotide sequence identity to at least one of the IAP amino acid encoding sequences of Figs. 1 through 6, or portions thereof. Preferably the region of sequence over which identity is measured is a region encoding at least one BIR domain and a ring zinc finger domain.

Mammalian IAP genes include nucleotide sequences isolated from any mammalian source. Preferably the mammal is a human.

10

20

The term "IAP gene" is meant to encompass any member of the family of genes that encode inhibitors of apoptosis. An IAP gene may encode a polypeptide that has at least 20%, preferably at least 30%, and most preferably at least 50% amino acid sequence identity with at least one of the conserved regions of one of the IAP members described herein (i.e., either the BIR or ring zinc finger domains from human or murine XIAP, HIAP1, and HIAP2). Representative members of the IAP gene family include, without limitation, the human and murine XIAP, HIAP1, and HIAP2 genes.

By "IAP protein" or "IAP polypeptide" is meant a polypeptide, or fragment thereof, encoded by an IAP gene.

Ù,

Pro-Xaal-Cys-Xaal-Phe-Val, wherein Xaal is any amino acid and Xaa2 is any amino acid or is absent (SEQ ID NO: 216). Preferably the sequence is substantially identical to one of the BIR domain sequences provided for XIAP, HIAP1, or HIAP2 herein.

5

10

15

20

Preferably the sequence is substantially identical to the RZF domains provided in U.S.S.N. 08/800,929, incorporated herein by reference, for the human or murine XIAP, HIAP1, or HIAP2.

By "enhancing apoptosis" is meant increasing the number of cells that apoptose in a given cell population. Preferably the cell population is selected from a group including ovarian cancer cells, breast cancer cells, pancreatic cancer cells, T cells, neuronal cells, fibroblasts, or any other cell line known to proliferate in a laboratory setting. It will be appreciated that the degree of apoptosis enhancement provided by an apoptosisenhancing compound in a given assay will vary, but that one skilled in the art can determine the statistically significant change in the level of apoptosis that identifies a compound that enhances apoptosis otherwise limited by an IAP. Preferably "enhancing apoptosis" means that the increase in the number of cells undergoing apoptosis is at least 25%, more preferably the increase is 50%, and most preferably the increase is at least one-fold. Preferably the sample monitored is a sample of cells that normally undergo insufficient apoptosis (i.e., cancer cells). Methods for

detecting a changes in the level of apoptosis (i.e., enhancement or reduction) are described herein.

By "proliferative disease" is meant a disease that is caused by or results in inappropriately high levels of cell division, inappropriately low levels of apoptosis, or both. For example, cancers such as lymphoma, leukemia, melanoma, ovarian cancer, breast cancer, pancreatic cancer, and lung cancer are all examples of proliferative disease.

By "IAP biological activity" is meant any activity known to be caused *in vivo* or *in vitro* by an IAP polypeptide.

10

15

20

25

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) an IAP polypeptide.

By "transgene" is meant any piece of DNA that is inserted by artifice into a cell, and becomes part of the genome of the organism that develops from that cell. Such a transgene may include a gene that is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell that includes a DNA sequence that is inserted by artifice into a cell and becomes part of the genome of the organism that develops from that cell. As used herein, the transgenic organisms are generally transgenic mammals (e.g., rodents, such as rats or mice) and the DNA (transgene) is inserted by artifice into the nuclear genome.

By "transformation" is meant any method for introducing foreign molecules, for example, an antisense nucleic acid, into a cell. Lipofection, calcium phosphate precipitation, retroviral delivery, electroporation, biolistic transformation, and penetratin are just a few of the teachings that may be used. For example, biolistic transformation is a method for

introducing foreign molecules into a cell using velocity driven microprojectiles such as tungsten or gold particles. Such velocity-driven methods originate from pressure bursts that include, but are not limited to, helium-driven, air-driven, and gunpowder-driven techniques. Biolistic transformation may be applied to the transformation or transfection of a wide variety of cell types and intact tissues including, without limitation, intracellular organelles (e.g., and mitochondria and chloroplasts), bacteria, yeast, fungi, algae, animal tissue, and cultured cells. In another example, a foreign molecule (e.g., an antisense nucleic acid) can be translocated into a cell using the penetratin system as described, for example, by Prochiantz (Nature Biotechnology 16: 819-820, 1998; and Derossi et al. (Trends Cell Biol. 8: 84-87, 1998). In this system a penetratin peptide contains a transduction sequence that carries the peptide and a conjugated partner, for example, a phosphorothicate antisense nucleic acid (that is cross-linked through a disulfide bridge to the peptide) across the plasma membrane into the cell. The disulfide band is reduced inside the cell, releasing the partner.

10

15

20

25

By "antisense," as used herein in reference to nucleic acids, is meant a nucleic acid sequence, regardless of length, that is complementary to the coding strand or mRNA of an IAP gene. Preferably the antisense nucleic acid is capable of enhancing apoptosis when present in a cell that normally does not undergo sufficient apoptosis. Preferably the increase is at least 10%, relative to a control, more preferably 25%, and most preferably 1-fold or more. Preferably an IAP antisense nucleic acid comprises from about 8 to 30 nucleotides. An IAP antisense nucleic acid may also contain at least 40, 60, 85, 120, or more consecutive nucleotides that are complementary to a IAP mRNA or DNA, and may be as long as a full-length IAP gene or mRNA. The antisense nucleic acid may contain a modified backbone, for example, phosphorothioate, phosphorodithioate, or

other modified backbones known in the art, or may contain non-natural internucleoside linkages.

By "ribozyme" is meant an RNA that has enzymatic activity, possessing site specificity and cleavage capability for a target RNA molecule. Ribozymes can be used to decrease expression of a polypeptide. Methods for using ribozymes to decrease polypeptide expression are described, for example, by Turner et al., (Adv. Exp. Med. Biol. 465:303-318, 2000) and Norris et al., (Adv. Exp. Med. Biol. 465:293-301, 2000).

10

15

20

25

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 50%, preferably 85%, more preferably 90%, and most preferably 95% homology to a reference amino acid or nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

Sequence identity is typically measured using sequence analysis software with the default parameters specified therein (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). This software program matches similar sequences by assigning degrees of homology to various substitutions, deletions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine, valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

By "substantially pure polypeptide" is meant a polypeptide that has been separated from the components that naturally accompany it.

Typically, the polypeptide is substantially pure when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably the polypeptide is an IAP polypeptide that is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, pure. A substantially pure IAP polypeptide may be obtained, for example, by extraction from a natural source (e.g., a fibroblast, neuronal cell, or lymphocyte) by expression of a recombinant nucleic acid encoding an IAP polypeptide, or by chemically synthesizing the protein. Purity can be measured by any appropriate method, e.g., by column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

10

15

20

25

A protein is substantially free of naturally associated components when it is separated from those contaminants that accompany it in its natural state. Thus, a protein that is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. Accordingly, substantially pure polypeptides include those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

By "substantially pure DNA" is meant DNA that is free of the genes that, in the naturally-occurring genome of the organism from which the DNA of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other

sequences. It also includes a recombinant DNA that is part of a hybrid gene encoding additional polypeptide sequence.

By "positioned for expression" is meant that the DNA molecule is positioned adjacent to a DNA sequence, that directs transcription and translation of the sequence (i.e., facilitates the production of, e.g., an IAP polypeptide, a recombinant protein or an RNA molecule).

5

10

15

20

25

By "reporter gene" is meant a gene whose expression may be assayed; such genes include, without limitation, glucuronidase (GUS), luciferase, chloramphenical transacetylase (CAT), and Beta-galactosidase.

By "promoter" is meant a minimal sequence sufficient to direct transcription. Also included in the invention are those promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell type-specific, tissue-specific or that are inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the native gene.

By "operably linked" is meant that a gene and one or more regulatory sequences are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequences.

By "conserved region" is meant any stretch of six or more contiguous amino acids exhibiting at least 30%, preferably 50%, and most preferably 70% amino acid sequence identity between two or more of the IAP family members, (e.g., between human HIAP1, HIAP2, and XIAP). Examples of preferred conserved regions include, without limitation, BIR domains and ring zinc finger domains.

By "detectably-labelled" is meant any means for marking and identifying the presence of a molecule, e.g., an oligonucleotide probe or primer, a gene or fragment thereof, or a cDNA molecule. Methods for detectably-labelling a molecule are well known in the art and include,

without limitation, radioactive labelling (e.g., with an isotope such as ³²P or ³⁵S) and nonradioactive labelling (e.g., chemiluminescent labeling or fluorescein labelling).

By "purified antibody" is meant an antibody that is at least 60%, by weight, free from proteins and naturally occurring organic molecules with which it is naturally associated. Preferably the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, e.g., an IAP-specific antibody. A purified antibody may be obtained, for example, by affinity chromatography using recombinantly-produced protein or conserved motif peptides and standard techniques.

By "specifically binds" is meant an antibody that recognizes and binds a protein but that does not substantially recognize and bind other molecules in a sample, e.g., a biological sample, that naturally includes protein.

10

15

20

25

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

Fig. 1 is the human XIAP cDNA sequence (SEQ ID NO: 218) and the XIAP polypeptide sequence (SEQ ID NO: 219).

Fig. 2 is the human HIAP1 cDNA sequence (SEQ ID NO: 220) and the HIAP1 polypeptide sequence (SEQ ID NO: 221).

Fig. 3 is the human HIAP2 cDNA sequence (SEQ ID NO: 222) and the HIAP2 polypeptide sequence (SEQ ID NO: 223). The sequence absent in the HIAP2 variant is boxed.

Fig. 4 is the murine XIAP (also referred to as "miap-3") cDNA sequence (SEQ ID NO: 224) and encoded murine XIAP polypeptide sequence (SEQ ID NO: 225).

Fig. 5 is the murine HIAP1 (also referred to as "miap-1") cDNA sequence (SEQ ID NO: 226) and the encoded murine HIAP1 polypeptide sequence (SEQ ID NO: 227).

Fig. 6 is the murine HIAP2 (also referred to as "miap-2") cDNA sequence (SEQ ID NO: 228) and the encoded murine HIAP2 polypeptide (SEQ ID NO: 229).

5

10

20

25

Figs.7A through 7L are graphs showing the effect of antisense XIAP oligonucleotides on XIAP protein expression, relative to total protein (Figs. 7A, 7C, 7E, 7G, 7I, and 7K). Figs. 7B, 7D, 7F, 7H, 7J, and 7L are the total protein concentration values for each oligonucleotide transfection compared to mock transfection results that were used to normalize the above XIAP protein results.

Figs. 8A through 8C are graphs showing the effects of various antisense XIAP oligonucleotides, alone or in combination, on XIAP RNA (Fig. 8A) and protein (Fig. 8B). Fig. 8C is a graph of the total protein concentration values for each oligonucleotide transfection compared to mock transfection results, which were used to normalize the XIAP protein results shown in Fig. 8B.

Figs. 9A through 9D are graphs of the effects of antisense XIAP oligonucleotides on cell viability (Figs. 9A, 9C, and 9D), and chemosensitization in the presence of adriamycin (Fig. 9B).

Fig. 10 is a graph showing the effects of HIAP1 antisense oligonucleotides on HIAP1 RNA levels.

Fig. 11A is a densitometric scan of a Western blot showing the effects of HIAP1 antisense oligonucleotides on a cell's ability to block cycloheximide-induced upregulation of HIAP1 protein.

Fig. 11 is a graph showing the effects of HIAP1 antisense oligonucleotides on a cell's ability to block cycloheximide-induced upregulation of HIAP1 protein.

Fig. 12 is a graph showing the effects of HIAP1 antisense oligonucleotides on cytotoxicity, as measured by total protein.

Fig. 13 is a graph showing the validation of the sequence specificity for HIAP1 antisense oligonucleotide APO 2.

Fig. 14 is a graph showing the effect of HIAP1 antisense oligonucleotides on the chemosensitization of drug-resistant SF295 glioblastomas.

5

10

15

20

25

Fig. 15 is the human XIAP sequence containing a 5' UTR, the coding region, and a 3' UTR (SEQ ID NO: 230).

Fig. 16 is the human HIAP1 sequence containing a 5' UTR, the coding region, and a 3' UTR (SEQ ID NO: 231).

Detailed Description of the Invention

The present invention provides IAP antisense nucleic acid sequences that inhibit IAP biological activity, regardless of length, and methods for using them to induce apoptosis in a cell. The antisense nucleic acids of the present invention may also be used to form pharmaceutical compositions. The invention also features methods for enhancing apoptosis in a cell by administering a negative regulator of the IAP anti-apoptotic pathway other than antisense. Such negative regulators include, for example, an IAP polypeptide comprising a ring zinc finger having no more than two BIR domains, and a compound that prevents cleavage of an IAP polypeptide. Such negative regulators may also be used to form a pharmaceutical composition. These pharmaceutical compositions may be used to treat, ameliorate, improve, sustain, or prevent a proliferative disease, for example, cancer, or a symptom of a proliferative disease.

Administration

10

15

20

25

An IAP antisense nucleic acid, or other negative regulator of the IAP anti-apoptotic pathway may be administered within a pharmaceutically-acceptable diluent, carrier, or excipient, in unit dosage form. Conventional pharmaceutical practice may be employed to provide suitable formulations or compositions to administer the compounds to patients suffering from a disease that is caused by excessive cell proliferation. Administration may begin before the patient is symptomatic. Any appropriate route of administration may be employed, for example, administration may be parenteral, intravenous, intraarterial, subcutaneous, intramuscular, intracranial, intraorbital, ophthalmic, intraventricular, intracapsular, intraspinal, intracisternal, intraperitoneal, intranasal, aerosol, suppository, or oral administration. For example, therapeutic formulations may be in the form of liquid solutions or suspensions; for oral administration, formulations may be in the form of tablets or capsules; and for intranasal formulations, in the form of powders, nasal drops, or aerosols.

Methods well known in the art for making formulations are found, for example, in "Remington's Pharmaceutical Sciences." Formulations for parenteral administration may, for example, contain excipients, sterile water, or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated napthalenes. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for IAP modulatory compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example,

polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

The formulations can be administered to human patients in therapeutically effective amounts (e.g., amounts which prevent, eliminate, or reduce a pathological condition) to provide therapy for a disease or condition. The preferred dosage of therapeutic agent to be administered is likely to depend on such variables as the type and extent of the disorder, the overall health status of the particular patient, the formulation of the compound excipients, and its route of administration.

If desired, treatment with an IAP antisense nucleic acid, IAP fragments, or other negative regulator of the anti-apoptotic pathway may be combined with more traditional therapies for the proliferative disease such as surgery or chemotherapy.

10

15

20

25

For any of the methods of application described above, the therapeutic antisense IAP nucleic acid or other negative regulator of the IAP anti-apoptotic pathway is preferably applied to the site of the needed apoptosis event (for example, by injection). However, it may also be applied to tissue in the vicinity of the predicted apoptosis event or to a blood vessel supplying the cells predicted to require enhanced apoptosis.

The dosage of an antisense IAP nucleic acid, or a negative regulator of the IAP anti-apoptotic pathway, for example, an IAP fragment, IAP mutant protein or an IAP antibody depends on a number of factors, including the size and health of the individual patient, but, generally, between 0.1 mg and 100 mg inclusive are administered per day to an adult in any pharmaceutically acceptable formulation. In addition, treatment by any IAP-modulating gene therapy approach may be combined with more traditional therapies.

Antisense Therapy

10

15

20

25

Anti-cancer therapy may be accomplished by direct administration of a therapeutic antisense IAP nucleic acid to a cell that is expected to require enhanced apoptosis. The antisense nucleic acid may be produced and isolated by any one of many standard techniques. Administration of IAP antisense nucleic acids to malignant cells can be carried out by any of the methods for direct nucleic acid administration, as described herein.

Retroviral vectors, adenoviral vectors, adeno-associated viral vectors, or other viral vectors with the appropriate tropism for cells likely requiring enhanced apoptosis (for example, breast cancer and ovarian cancer cells) may be used as a gene transfer delivery system for a therapeutic antisense IAP gene construct. Numerous vectors useful for this purpose are generally known (Miller, Human Gene Therapy 15-14, 1990; Friedman, Science 244:1275-1281, 1989; Eglitis and Anderson, BioTechniques 6:608-614, 1988; Tolstoshev and Anderson, Current Opinion in Biotechnology 1:55-61, 1990; Sharp, The Lancet 337:1277-1278, 1991; Cornetta et al., Nucleic Acid Research and Molecular Biology 36:311-322, 1987; Anderson, Science 226:401-409, 1984; Moen, Blood Cells 17:407-416, 1991; Miller et al., BioTechniques 7:980-990, 1989; Le Gal La Salle et al., Science 259:988-990, 1993; and Johnson, Chest 107:77S-83S, 1995).

Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg et al., N. Engl. J. Med 323:370, 1990; Anderson et al., U.S. Patent No. 5,399,346). Non-viral approaches may also be employed for the introduction of therapeutic DNA into cells otherwise predicted to undergo apoptosis. For example, IAPs may be introduced into a cell by lipofection (Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413, 1987; Ono et al., Neurosci. Lett. 117:259, 1990; Brigham et al., Am. J. Med. Sci. 298:278, 1989; Staubinger et al., Meth. Enz. 101:512,

1983), the penetratin system (Allinquant et al., J. Cell Biol. 128:919-927, 1995; Prochiantz, Curr. Opin. Neurobiol. 6:629-634, 1996), asialorosonucoid-polylysine conjugation (Wu et al., J. Biol. Chem. 263:14621, 1988; Wu et al., J. Biol. Chem. 264:16985, 1989); or, less preferably microinjection under surgical conditions (Wolff et al., Science 247:1465, 1990).

In the therapeutic nucleic acid constructs described, nucleic acid expression can be directed from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element. For example, if desired, enhancers known to preferentially direct gene expression in ovarian cells, breast tissue, neural cells, T cells, or B cells may be used to direct expression. Enhancers include, without limitation, those that are characterized as tissue- or cell-specific in their expression. Alternatively, if a clone is used as a therapeutic construct, regulation may be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

20 Therapeutic Products

25

For IAP related therapies one may employ the paradigms utilized for Bcl-2 and Ras antisense development, although accommodation of an IAP mutation is not required (in contrast to Ras antisense). Most useful are antisense constructs that enhance apoptosis at least 10%, preferably by enhancing degradation of the RNA in the nucleus.

Manipulation of cancer chemotherapeutic drug resistance using an antisense oligonucleotide and fragment approaches

We have documented that overexpression of the IAPs renders cell lines resistant to serum growth factor withdrawal, tumor necrosis factor alpha (TNF) and menadione exposure, all of which are treatments that normally induce apoptosis. Herein, we describe the extension of these studies to cancer cell lines using apoptotic triggers used in clinical situations, such as doxorubicin, adriamycin, and methotrexate. Our findings have led up to the design of antisense RNA therapeutics. Rapid screening of multiple cell lines for apoptotic response has been made 10 feasible through the generation of a series of sense and antisense adenoviral IAP and expression vectors, as well as control lacZ viruses. One may now show enhanced drug resistance using the expression constructs. In addition, anti-sense adenovirus constructs may be developed and used to test reversal of the drug resistant phenotype of 15 appropriate cell lines. We have designed a series of antisense oligonucleotides to various regions of each of the iaps. These oligonucleotides may be used to enhance drug sensitivity after testing in an assay system, i.e., with the adenoviral vectors system. Animal modeling of the effectiveness of antisense IAP oligonucleotides may also 20 be employed as a step in testing and appropriate transgenic mammals for this are described in U.S.S.N. 08/800,929, incorporated herein by reference, and are also generally available in the art.

Characterization of IAP Activity and Intracellular Localization Studies

25

The ability of IAPs to modulate apoptosis can be defined *in vitro* systems in which alterations of apoptosis can be detected. Mammalian expression constructs carrying IAP cDNAs, which are either full-length, truncated, or antisense constructs can be introduced into cell lines, such as

CHO, NIH 3T3, HL60, Rat-1, or Jurkat cells. In addition, SF21 insect cells may be used, in which case the IAP gene is preferentially expressed using an insect heat shock promoter. Following transfection, apoptosis can be induced by standard methods, which include serum withdrawal, or application of staurosporine, menadione (which induces apoptosis via free radial formation), or anti-Fas antibodies. As a control, cells are cultured under the same conditions as those induced to undergo apoptosis, but either not transfected, or transfected with a vector that lacks an IAP insert. The ability of each IAP related construct to inhibit or enhance apoptosis upon expression can be quantified by calculating the survival index of the cells, i.e., the ratio of surviving transfected cells to surviving control cells. These experiments can confirm the presence of apoptosis inhibiting activity and, as discussed below, can also be used to determine the functional region(s) of an IAP that may be employed to achieve enhancement of apoptosis. These assays may also be performed in combination with the application of additional compounds in order to identify compounds that enhance apoptosis via IAP expression.

Apoptosis Assays

5

15

20

25

Specific examples of apoptosis assays are provided in the following references. Assays for apoptosis in lymphocytes are disclosed by: Li et al., "Induction of apoptosis in uninfected lymphocytes by HIV-1 Tat protein", Science 268:429-431, 1995; Gibellini et al., "Tat-expressing Jurkat cells show an increased resistance to different apoptotic stimuli, including acute human immunodeficiency virus-type 1 (HIV-1) infection", Br. J. Haematol. 89:24-33, 1995; Martin et al., "HIV-1 infection of human CD4⁺ T cells *in vitro*. Differential induction of apoptosis in these cells." J. Immunol. 152:330-342, 1994; Terai et al., "Apoptosis as a mechanism of cell death in cultured T lymphoblasts acutely infected with HIV-1", J.

Clin. Invest. 87:1710-1715, 1991; Dhein et al., "Autocrine T-cell suicide mediated by APO-l/(Fas/CD95)", Nature 373:438-441, 1995; Katsikis et al., "Fas antigen stimulation induces marked apoptosis of T lymphocytes in human immunodeficiency virus-infected individuals", J. Exp. Med. 1815:2029-2036, 1995; Westendorp et al., "Sensitization of T cells to CD95-mediated apoptosis by HIV-1 Tat and gp120", Nature 375:497, 1995; and DeRossi et al., Virology 198:234-44, 1994.

Assays for apoptosis in fibroblasts are disclosed by: Vossbeck et al., "Direct transforming activity of TGF-beta on rat fibroblasts", Int. J. Cancer 61:92-97, 1995; Goruppi et al., "Dissection of c-myc domains involved in S phase induction of NIH3T3 fibroblasts", Oncogene 9:1537-1544, 1994; Fernandez et al., "Differential sensitivity of normal and Haras transformed C3H mouse embryo fibroblasts to tumor necrosis factor: induction of bcl-2, c-myc, and manganese superoxide dismutase in resistant cells", Oncogene 9:2009-2017, 1994; Harrington et al., "c-Myc-induced apoptosis in fibroblasts is inhibited by specific cytokines", EMBO J., 13:3286-3295, 1994; and Itoh et al., "A novel protein domain required for apoptosis. Mutational analysis of human Fas antigen", J. Biol. Chem. 268:10932-10937, 1993.

10

15

20

25

Assays for apoptosis in neuronal cells are disclosed by: Melino et al., "Tissue transglutaminase and apoptosis: sense and antisense transfection studies with human neuroblastoma cells", Mol. Cell. Biol. 14:6584-6596, 1994; Rosenbaum et al., "Evidence for hypoxia-induced, programmed cell death of cultured neurons", Ann. Neurol. 36:864-870, 1994; Sato et al., "Neuronal differentiation of PC12 cells as a result of prevention of cell death by bcl-2", J. Neurobiol. 25:1227-1234, 1994; Ferrari et al., "N-acetylcysteine D- and L-stereoisomers prevents apoptotic death of neuronal cells", J. Neurosci. 1516:2857-2866, 1995; Talley et al., "Tumor necrosis factor alpha-induced apoptosis in human neuronal cells:

protection by the antioxidant N-acetylcysteine and the genes bcl-2 and crmA", Mol. Cell Biol. 1585:2359-2366, 1995; and Walkinshaw et al., "Induction of apoptosis in catecholaminergic PC12 cells by L-DOPA. Implications for the treatment of Parkinson's disease.", J. Clin. Invest. 95:2458-2464, 1995.

Assays for apoptosis in insect cells are disclosed by: Clem et al., "Prevention of apoptosis by a baculovirus gene during infection of insect cells", Science 254:1388-1390, 1991; Crook et al., "An apoptosis-inhibiting baculovirus gene with a zinc finger-like motif", J. Virol. 67:2168-2174, 1993; Rabizadeh et al., "Expression of the baculovirus p35 gene inhibits mammalian neural cell death", J. Neurochem. 61:2318-2321, 1993; Birnbaum et al., "An apoptosis inhibiting gene from a nuclear polyhedrosis virus encoding a polypeptide with Cys/His sequence motifs", J. Virol. 68:2521-2528, 1994; and Clem et al., "Control of programmed cell death by the baculovirus genes p35 and IAP", Mol. Cell. Biol. 14:5212-5222, 1994.

The following examples are to illustrate the invention. They are not meant to limit the invention in any way.

20

25

15

5

. 10

Example 1: Testing of antisense oligonucleotides

1. Complete panel of adenovirus constructs. The panel may consist of approximately four types of recombinant virus. A) Sense orientation viruses for each of the IAP open reading frames. These viruses are designed to massively overexpress the recombinant protein in infected cells. XIAP, HIAP1, HIAP2, and NAIP. B) Antisense orientation viruses in which the viral promoter drives the synthesis of an mRNA of opposite polarity to the *iap* mRNA, thereby shutting off host cell synthesis of the

targeted protein coding region. XIAP, HIAP1, HIAP2, and NAIP
"antisense" constructs are used for production of such antisense IAPs. C)
Sub-domain expression viruses. These constructs express only a partial
IAP protein in infected cells. We have data indicating that deletion of the
zinc finger of XIAP renders the protein more potent in protecting cell
against apoptotic triggers. This data also indicates that expression of the
zinc finger alone will indicate apoptosis by functioning as a dominantnegative repressor of XIAP function. XIAP- ZF and XIAP- BIR viruses
are required. D) Control viruses. Functional analysis of the IAPs requires
suitable positive and negative controls for comparison. Bcl-2 sense, Bcl-2
antisense, p53 sense, and Lac Z (negative control) viruses may be utilized.

- Confirmation of recombinant adenovirus function. Verification of the sense adenovirus function involves infection of tissue culture cells and determination of protein expression levels. We have performed Western blot analysis of several of the recombinant adenoviruses, including NAIP, XIAP and XIAP- ZF. The remaining viruses may be readily assessed for protein expression using the polyclonal IAP antibodies. Functional analysis of the antisense viruses may be done at the RNA level using either Northern blots of total RNA harvested from infected tissue culture cells or ribonuclease protection assays. Western blot analysis of infected cells will be used to determine whether the expressed antisense RNA interferes with IAP expression in the host cell.
- 25 3. Documentation that IAP overexpression results in increased drug resistance. We have optimized cell death assays to allow high through-put of samples with minimal sample variation. Testing of the sense IAP adenoviruses for their ability to alter drug sensitivity of breast and pancreatic adenocarcinoma cell lines may be accomplished as follows.

Cancer cell lines are infected with the recombinant viruses, cultured for 5 days, then subdivided into 24 well plates. Triplicate cell samples each receive increasing concentrations of the anti-cancer drug under investigation. Samples are harvested at 24, 48, and 72 hours post-exposure, and assayed for the number of viable cells in the well. The dose response curve is then compared to uninfected and control virus (both positive and negative) infected cells. One may document a dramatic increase in the relative resistance of the cancer cell lines when infected with the sense viruses, confirming our hypothesis that overexpression of the IAP proteins contributes to the anti-apoptotic phenotype of cancer cells. Initial experiments utilize the drugs doxorubicin and adriamycin.

4. Documentation that antisense IAP overexpression results in increased drug sensitivity. Having confirmed that IAP overexpression renders cancer cells more resistant to chemotherapeutic drugs, one may examine whether the antisense adenoviruses render the same cells more sensitive. The effectiveness of antisense IAP viruses relative to antisense Bcl-2 virus will also be assessed as a crucial milestone.

15

5. Identification of antisense oligonucleotides. Concomitant to the adenovirus work, we have designed a series of antisense oligonucleotides to various regions of each of the IAPs. A generally accepted model of how antisense oligonucleotides function proposes that the formation of RNA/DNA duplexes in the nucleus activates cellular RnaseH enzymes which then enzymatically degrade the mRNA component of the hybrid. Virtually any region of the mRNA can be targeted, and therefore choosing an appropriate sequence to target is somewhat empirical.

6. Optimization of oligonucleotides. A secondary round of oligonucleotides may be made when effective target regions have been identified. These oligonucleotides target sequences in the immediate vicinity of the most active antisense oligonucleotides identified using methods such as those provided above. A second round of testing by Northern blot analysis may be required.

- 7. Testing antisense oligonucleotides in vitro. Following successful identification and optimization of targeting oligonucleotides, one may test these in the tissue culture model system using the optimal cell lines such 10 as those described in the cancer survey described in U.S.S.N. 08/800,929, incorporated herein by reference. Experimental procedures may parallel those used in the recombinant antisense adenovirus work. Negative control oligonucleotides with miss-match sequences are used to establish base line or non-specific effects. Assisted transfection of the 15 oligonucleotides using cationic lipid carriers may be compared to unassisted transfection. Confirmation of the effectiveness of specific antisense oligonucleotides prompts synthesis of oligonucleotides with modified phosphodiester linkages, such as phosphorothioate or methylimino substituted oligonucleotides. These may also be tested in 20 vitro.
 - 8. Animal modeling of antisense oligonucleotide therapies. Animal modeling of the effectiveness of the antisense IAP approach is described here. Cell lines are routinely assessed for their tumorigenic potential in "nude" mice, a hairless strain of mouse that is immunocompromised, and thus extremely susceptible to developing tumors. In the nude mouse assay, cancer cells are grown in tissue culture and then injected under the skin at multiple sites. The frequency with which these cells give rise to

25

palpable tumors within a defined period of time provides an index of the tumorigenic potential of the cell line in the absence of interference by a functional immune system. Preliminary assessment of an antisense IAP therapeutic involves injection of cancer cells infected with the recombinant adenoviruses (sense, antisense, and control viruses) under the skin, and the tumorigenic index compared to that of untreated cells. One may also use this model to assess the effectiveness of systemic administration of antisense oligonucleotides in increasing the efficacy of anti-cancer drugs in the nude mouse model. Phosphorothioate or methylimino substituted oligonucleotides will be assessed at this stage. This type of antisense oligo has demonstrated enhanced cell permeability and slower clearance rates from the body in experimental animal models.

Example 2: Antisense oligonucleotide (ODN) selection

15

20

25

We selected 96 or 98, mostly non-overlapping, 19-mer antisense oligonucleotide (ODN) sequences for XIAP and HIAP1, respectively, based on the selection criteria listed below. In the case of XIAP, we selected 96 sequences (each being 19 nucleobases in length) (SEQ ID NOS: 1 through 96; Table 1), from a region approximately 1 kb upstream of the start codon to approximately 1 kb downstream of the stop codon of the cDNA sequence (Fig. 15). This blanketed approximately 50% of the coding region, and immediate 5' and 3' UTR sequences (i.e., 96 19-mers span 1.8 kb of sequence, while the targeted region is approximately 3.5 kb in length, comprised of a coding region of 1.5 kb plus 1 kb at either side of UTR sequences).

Table 1. XIAP Antisense Oligonucleotides

SEQ		Position in XIAP	Antisense Oligonucleotide
ID NO:	Code	Sequence	Sequence
1	A1	2	AAAATTCTAAGTACCTGCA
2	B1	21	TCTAGAGGGTGGCTCAGGA
3	C1	44	CAGATATATATGTAACACT
4	D1	78	TGAGAGCCCTTTTTTTGTT
5	E1	110	AGTATGAAATATTTCTGAT
6	F1	134	ATTGGTTCCAATGTGTTCT
7	G1	160	TTAGCAAAATATGTTTTAA
8	H1	185	TGAATTAATTTTTAATATC
9	A2	238	ATTCAAGGCATCAAAGTTG
10	B2	326	GTCAAATCATTAATTAGGA
11	C2	370	AATATGTAAACTGTGATGC
12	D2	411	GCAGAATAAAACTAATAAT
13	E2	430	GAAAGTAATATTTAAGCAG
14	F2	488	TTACCACATCATTCAAGTC
15	G2	508	CTAAATACTAGAGTTCGAC
16	H2	535	ACACGACCGCTAAGAAACA
17	A3	561	TATCCACTTATGACATAAA
18	В3	580	GTTATAGGAGCTAACAAAT
19	C3	607	AATGTGAAACACAAGCAAC
20	D3	638	ACATTATATTAGGAAATCC
21	E3	653	CTTGTCCACCTTTTCTAAA
22	F3	673	ATCTTCTCTTGAAAATAGG
23	G3	694	CCTTCAAAACTGTTAAAAG
24	Н3	721	ATGTCTGCAGGTACACAAG
25	A4	759	ATCTATTAAACTCTTCTAC

SEQ		Position in XIAP	Antisense Oligonucleotide
ID NO:	Code	Sequence	Sequence
26	B4	796	ACAGGACTACCACTTGGAA
27	C4	815	TGCCAGTGTTGATGCTGAA
28	D4	835	GTATAAAGAAACCCTGCTC
29	E4	856	CGCACGGTATCTCCTTCAC
30	F4	882	CTACAGCTGCATGACAACT
31	G4	907	GCTGAGTCTCCATATTGCC
32	H4	930	ATACITTCCTGTGTCTTCC
33	A5	950	GATAAATCTGCAATTTGGG
34	B5	990	TTGTAGACTGCGTGGCACT
35	C5	1010	ACCATTCTGGATACCAGAA
36	D5	1029	AGTTTTCAACTTTGTACTG
37	E5	1059	ATGATCTCTGCTTCCCAGA
38	F5	1079	AGATGGCCTGTCTAAGGCA
39	G5	1100	AGTTCTCAAAAGATAGTCT
40	H5	1126	GTGTCTGATATATCTACAA
41	A6	1137	TCGGGTATATGGTGTCTGA
42	B6	1146	CAGGGTTCCTCGGGTATAT
43	C6	1165	GCTTCTTCACAATACATGG
44	D6	1192	GGCCAGTTCTGAAAGGACT
45	E6	1225	GCTAACTCTCTTGGGGTTA
46	F6	1246	GTGTAGTAGAGTCCAGCAC
47	Ģ6	1273	AAGCACTGCACTTGGTCAC
48	H6	1294	TTCAGTTTTCCACCACAAC
49	A7	1316	ACGATCACAAGGTTCCCAA
50	B7	1337	TCGCCTGTGTTCTGACCAG
51	C7	1370	CCGGCCCAAAACAAAGAAG

SEQ		Position in XIAP	Antisense Oligonucleotide
ID NO:	Code	Sequence	Sequence
52	D7	1393	GATTCACTTCGAATATTAA
53	E7	1413	TATCAGAACTCACAGCATC
54	F7	1441	GGAAGATTTGTTGAATTTG
55	G7	1462	TCTGCCATGGATGGATTTC
56	H7	1485	AAGTAAAGATCCGTGCTTC
57	A8	1506	CTGAGTATATCCATGTCCC
58	В8	1525	GCAAGCTGCTCCTTGTTAA
59	C8	1546	AAAGCATAAAATCCAGCTC
60	D8	1575	GAAAGCACTTTACTTTATC
61	Н8	1610	ACTGGGCTTCCAATCAGTT
62	E8	1629	GITGITCCCAAGGGTCTTC
63	F8	1650	ACCCTGGATACCATTTAGC
64	G8	1669	TGTTCTAACAGATATTTGC
65	A9	1688	TATATATTCTTGTCCCTTC
66	B9	1696	AGTTAAATGAATATTGTTT
67	C9	1725	GACACTCCTCAAGTGAATG
68	D9	1745	TTTCTCAGTAGTTCTTACC
69	E9	1759	GTTAGTGATGGTGTTTTCT
70	F9	1782	AGATGGTATCATCAATTCT
71	G9	1801	TGTACCATAGGATTTTGGA
72	Н9	1820	CCCCATTCGTATAGCTTCT
73	A10	1849	ATTATTTTCTTAATGTCCT
74	B10	1893	CAAGTGATTTATAGTTGCT
75	C10	1913	TAGATCTGCAACCAGAACC
76	D10	1945	CATCTTGCATACTGTCTTT
77	E10	1997	CCTTAGCTGCTCTTCAGTA

SEQ		Position in XIAP	Antisense Oligonucleotide
ID NO:	Code	Sequence	Sequence
78	F10	2018	AAGCTTCTCCTCTTGCAGG
79	G10	2044	ATATTTCTATCCATACAGA
80	H10	2076	CTAGATGTCCACAAGGAAC
81	A11	2096	AGCACATTGTTTACAAGTG
82	B11	2123	AGCACATGGGACACTTGTC
83	C11	2144	CTTGAAAGTAATGACTGTG
84	D11	2182	CCTACTATAGAGTTAGATT
85	E11	2215	ATTCAATCAGGGTAATAAG
86	F11	2234	AAGTCAGTTCACATCACAC
87	G11	2375	CAGTAAAAAAAATGGATAA
88	H11	2428	TTCAGTTATAGTATGATGC
89	A12	2471	TACACTTAGAAATTAAATC
90	B12	2630	TCTCTATCTTTCCACCAGC
91	C12	2667	AGAATCCTAAAACACAACA
92	D12	2709	ATTCGCACAAGTACGTGTT
93	E12	2785	TGTCAGTACATGTTGGCTC
94	F12	2840	ACATAGTGTTTTGCCACTT
95	G12	2861	CTTTGATCTGGCTCAGACT
96	H12	2932	GAAACCACATTTAACAGTT

Note that the three most 5' and the three most 3' nucleobases may comprise DNA residues, or RNA residues, such as 2'-O methyl RNA residues. For example, the antisense oligonucleotide sequence of SEQ ID NO: 3 may be CAGATATATATGTAACACT or CAGATATATGTAACACU.

A similar approach was taken for designing antisense oligonucleotides against HIAP1. Ninety-eight 19-mer sequences were chosen, with some of the latter sequences picked using less stringent criteria than the originally defined selection criteria (listed below), to increase the number of candidate sequences to study (SEQ ID NOS: 97 through 194; Table 2). Of these 98 sequences targeted to the HIAP1 sequence of Fig. 16, 15 (SEQ ID NOS: 97 through 104, 107, 113, 136, 156, 157, 181, and 193) were selected to evaluate the efficacy of decreasing HIAP1 expression. These 15 candidate sequences consisted of 4 sequences targeting the coding region (SEQ ID NOS: 136, 156, 157, and 181), 1 sequence targeting the 3' UTR (SEQ ID NO: 193), and 7 sequences targeting the 5'UTR (SEQ ID NOS: 100 through 104, 107, and 113; one of the 7 oligonucleotides overlapped the start codon), and 3 other oligonucleotides (SEQ IDs 97 through 99) that were designed to target an intronic segment of the 5'UTR (the value of which is discussed in Example 7). These above-described 15 HIAP1 antisense oligonucleotides were synthesized and tested.

5

Table 2. HIAP1 Antisense Oligonucleotides

		Position in HIAP1	Antisense Oligonucleotide
SEQ ID NO	Code	Sequence	Sequence
97	APO 1	1152	TCATTTGAGCCTGGGAGGU
98	APO 2	1172	CGGAGGCTGAGGCAGGAGA
99	APO 3	1207	GGTGTGGTGGTACGCGCCT
100	APO 4	1664	ACCCATGCACAAAACTACC
101	APO 5	1865	AGAATGTGCCAGTAGGAGA
102	APO 6	2440	TCTCACAGACGTTGGGCTT
103	APO 7	2469	CCAGTGGTTTGCAAGCATG
104	APO 8	3695	GAAATTTAGTGGCCAGGAA
105		4013	AGAAATACACAATTGCACC
106		4032	TACTGATACATTTTAAGGA
107	APO 9	4057	TTCAACATGGAGATTCTAA
108		4076	ATTTCTATGCATTTAGAGT
109		4121	AATACTAGGCTGAAAAGCC
110		4142	GGCTTTGCTTTTATCAGTT
111		4165	TCTAGGGAGGTAGTTTTGT

		Position in HIAP1	Antisense Oligonucleotide
SEQ ID NO	Code	Sequence	Sequence
112	- 0000	4189	GGGAAGAAAGGGACTAGC
113	APO 10	4212	GTTCATAATGAAATGAATG
114	ALO IV	4233	ATAAGAATATGCTGTTTTC
115		4265	TTCAAACGTGTTGGCGCTT
116	 	4283	ATGACAAGTCGTATTTCAG
117	l	4317	AAGTGGAATACGTAGACAT
118		4338	AGACAGGAACCCCAGCAGG
119		4357	CGAGCAAGACTCCTTTCTG
120		4376	AGTGTAATAGAAACCAGCA
121	 	4395	TGACCTTGTCATTCACACC
122	 	4426	TTATCCAGCATCAGGCCAC
123	 	4445	ACTGTCTCCTCTTTTCCAG
124		4464	TTTTATGCTTTTCAGTAGG
	+	4489	ACGAATCTGCAGCTAGGAT
125			CAAGTTGTTAACGGAATTT
126		4517	
127		4536	TAGGCTGAGAGGTAGCTTC
128	 	4555	GTTACTGAAGAAGGAAAAG
129		4574	GAATGAGTGTGTGGAATGT
130		4593	TGTTTTCTGTACCCGGAAG
131		4612	GAGCCACGGAAATATCCAC
132		4631	TGATGGAGAGTTTGAATAA
133	ļ	4656	GATTTGCTCTGGAGTTTAC
134		4670	GGCAGAAAATTCTTGATTT
135		4696	GGACAGGGGTAGGAACTTC
136	APO 11	4714	GCATTTTCGTTATTCATTG
137		4733	CTGAAAAGTAAGTAATCTG
138		4759	GGCGACAGAAAAGTCAATG
139		4812	CCACTCTGTCTCCAGGTCC
140		4831	CCACCACAGGCAAAGCAAG
141	ļ	4855	TTCGGTTCCCAATTGCTCA
142		4874	TTCTGACATAGCATTATCC
143		4893	TGGGAAAATGTCTCAGGTG
144		4907	TATAAATGGGCATTTGGGA
145	ļ	4926	TGTCTTGAAGCTGATTTTC
146		4945	GAAACTGTGTATCTTGAAG
147		4964	TGTCTGCATGCTCAGATTA
148		4988	GAATGTTTTAAAGCGGGCT
149		5007	CACTAGAGGGCCAGTTAAA
150		5040	CCGCACTTGCAAGCTGCTC
151		5070	CATCATCACTGTTACCCAC
152		5095	CCACCATCACAGCAAAAGC
153		5117	TCCAGATTCCCAACACCTG
154		5130	CCCATGGATCATCTCCAGA
155		5149	AACCACTTGGCATGTTGAA
156	APO 12	5168	CAAGTACTCACACCTTGGA
157	APO 13	5187	CCTGTCCTTTAATTCTTAT
158		5206	TGAACTTGACGGATGAACT
159	1	5225	TAGATGAGGGTAACTGGCT
160	ļ · · · · · · · · · · · · · · · · · · ·	5244	TGGATAGCAGCTGTTCAAG
161	1	5271	CATTTTCATCTCCTGGGCT

	T	Position in HIAP1	Antisense Oligonucleotide
SEQ ID NO	Code	Sequence	Sequence
162		529	TGGATAATTGATGACTCTG
163		5309	GTCTTCTCCAGGTTCAAAA
164		5337	TATTCATCATGATTGCATC
165		5366	CATTTCCACGGCAGCATTA
166		5367	CCAGGCTTCTACTAAAGCC
167		5416	GCTAGGATTTTTCTCTGAA
168		5435	TCTATAATTCTCTCCAGTT
169		5454	ACACAAGATCATTGACTAG
170		5473	TCTGCATTGAGTAAGTCTA
171		5492	CTCTTCCCTTATTTCATCT
172		5515	TCCTCAGTTGCTCTTTCTC
173		5560	GCCATTCTATTCTTCCGGA
174		5579	AGTCAAATGTTGAAAAAGT
175		5598	CCAGGATTGGAATTACACA
176		5622	ATTCCGGCAGTTAGTAGAC
177		5646	TAACATCATGTTCTTGTTC
178		5675	GTCTGTGTCTTCTGTTTAA
179		5684	TTCTCTTGCTTGTAAAGAC
180		5703	CTAAAATCGTATCAATCAG
181	APO 14	5723	GGCTGCAATATTTCCTTTT
182		5742	GAGAGTTTCTGAATACAGT
183		5761	ACAGCTTCAGCTTCTTGCA
184		5780	AAATAAATGCTCATATAAC
185		5821	GAAACATCTTCTGTGGGAA
186		5841	GTTCTTCCACTGGTAGATC
187		5862	CTTCTTGTAGTCTCCGCAA
188		5890	TTGTCCATACACACTTTAC
189		6097	AACCAAATTAGGATAAAA
	_ }	<u></u>	G
190		6181	ATGTTCATATGGTTTAGAT
191		6306	TAAGTTTTACTTCACTTAC
192		6369	ATGTTCCCGGTATTAGTAC
193	APO 15	6432	GGGCTCAAGTAATTCTCTT
194		6455	GCCCAGGATGGATTCAAAC

Oligonucleotide selection criteria

The computer program OLIGO (previously distributed by National Biosciences Inc.) was used to define suitable antisense oligonucleotides based on the following criteria: 1) no more than 75% GC content, and no more than 75% AT content; 2) preferably no oligonucleotide with 4 or more consecutive G residues (due to reported toxic effects, although one was chosen as a toxicity control); 3) no oligonucleotides with the ability

to form stable dimers or hairpin structures; and 4) sequences around the translation start site are a preferred region. In addition, accessible regions of the mRNA were predicted with the help of the RNA secondary structure folding program mfold, by M. Zuker (website 1999-2000:

http://mfold2.wustl.edu/~mfold/rna/form1.cgi). Sub-optimal folds with a free energy value within 5% of the predicted most stable fold of the mRNA were predicted using a window of 200 bases within which a residue can find a complimentary base to form a base pair bond. Open regions that did not form a base pair were summed together with each suboptimal fold and areas that consistently were predicted as open were considered more accessible to the binding of antisense oligonucleotides. Additional oligonucleotides that only partially fulfilled some of the above selection criteria (1-4), were also chosen as possible candidates if they recognized a predicted open region of the target mRNA.

15

20

25

10

Example 3: Antisense oligonucleotide synthesis

The antisense oligonucleotides were synthesized by IDT (Integrated DNA Technologies, USA) as chimeric, second-generation oligonucleotides, consisting of a core of phosphodiester DNA residues flanked on either side by two 2'-O methyl RNA residues with a phosphorothioate linkage between the flanking RNA residues. The oligonucleotides were provided in a 96-well plate, as well as matching tubes, with a minimum of 12 ODs of oligo DNA, which provided ample material for transfections (greater than a hundred assays in the 96-well format) when the detection method is a sensitive method, such as TaqMan quantitative PCR, or an ELISA. Once the positive hits were identified (see below), the antisense oligonucleotides were re-synthesized with 3, instead of 2, flanking RNA residues to further increase stability/nuclease resistance. In addition, for validation purposes, appropriate controls (such

as scrambled, 4-base mismatch, and reverse polarity oligonucleotides) were synthesized for some of the antisense targets that yielded the highest antisense activity.

5 Example 4: Screening assays and optimization of antisense oligonucleotide sequences

15

20

25

Our approach to identifying IAP antisense oligonucleotides was to screen the above-described antisense oligonucleotide libraries for specific decreases (knock-down) of the RNA and protein for the specific IAP gene targeted. Any number of standard assays may be used to detect RNA and protein levels in cells that have been administered an IAP antisense nucleic acid. For example, RNA levels can be measured using standard Northern blot analysis or RT-PCR techniques. In addition, protein levels can be measured, for example, by standard Western blot analyses or immunoprecipitation techniques. Alternatively, cells administered an antisense IAP nucleic acid may be examined for cell viability, according to methods described for example, in U.S. Patent No. 5,919,912, or U.S.S.Ns. 08/576,956, 08/800,929, incorporated herein by reference.

We used TaqMan quantitative PCR conditions (described below) to assay for changes in mRNA levels after antisense oligonucleotide treatment, as well as our ELISA method for XIAP and Western blotting (described below) for changes in HIAP1 protein levels, using a polyclonal anti-RIAP1 antibody (rat HIAP1 ortholog; AEgera Therapeutics, Inc.) in the latter case. Transfection conditions were optimized with LipofectAMINE PLUS (Life Technologies, Canada) on T24 bladder carcinoma cells, or lipofectin on SF-295 glioblastoma cells, using a fluorescein-tagged control sense oligo from XIAP spanning the start codon (mGmAG AAG ATG ACT GGT AAmC mA; SEQ ID NO:

195). The results were visualized and gauged by epi-fluorescence microscopy. In addition, in the case of T24 cells, transfections were further optimized based on the ability of a published antisense oligonucleotide to downregulate survivin expression (Li et al., Nat. Cell Biol. 1:461-466, 1999) (U/TGT GCT ATT CTG TGA AU/TU/T SEQ ID NO: 196). We optimized the transfection conditions based on the TaqMan results of survivin RNA knock-down detected with PCR primers and fluorescent probe, described in detail below. Optimal conditions for oligo uptake by the cells were found to be 940 nM oligonucleotide and 40 μ L PLUS reagent and 0.8 μ L LipofectAMINE in a total of 70 μ L for 3 hours. We then applied these conditions to screen for XIAP protein knock-down using the oligo library against T24 cells.

10

15

25

HIAP1 knock-down was studied in SF-295 cells because these cells had easily detectable and discernable 70 kDa HIAP1 protein, while many cell lines do not express high levels of the protein, or are not distinguishable from the large amounts of the similarly sized 68 kDa HIAP2 protein. In fact, there are a number of published errors involving HIAP1 and HIAP2 in the literature because of naming errors in the databases, and because of the poor quality and high crossreactivity, of the various commercial antibodies to HIAP1/cIAP2. The best way to distinguish HIAP1 from HIAP2 is to perform an immunoprecipitation experiment with an IAP antibody (Aegera Therapeutics, Inc.), separate the proteins by 2-dimensional gel electrophoresis, and to then carry out mass spectroscopy analysis of the spots migrating in the 68 to 70 kDa range to verify the identity of the HIAP1 and HIAP2 bands, using standard methods known in the art. This method determines if HIAP1 and HIAP2 co-migrate at the 68 kDa position, and if the 70 kDa form of HIAP1 results from a splice variant or a post-translational modification.

Real-time PCR

RNA was extracted from cells lysed in RLT buffer (QIAGEn, Inc., Canada), and purified using QIAGEN RNeasy columns/kits. Real-time quantitative PCR was performed on a Perkin-Elmer ABI 7700 Prism PCR machine. RNA was reverse transcribed and amplified according to the TaqMan Universal PCR Master Mix protocol of PE Biosystems, using primers and probes designed to specifically recognize XIAP, HIAP1, survivin, or GAPDH. For human survivin, the forward primer was 5'-TCT GCT TCA AGG AGC TGG AA-3', the reverse primer was 5'-GAA AGG AAA GCG CAA CCG-3', and the probe was 5'-(FAM) AGC CAG ATG ACG ACC CCA TAG AGG AAC ATA(TAMRA)-3' (SEQ ID NOS: 197 through 199). For human HIAP1, the forward primer was 5'-TGG AGA TGA TCC ATG GGT TCA-3', the reverse primer was 5'-GAA CTC CTG TCC TTT AAT TCT TAT CAA GT-3', and the probe was 5'-(FAM) CTC ACA CCT TGG AAA CCA CTT GGC ATG(TAMRA)-3' (SEQ ID NOS: 200 through 202). For human XIAP, the forward primer was 5'-GGT GAT AAA GTA AAG TGC TTT CAC TGT-3', the reverse primer was 5'-TCA GTA GTT CTT ACC AGA CAC TCC TCA A-3', and the probe was 5'-(FAM) CAA CAT GCT AAA TGG TAT CCA GGG TGC AAA TAT C(TAMRA)-3' (SEQ ID NOS: 203 through 205). For human GAPDH, the forward primer was 5'-GAA GGT GAA GGT CGG AGT C-3', the reverse primer was 5'-GAA GAT GGT GAT GGG ATT C-3', and the probe was 5'-(JOE) CAA GCT TCC CGT TCT CAG CC(TAMRA)-3' (SEQ ID NOS: 206 through 208). 25

Relative quantitation of gene expression was performed as described in the PE Biosystems manual using GAPDH as an internal standard. The comparative Ct (cycle threshold) method was used for relative quantitation of IAP mRNA levels compared to GAPDH mRNA

levels. Briefly, real-time fluorescence measurements were taken at each PCR cycle and the threshold cycle (Ct) value for each sample was calculated by determining the point at which fluorescence exceeded a threshold limit of 30 times the baseline standard deviation. The average baseline value and the baseline SD are calculated starting from the third cycle baseline value and stopping at the baseline value three cycles before the signal starts to exponentially rise. The PCR primers and/or probes for ecific IAPs were designed to span at least one exon-intron boundary

y 1 or more kb of genomic DNA, to reduce the possibility of amplifying and detecting genomic DNA contamination. The specificity of the signal, and possible contamination from DNA, were verified by some RNA samples with either DNase or RNase, prior to g the reverse transcription and PCR reaction steps.

XIAP ELISA and HIAP1 Western immunoblots

15

20

25

A standard colorimetric XIAP ELISA assay was performed using an affinity-purified rabbit polyclonal antibody to XIAP (Aegera Therapeutics, Inc.) as a capture antibody, and was detected with a XIAP monoclonal antibody (MBL, Japan) and a biotinylated anti-mouse Ig antibody and horseradish peroxidase-conjugated streptavidin and TMB substrate. Alternatively, a polyclonal XIAP or HIAP1 antibody may be used to measure XIAP or HIAP1 protein levels, respectively.

HIAP1 was detected on a Western immunoblot using an affinity-purified anti-RIAP1 rabbit polyclonal antibody as a primary antibody and was detected by ECL (Amersham) on X-ray film with a secondary horseradish-peroxidase-conjugated anti-rabbit Ig antibody and a chemiluminescent substrate. The anti-RIAP1 polyclonal antibody is raised against a GST-fusion of the rat ortholog of HIAP1. This antibody cross-reacts with both human and murine HIAP1 and HIAP2.

Example 5: Antisense XIAP oligonucleotides decrease XIAP RNA and polypeptide expression

The XIAP synthetic library of 96 antisense oligonucleotides was first screened for decreases in XIAP protein levels. Specifically, T24 cells (1.5 x 10⁴ cells/well) were seeded in wells of a 96-well plate on day 1, and were cultured in antibiotic-free McCoy's medium for 24 hours. On day 2, the cells were transfected with XIAP antisense oligonucleotides as described above (oligonucleotides are labeled according to their plated position, i.e., A1 to H12, and include 2 repeats, A13 and B13 that contain lyophilized DNA pellets that stuck to the sealing membrane). Briefly, the oligos were diluted in 10 μl/well of serum-free, antibiotic-free McCoy's medium and then PLUS reagent was added. LipofectAMINE was diluted in 10 μl/well of serum-free, antibiotic-free McCoy's medium, and both mixes were incubated for 15 minutes at room temperature. The mixes were then combined and incubated for 15 minutes at room temperature.

10

15

In the meantime, the complete medium was removed from the cells and 50 μ l/well of serum-free, antibiotic-free medium was added to the cells. The transfection mixes were added to the well, and the cells were incubated for 3 hours. Then 30 μ l/well of serum-free, antibiotic-free medium and 100 μ l/well of antibiotic-free complete medium, containing 2X fetal bovine serum were added to each well.

At day 3, XIAP RNA levels were measured using quantitative real-time PCR techniques, as described above. At day 4, XIAP protein levels were measured by ELISA (Figs. 7A, 7C, 7E, 7G, 7I, and 7K), and total cellular protein was measured biochemically (Figs. 7B, 7D, 7F, 7H, 7J, and 7L; used to normalize the XIAP protein levels). The results were compared to a mock transfection sample (treated with the transfection agent but no oligonucleotide DNA was added, and then processed as for

the other samples). Time course experiments determined that the optimal time for protein knock-down to be around 12 to 24 hours.

The library was also screened for decreases in RNA levels, using TaqMan-specific PCR primers and fluorescent probes at the appropriate optimal time, using the primers and probes described above. Time course experiments determined mRNA to be optimally decreased at 6 to 9 hours. These results agree well with the protein results.

10

20

25

The first screen (although performed at a sub-optimal time point when XIAP levels are returning to normal, possibly due to an outgrowth of non-transfected cells) identified 16 antisense oligonucleotides (ODNs C2, E2, E3, F3, C4, D4, E4, F4, G4, C5, D5, B6, F6, D7, D8, F8) out of the total 96 antisense oligonucleotides tested that showed some decrease in XIAP protein levels relative to total protein, compared to mock (no ODN) transfection levels (Fig. 7A, 7C, 7E, 7G, 7I, and 7K). Interestingly, total protein was decreased for each of these 16 ODNs, which indicates a toxic or cytostatic effect of these ODNs (Fig. 7B, 7D, 7F, 7H, 7J, 7L). Note that ODNs B9 and C9 showed a clear drop in total protein but no relative drop in XIAP protein levels. These 16 hits were then validated more rigorously at more optimal time points XIAP protein and RNA knock-down results at 12 hours after the start of transfection.

The 16 antisense ODNs that showed some decrease in relative XIAP protein levels compared to mock transfection, were re-tested alone or in combination, with one control oligo (D2) included, for their ability to knock-down XIAP protein at a more optimal time point (12 hours) based on the above described time course studies (Fig. 8B). these ODNs were also examined for their ability to decrease XIAP mRNA levels at 12 hours, normalized against GAPDH levels, and compared to mock transfection. Total protein concentrations at 12 hours were also determined (Fig. 8C).

There was a good correlation between the ability of an antisense ODN to decrease XIAP protein levels (Fig. 8B) with its ability to decrease XIAP mRNA levels (Fig. 8A). In addition, there is no major loss of total protein at this early time point, and the decrease in XIAP mRNA and protein precede the decrease in total protein that is seen at later time points. The ODNs that showed greater than 50% loss of XIAP protein or mRNA levels alone, or in a combination of two ODNs added at a 0.5:0.5 ratio, were identified as the best ODNs and validated further. Of these 16 oligonucleotides, 10 of them (ODNs E2, E3, F3, E4, F4, G4, C5, B6, D7, F8) showed a consistent ability to decrease XIAP protein or RNA levels by more than 50%, depending on the transfection conditions used, or when used in combination, as for ODNs C5 and G4.

Interestingly, these 16 oligonucleotides that demonstrated antisense activity clustered in 4 different target regions of the XIAP mRNA, with adjacent ODNs showing some knock-down activity. No antisense activity was observed by oligonucleotides that target sequences between these regions or islands of sensitivity. Presumably, these regions represent open areas on the mRNA that are accessible to antisense ODNs inside the cell. Two antisense oligonucleotides, E3 and F3, target XIAP just upstream of the start codon in the intervening region between the IRES and the translation start site, and partially overlap the end of the IRES element. ODNs C2, D2, and E2 target a XIAP region upstream of the minimal IRES element, providing further evidence that the minimal IRES region is a highly structured region of RNA which is not readily accessible to antisense ODNs in vivo. All the other antisense ODN hits fall within the coding region, including a cluster of activity at positions 856-916 of the HIAP sequence of Fig. 15 (ODNs E4, F4, and G4) and smaller separate areas, as demonstrated by ODNs C5 and D5, for example.

15

20

Example 6: XIAP antisense oligonucleotides increase cytotoxicity and chemosensitization

We also investigated if XIAP antisense ODNs could chemosensitize the highly drug resistant T24 cells to traditional chemotherapeutic drugs, such as adriamycin or cisplatin. Antisense ODNs were chosen to represent some of the different XIAP target regions and were tested for their cytotoxic effects, alone or in combination with other ODNs or drugs. Five of the ten best XIAP antisense oligonucleotides were tested for their ability to kill or chemosensitize T24 bladder carcinoma cells, and were compared to the effects of three corresponding scrambled control ODNs.

10

15

20

25

T24 cells were transfected with XIAP antisense oligonucleotides, scrambled oligonucleotides, no oligonucleotides (mock transfected), or were left untreated. The cells were tested for viability 20 hours after transfection (with the exception of the untreated control) using the WST-1 tetrazolium dye assay in which WST-1 tetrazolium dye is reduced to a colored formazan product in metabolically active cells (Fig 9A). Alternatively, cell viability is tested using any one of the above described apoptosis methods.

The occurrence of cytoxicity induced by the antisense XIAP ODN E4 was examined by visually inspecting T24 cells that were left untreated, mock transfected, or transfected with E4 antisense ODNs, E4 scrambled ODNs, E4 reverse polarity, or E4 mismatched ODNs. Twenty hours after transfection, the cells were examined for morphology (Fig. 9D). Only the cell transfected with antisense E4 ODNs showed signs of toxicity.

To examine the effects of the oligonucleotides on the chemosensitization of the T24 cells to cisplatin or adriamycin, oligonucleotides were tested for their ability to further kill T24 cells in the presence of a fixed dose of adriamycin (0.5 μ g/ml). Cells were first

transfected with the oligonucleotides, then adriamycin was added for another 20 hours. Viability was measured by WST-1 at the end of the 20 hour drug treatment (Fig. 9B). Values are shown as percentage viability compared to their ODN treatment alone results shown in Fig. 9C. Fig. 9C is essentially a repeat of Fig. 9A, but with the actual corresponding values used in calculating the results for the chemosensitization experiment in Fig. 9B.

All 5 oligonucleotides tested (ODNs F3, E4, G4, C5, D7, or the combinations of E4+C5, or G4+C5) killed the T24 cells, leaving only 10-15% surviving cells after 24 hours, as compared to the mock (no ODN) 10 transfected cells, or to cells transfected with 3 corresponding scrambled controls to F3 (mCmAmG AGA TTT CAT TTA AmCmG mU; SEQ ID NO: 209), E4 (mCmUmA CgC TCg CCA TCg TmUmC mA; SEQ ID NO: 210) and C5 (mUmGmC CCA AGA ATA CTA GmUmC mA; SEQ ID NO: 211)(Figs. 9A and C). Therefore, the toxicity is sequence-specific 15 to those ODNs that reduce XIAP levels, and not to a non-sequence specific toxicity due to ODNs of this chemistry in general, as three scrambled controls did not show any more toxicity compared to the mock transfected control. This cytotoxicity may result from the combined effect of XIAP protein knock-down (and the expected loss of anti-apoptotic 20 protection afforded by XIAP) and the cytotoxicity of the transfection itself. Both mock (no ODN) and scrambled ODN transfections resulted in an approximately 40% decrease in survival as compared to untreated cells (Fig. 9A). This is not unexpected, as the opposite is true (i.e., overexpression of IAPs protect insect cells from cytofectin-mediated cell 25 death, a liposomal transfection agent similar to the ones used in these studies (Jones et al., J. Biol. Chem. 275:22157-22165, 2000)

The addition of a fixed dose of adriamycin or cisplatin at the end of the 3 hour transfection period resulted in a further decrease in survival for

some of the tested oligonucleotides, a further 40% drop in survival after 20 hours for ODNs F3, D7 and G4+C5 combination (Fig. 9B), compared to their corresponding ODNs treated values (Fig. 9C). Note that the values in Fig. 9B (ODN plus drug) are compared to their corresponding ODN survival (ODN alone) in Fig. 9C, which is set a 100% for each ODN. Only the results for adriamycin chemosensitization are shown; however, similar results were obtained when the cells were chemosensitized with cisplatin. At the fixed doses used, the mock and scrambled control transfections did not show any increased loss of survival when either treated with adriamycin (Fig. 9B). Chemosensitization is only seen when XIAP levels are decreased by a specific antisense ODN.

Example 7: Antisense HIAP1 oligonucleotides decrease HIAP1 RNA and polypeptide expression

15

20

The smaller library of 15 HIAP1 antisense oligonucleotides was screened for protein knock-down by Western and for RNA knock-down by TaqMan, using the primers and probes described above, under two different conditions. Alternatively, HIAP1 RNA levels may be detected using standard Northern blot analyses or RT-PCR techniques. The antisense oligonucleotides were administered to cells under basal conditions or under cycloheximide-induction conditions (24 hour treatment with sub-toxic doses). We have discovered that cycloheximide (CHX) can lead to a 10- to 200-fold induction of HIAP1 mRNA depending on the cell line treated. This in turn leads to an increase in HIAP1 protein, as seen on a Western blot (70 kDa band). We have also discovered that this effect of CHX is via two distinct mechanisms of action. First, CHX activates NFkB, a known transcriptional inducer of HIAP1, by blocking the *de novo* synthesis of a labile protein, IkB, which

is an inhibitor of NFkB. This effect is mimicked by puromycin, another protein synthesis inhibitor, and by TNF-alpha, which induces a signaling cascade leading to the phosphorylation, ubiquination, and degradation of IkB. However, only CHX leads to a further stabilization of the HIAP1 mRNA, as seen by the decreased rate of disappearance of HIAP1 message in the presence of actinomycin D, to block *de novo* transcription, and CHX, as opposed to actinomycin D and puromycin or TNF-alpha combined.

SF295 glioblastoma cells were transfected with lipofectin and ODN (scrambled survivin, no oligo or mock, antisense APO1 to APO15) or left untreated. RNA was isolated from the cells 6 hours after transfection and the level of HIAP1 mRNA was measured by quantitative PCR (TaqMan analysis), normalized for GAPDH mRNA, with the value for the scrambled survivin ODN transfection set as 1.0.

10

15

20

25

The results of this experiment, a compilation of three separate experiments, are shown in Fig. 10. The scrambled survivin ODN, the mock transfection, and untreated (non-transfected) cells, all showed similar HIAP1 mRNA levels. Of the 15 antisense ODNs, 7 oligonucleotides (ODNs APO 1, -2, -7, -8, -9, -12, -15) showed an almost 50% decrease when compared to mock transfection or survivin scrambled control (mUmAmA GCT GTT CTA TGT GmUmU mC; SEQ ID NO: 212) ODN transfection (Fig. 10). Some of the ODNs led to an induction in HIAP1 mRNA, which may be a stress response to a non-specific toxic ODN. The antisense ODN, however, may still be effective at knocking down HIAP1 protein levels even if the message is increased if the ODN is able to interfere with the translation process.

The effect of HIAP1 antisense oligonucleotides on HIAP1 protein and mRNA expression was also examined in cells induced to express HIAP1. SF295 cells were transfected with ODNs, or were mock

transfected. The transfected cells were then treated with 10 µg/ml cycloheximide for 24 hours to induce 70 kDa HIAP1 mRNA and protein. Protein levels were measured by Western immunoblot analysis with an anti-RIAP1 polyclonal antibody, and normalized against actin protein in a re-probing of the same blots. Scans of the Western blot results are shown in Fig. 11A. The densitometric scan results were plotted against the mock results (set at 100%) in Fig. 11B. A line is drawn at 50% to easily identify the most effective antisense ODNs. The transfection process itself (e.g., mock or scrambled survivin) induces HIAP1 protein compared to the untreated sample as shown on the Western immunoblot.

10

15

25

Of the 15 tested ODNs, 6 of them (APO 1, -2, -7, -8, -12, and -15) showed the strongest activity, or had significant activity in both the protein and mRNA assays, and did not cause a stress-induced increase in HIAP1 mRNA, such as that seen with ODNs APO 4, -6, -11, -13, -14 (Fig. 10), and by control ODNs to APO 2 (mismatch or reverse polarity, see text below and Figs.12 and 13). Note that APO 6 also showed evidence of toxicity as seen by the general decrease in total protein (Fig. 12).

To further investigate the efficacy of HIAP1 antisense oligonucleotides under cycloheximide induction conditions, changes in HIAP1 mRNA were measured by TaqMan real time PCR 6 hours after transfection with ODN APO 2, which targets an Alu repeat within an intron of HIAP1 and results in the greatest block of CHX-induced upregulation of HIAP1 mRNA and protein. Controls for this experiment were three ODNs for APO 2: one scrambled sequence (same base composition but random order, AAG GGC GGC GGA GTG AGA C; SEQ ID NO: 213), one reverse polarity (same base composition, same sequential order but in the opposite direction, AGA GGA CGG AGT CGG

AGG C; SEQ ID NO: 214), and one mismatch sequence (containing 4 base mismatches out of 19 bases, CGG AGC GTG AGG ATG GAG A; SEQ ID NO: 215).

5

10

15

20

25

Transfection of the APO 2 antisense into cells resulted in a 50% decrease in mRNA compared to a scrambled survivin control and matched perfectly with the protein results, while the scrambled control for APO 2 (H1 sc apo 2 in Fig. 13) did not change HIAP1 mRNA levels at all (repeated twice here, and in two different experiments). However, the mismatch control ODN (H1 mm apo 2) and the reverse polarity control ODN (H1 RV apo 2) showed an induction of 6 to 7 fold in HIAP1 mRNA at 6 hours. These ODNs no longer targeted HIAP1, as expected, but may still target Alu repeats because of the degeneracy and repeat nature of these sequences. Therefore, it is possible that these two controls are toxic to the cell and cause a stress response that leads to the induction of HIAP1. This effect may also occur with the antisense APO 2 ODN, but in this case, the APO 2 ODN also causes the degradation of the induced HIAP1 mRNA which results in a relative decrease of HIAP1 mRNA, compared to a scrambled survivin control, as well as decreasing the relative fold induction of HIAP1 protein after transfection and CHX treatment, compared to scrambled survivin control ODN.

The 6 optimal antisense HIAP1 ODNs include two very effective antisense ODNs against an intronic sequence (APO 1, and -2; with APO 2 demonstrating the best activity). These oligonucleotides have some interesting properties that could be of great use therapeutically for cancer or autoimmune disorders. The oligonucleotides against an intronic sequence would likely only target pre-mRNA (very short-lived target) and not the mature, processed form of HIAP1. Typically, introns are not targeted for antisense except when one wants to alter splicing by targeting the intron-exon boundaries or the branching point. These usually result in

the skipping of an exon rather than RNase-mediated degradation of the message. Both mechanisms would likely be favorable for the enhancement of apoptosis, as the skipping would result in the loss of the exon encoding the first two important BIR domains of HIAP1. The APO-2 antisense oligo also targets an intron of survivin for 18 consecutive bases out of 19, but we did not see any loss of survivin protein; only HIAP1 was decreased after the oligo treatment, demonstrating the specificity of the HIAP1 antisense oligonucleotide. These antisense oligonucleotides hit Alu sequences in the HIAP1 intron and potentially in many other genes, and induce the cancer cells to die (see below), which may be as a result of down regulating HIAP1 and some other critical genes, and thus of therapeutic value if it is not too toxic to normal cells.

Cancer cells have reportedly more Alu-containing transcripts and may therefore be more sensitive to apoptosis induction with an Alu targeting antisense ODN. Furthermore, this killing effect of APO 1 and APO 2 ODNs may be due to the combined effect of both targeting Alu sequences and HIAP1 simultaneously. This dual effect would result in an effective way to prevent the normal stress response of HIAP1 induction through the NFkB pathway, when the cell is exposed to certain toxic agents. This stress response is most likely part of the cancer cell's anti-apoptotic program. By blocking HIAP1 expression, we counter this anti-apoptotic stress response and precipitate the cancer cell's demise.

15

20

25

Example 8: HIAP1 antisense oligonucleotides increase cytotoxicity and chemosensitization

The effect of HIAP1 antisense oligonucleotides on the chemosentization of SF295 cells was also evaluated. Cells were transfected with one of 3 different antisense ODNs (APO 7, APO 15, and Scrambled APO 2 (control)). Twenty-four hours after transfection with the

ODNs, the cells were incubated with adriamycin for an additional 24 hours before assaying by for cell survival by assaying WST-1.

The WST-1 survival curves for SF295 cells transfected with the above-described HIAP1 ODNs and then treated with increasing concentrations of adriamycin are shown in Fig. 14. The two ODNs that resulted in a decrease in HIAP1 mRNA also showed a decrease in survival when treated with adriamycin compared to cells treated with an ODN which did not reduce HIAP1 mRNA levels. Therefore, reducing HIAP1 levels by antisense, or other means, can chemosensitize a glioblastoma cell line that is highly resistant to the cytotoxic action of many chemotherapeutic drugs.

Example 9: In vivo analyses of IAP antisense oligonucleotides

10

20

25

Antisense oligonucleotides that decrease expression of IAP in cell culture models can be tested in animals. For example, the antisense oligonucleotide can be tested in mice according to the method of Lopes de Menezes et al. (Clin. Cancer Res. 6: 2891-2902, 2000) or Klasa et al. (Clin. Cancer Res. 6: 2492-2500, 2000). Antisense and control ODNs are tested, for example, in sub-cutaneous human xenografts of breast cancer, colon cancer, lung cancer, squamous cell carcinoma or prostate cancer in SCID mice. The antisense oligonucleotides are also tested in an orthotopic model for the prostate, as well as in a disseminated non-Hodgkin's lymphoma model. The mouse's tolerance to cisplatin, taxol, doxorubicin, and cyclophosphamide is known for each of these models.

In vivo testing of the antisense oligonucleotides involves 15 intraperitoneal injections (once a day, on days 3 through 7, 10 through 14, and 17 through 21) of naked ODN of (5 mg/kg), with or without a chemotherapeutic drug. Alternatively, liposomal type carriers for the ODNs may also be employed. Oligos are injected shortly after tumors

cells have been seeded in the mouse, or when the tumor has established and grown to a size of 0.1-0.15 g. Tumor size is then monitored to determine if the ODN treatments or ODN plus drug treatments reduce the growth rate of the tumor, lead to regression, or have no effect at all. In another alternative, ODNs in liposomal formulation are injected directly into the tumors.

Example 10: Anti-IAP antibodies

5

10

15

20

25

In order to generate IAP-specific antibodies, an IAP coding sequence (e.g., amino acids 180-276) can be expressed as a C-terminal fusion with glutathione S-transferase (GST; Smith et al., Gene 67:31-40, 1988). The fusion protein can be purified on glutathione-Sepharose beads, eluted with glutathione, and cleaved with thrombin (at the engineered cleavage site), and purified to the degree required to successfully immunize rabbits. Primary immunizations can be carried out with Freund's complete adjuvant and subsequent immunizations performed with Freund's incomplete adjuvant. Antibody titres are monitored by Western blot and immunoprecipitation analyses using the thrombin-cleaved IAP fragment of the GST-IAP fusion protein. Immune sera are affinity-purified using CNBr-Sepharose-coupled IAP protein. Antiserum specificity is determined using a panel of unrelated GST proteins (including GSTp53, Rb, HPV-16 E6, and E6-AP) and GST-trypsin (which was generated by PCR using known sequences).

As an alternate or adjunct immunogen to GST fusion proteins, peptides corresponding to relatively unique hydrophilic regions of IAP may be generated and coupled to keyhole limpet hemocyanin (KLH) through an introduced C-terminal lysine. Antiserum to each of these peptides is similarly affinity-purified on peptides conjugated to BSA, and specificity is tested by ELISA and Western blotting, using peptide

conjugates, and by Western blotting and immunoprecipitation using IAP expressed as a GST fusion protein.

Alternatively, monoclonal antibodies may be prepared using the IAP proteins described above and standard hybridoma technology (see, e.g., Kohler et al., Nature 256:495, 1975; Kohler et al., Eur. J. Immunol. 6:511, 1976; Kohler et al., Eur. J. Immunol. 6:292, 1976; Hammerling et al., In Monoclonal Antibodies and T Cell Hybridomas, Elsevier, New York, NY, 1981; Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, 1994). Once produced, monoclonal antibodies are also tested for specific IAP recognition by Western blot or immunoprecipitation analysis (by the methods described in Ausubel et al., supra).

10

15

20

25

Antibodies that specifically recognize IAPs or fragments of IAPs, such as those described in U.S.S.N. 08/800,929, incorporated herein by reference, containing one or more BIR domains (but not a ring zinc finger domain), or that contain a ring zinc finger domain (but not a BIR domain) are considered useful in the invention. They may, for example, be used in an immunoassay to monitor IAP expression levels or to determine the subcellular location of an IAP or IAP fragment produced by a mammal. Antibodies that inhibit the 26 kDa IAP cleavage product described herein (which contains at least one BIR domain) may be especially useful in inducing apoptosis in cells undergoing undesirable proliferation.

Preferably antibodies of the invention are produced using IAP sequence that does not reside within highly conserved regions, and that appears likely to be antigenic, as analyzed by criteria such as those provided by the Peptide structure program (Genetics Computer Group Sequence Analysis Package, Program Manual for the GCG Package, Version 7, 1991) using the algorithm of Jameson and Wolf (CABIOS 4:181, 1988). Specifically, these regions, which are found between BIR1

and BIR2 of all IAPs, are: from amino acid 99 to amino acid 170 of HIAP1, from amino acid 123 to amino acid 184 of HIAP2, and from amino acid 116 to amino acid 133 of either XIAP or m-XIAP. These fragments can be generated by standard techniques, e.g., by the PCR, and cloned into the pGEX expression vector (Ausubel et al., supra). Fusion proteins are expressed in E. coli and purified using a glutathione agarose affinity matrix as described in Ausubel et al. (supra). In order to minimize the potential for obtaining antisera that is non-specific, or exhibits low-affinity binding to IAP, two or three fusions are generated for each protein, and each fusion is injected into at least two rabbits. Antisera are raised by injections in series, preferably including at least three booster injections.

10

15

25

Example 11: Comparison of cell survival following transfection with full length vs. partial IAP constructs

In order to investigate the mechanism whereby human IAPs, including XIAP, HIAP1, and HIAP2, afford protection against cell death, expression vectors were constructed that contained either: (1) full-length IAP cDNA (as described in U.S.S.N. 08/800,929), (2) a portion of an IAP gene that encodes the BIR domains, but not the RZF, or (3) a portion of an IAP gene that encodes the RZF, but not the BIR domains. Human and murine XIAP cDNAs were tested by transient or stable expression in HeLa, Jurkat, and CHO cell lines. Following transfection, apoptosis was induced by serum withdrawal, application of menadione, or application of an anti-Fas antibody. Cell death was then assessed by trypan blue exclusion. As a control for transfection efficiency, the cells were cotransfected with a Beta-gal expression construct. Typically, approximately 20% of the cells were successfully transfected.

When CHO cells were transiently transfected, constructs containing full-length human or mouse XIAP cDNAs conferred modest but definite protection against cell death. In contrast, the survival of CHO cells transfected with constructs encoding only the BIR domains (i.e., lacking the RZF domain) was markedly enhanced 72 hours after serum deprivation. Furthermore, a large percentage of cells expressing the BIR domains were still viable after 96 hours, at which time no viable cells remained in the control, i.e. non-transfected, cell cultures, and less than 5% of the cells transfected with the vector only, i.e., lacking a cDNA insert, remained viable. Deletion of any of the BIR domains results in the complete loss of apoptotic suppression, which is reflected by a decrease in the percentage of surviving CHO cells to control levels within 72 hours of serum withdrawal.

10

15

20

25

Stable pools of transfected CHO cells, which were maintained for several months under G418 selection, were induced to undergo apoptosis by exposure to 10 µM menadione for 2 hours. Among the CHO cells tested were those that were stably transfected with: (1) full-length murine XIAP cDNA (miap), (2) full-length XIAP cDNA (XIAP), (3) full-length bcl-2 cDNA (Bcl-2), (4) cDNA encoding the three BIR domains (but not the RZF) of murine XIAP (BIR), and (5) cDNA encoding the RZF (but not BIR domains) of m-XIAP (RZF). Cells that were non-transfected (CHO) or transfected with the vector only (pcDNA3), served as controls for this experiment. Following exposure to 10 μM menadione, the transfected cells were washed with phosphate buffered saline (PBS) and cultured for an additional 24 hours in menadione-free medium. Cell death was assessed, as described above, by trypan blue exclusion. Less than 10% of the non-transfected or vector-only transfected cells remained viable at the end of the 24 hour survival period. Cells expressing the RZF did not fare significantly better. However, expression of full-length

murine XIAP, human XIAP, or bcl-2, and expression of the BIR domains, enhanced cell survival. When the concentration of menadione was increased from 10 µM to 20 µM (with all other conditions of the experiment being the same as when 10 µM menadione was applied), the percentage of viable CHO cells that expressed the BIR domain cDNA construct was higher than the percentage of viable cells that expressed either full-length murine XIAP or bcl-2.

Example 12: Analysis of the subcellular location of expressed RZF and BIR domains

10

15

20

25

The assays of cell death described above indicate that the RZF acts as a negative regulator of the anti-apoptotic function of IAPs. One way in which the RZF, and possibly other IAP domains, may exert their regulatory influence is by altering the expression of genes, whose products function in the apoptotic pathway.

In order to determine whether the subcellular locations of expressed RZF and BIR domains are consistent with roles as nuclear regulatory factors, COS cells were transiently transfected with the following four constructs, and the expressed polypeptide was localized by immunofluorescent microscopy: (1) pcDNA3-6myc-xiap, which encodes all 497 amino acids of SEQ ID NO:219, (2) pcDNA3-6myc-m-xiap, which encodes all 497 amino acids of mouse XIAP (SEQ ID NO:225), (3) pcDNA3-6myc-mxiap-BIR, which encodes amino acids 1 to 341 of m-xiap (SEQ ID NO:225), and (4) pcDNA3-6myc-mxiap-RZF, which encodes amino acids 342-497 of murine xiap (SEQ ID NO:225). The cells were grown on multi-well tissue culture slides for 12 hours, and then fixed and permeabilized with methanol. The constructs used (here and in the cell death assays) were tagged with a human Myc epitope tag at the N-terminus. Therefore, a monoclonal anti-Myc antibody and a secondary

goat anti-mouse antibody, which was conjugated to FITC, could be used to localize the expressed products in transiently transfected COS cells. Full-length XIAP and MIAP were located in the cytoplasm, with accentuated expression in the peri-nuclear zone. The same pattern of localization was observed when the cells expressed a construct encoding the RZF domain (but not the BIR domains). However, cells expressing the BIR domains (without the RZF) exhibited, primarily, nuclear staining. The protein expressed by the BIR domain construct appeared to be in various stages of transfer to the nucleus.

10

15

20

Other Embodiments

All publications and patent applications mentioned in this specification, including U.S. Patent No. 5,919,912 and U.S.S.Ns. 08/576,956 and 08/800,929 are herein incorporated by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth.

Claims

 An inhibitor of apoptosis (IAP) antisense nucleic acid that inhibits IAP biological activity, regardless of length of said antisense nucleic acid.

- 2. The antisense IAP nucleic acid of claim 1, wherein said IAP is XIAP.
- 3. The antisense IAP nucleic acid of claim 1, wherein said IAP is HIAP1.
 - 4. The antisense IAP nucleic acid of claim 1, wherein said IAP is HIAP2.

15

- 5. The antisense IAP nucleic acid of claim 1, wherein said antisense nucleic acid is mammalian.
- 6. The antisense IAP nucleic acid of claim 5, wherein said antisense nucleic acid is human.
 - 7. The antisense nucleic acid of claim 1, wherein said antisense nucleic acid is between 8 and 30 nucleotides in length.
- 25 8. The antisense IAP nucleic acid of claim 2, wherein said antisense is chosen from any one of SEQ ID NOS: 1 through 96.
 - 9. The antisense IAP nucleic acid of claim 3, wherein said antisense is chosen from any one of SEQ ID NOS: 97 through 194.

10. The antisense IAP nucleic acid of claim 1, wherein said IAP biological activity is inhibition of apoptosis.

- 11. The antisense IAP nucleic acid of claim 1, wherein said IAP5 biological activity is inhibition of IAP polypeptide expression.
 - 12. The antisense IAP nucleic acid of claim 1, wherein said antisense nucleic acid comprises at least one modified internucleoside linkage.

10

13. The antisense IAP nucleic acid of claim 12, wherein said modified internucleoside linkage is selected from the group consisting of phosphorothicate, methylphosphonate, phosphotriester, phosphorodithicate, and phosphoselenate linkages.

- 14. The antisense IAP nucleic acid of claim 1, wherein said antisense nucleic acid comprises at least one modified sugar moiety.
- 15. The antisense IAP nucleic acid of claim 14, wherein said modified sugar moiety is a 2'-O methyl group.
 - 16. The antisense IAP nucleic acid of claim 1, wherein said antisense nucleic acid is a chimeric nucleic acid.
- 25 17. The antisense IAP nucleic acid of claim 16, wherein said chimeric nucleic acid comprises DNA residues linked together by phosphorothioate linkages, said DNA residues flanked on each side by at least one 2'-O methyl RNA residues linked together by phosphorothioate linkages.

18. The antisense IAP nucleic acid of claim 17, wherein said DNA residues are flanked on each side by at least 3 2'-O methyl RNA residues.

- 19. The antisense nucleic acid of claim 1, wherein said antisense nucleic acid is a ribozyme.
 - 20. A method of enhancing apoptosis in a cell, said method comprising administering a negative regulator of the IAP anti-apoptotic pathway to said cell.
- 21. The method of claim 20, wherein said negative regulator is an antisense IAP nucleic acid.

10

15

- 22. The method of claim 21, wherein said IAP is XIAP.
- 23. The method of claim 21, wherein said IAP is HIAP1.
- 24. The method of claim 21, wherein said IAP is HIAP2.
- 25. The method of claim 21, wherein said antisense nucleic acid is mammalian.
 - 26 The method of claim 25, wherein said antisense nucleic acid is human.
 - 27. The method of claim 22, wherein said antisense is chosen from any one of SEQ ID NOS: 1 through 96.

28. The method of claim 23, wherein said antisense is chosen from any one of SEQ ID NOS: 97 through 194.

- 29. The method of claim 21, wherein said antisense nucleic acid comprises at least one modified internucleoside linkage.
 - 30. The method of claim 29, wherein said modified internucleoside linkage is selected from the group consisting of phosphorothioate, methylphosphonate, phosphotriester, phosphorodithioate, and phosphoselenate linkages.
 - 31. The method of claim 21, wherein said antisense nucleic acid comprises at least one modified sugar moiety.
- 32. The method of claim 31, wherein said modified sugar moiety is a 2'-O methyl group.
 - 33. The method of claim 21, wherein said antisense nucleic acid is a chimeric nucleic acid.
 - 34. The method of claim 33, wherein said chimeric nucleic acid comprises DNA residues linked together by phosphorothioate linkages, said DNA residues flanked on each side by at least one 2'-O methyl RNA residues linked together by phosphorothioate linkages.
 - 35. The method of claim 34, wherein said DNA residues are flanked on each side by at least 3 2'-O methyl RNA residues.

25

20

36. The method of claim 21, wherein said administration sensitizes said cell to chemotherapy.

- 37. The method of claim 21, wherein said administration sensitizes said cell to radiotherapy.
 - 38. The method of claim 20, wherein said negative regulator is an antibody that specifically binds an IAP polypeptide.
- 39. The method of claim 20, wherein said negative regulator is an IAP polypeptide comprising a ring zinc finger, said polypeptide having no more than two BIR domains.
- 40. The method of claim 20, wherein said negative regulator is a nucleic acid encoding the ring zinc finger domain of an IAP polypeptide.
 - 41. The method of claim 20, wherein said negative regulator is a compound that prevents cleavage of the IAP polypeptide.
- 20 42. The method of claim 20, wherein said cell is in vitro.
 - 43. The method of claim 20, wherein said cell is in vivo.
- 44. The method of claim 43, wherein said cell is in a mammal diagnosed with a proliferative disease.
 - 45. A pharmaceutical composition comprising a mammalian IAP antisense nucleic acid.

46. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the human XIAP gene or mRNA.

- 5 47. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the human HIAP1 gene or mRNA.
- 48. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the human HIAP2 gene or mRNA.
 - 49. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the murine XIAP gene or mRNA.
 - 50. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the murine HIAP1 gene or mRNA.

- 51. The pharmaceutical composition of claim 45, wherein said antisense nucleic acid binds a target sequence of the murine HIAP2 gene or mRNA.
- 52. The pharmaceutical composition of claim 45, wherein said mammalian antisense IAP nucleic acid is human antisense nucleic acid.
 - 53. The pharmaceutical composition of claim 46, wherein said antisense is chosen from any one of SEQ ID NOS: 1 through 96.

54. The pharmaceutical composition of claim 47, wherein said antisense is chosen from any one of SEQ ID NOS: 97 through 194.

1/67

FIG. 1A	₹									E	HUMAN xiap	z Xia	<u>Q</u>									
	4	gaaa	aaa.	99t)	gga -+-	Caac	Jt C	ctaa	at t	Cas	caagagaagatgacttttaa +	jaag	atg	act	ttt	aac	agt 	1 t t	cagttttgaag	1996	aggtggacaagtcctaatttcaagagaagatgacttttaacagttttgaaggatct	9
ಹ			•										×	E	Ct.	z	S	(tı	臼	O	S	ı
	61		aaaac	ttg	tgt -+-	acc	rg i	# + + + + + + + + + + + + + + + + + + +	cato	aa	caac		gaa	ıgaa	t t	g Fa	gaa	gaç	Jt t	aa	cttgtgtacctgcagacatcaataaggəagaagaatttgtagaagagtttaataga	120
ď		×	E	U	>	Δ,	4	Ω	н	Z	×	网	凹	E	ſz,	>	(H)	臼	ſz,	z	œ	
	121		ttaaa	aac	1 1 1 1	acttttgctaattttcc	taa	######################################	i Ö	aagt	Egg!	ggtagt	tcct	gt	+ t C3	g Ca	r Ca	ttcagcatcaacact -+	act c	990	aaacttttgctaattttccaagtggtagtcctgtttcagcatcaacactggcacga	180
rđ		,a	×	E٠	(24	Æ	z	Ŀ	Ω,	c)	Ø	ល	Ω,	>	(C)	4	ഗ	E	H	4	ĸ	•
•	181		gcagg	gtt	# # # # # # # # # # # # # # # # # # #	tcttta -+	tac	tgg +	tggtgaag +	agga 	aga	gataco	cgtgcg	JCG	gtga	ottt 	agt	tgt +	ttgtcat	T G	ggtttctttatactggtgaaggagataccgtgcggtgctttagttgtcatgcagct	240
đ		Æ	Ö	Ţ	H	>	E	Ö	បា	O	Ω	E	>	α¢	U	£14	ល	U	Ħ	K	æ	1
	241		gtaga	tag	atgg -+	gca	ata 	tgg +	atatggagactcagcagtt 	ot C	900	agt (ස ය -	acaca -+	ago	Jaac	igt.	gtato	CCCa	atagatggcaatatggagactcagcagttggaagacacaggaaagtatccccaaat	300
rð		>	Ω	C	3	ø	> +	Ö	Ω	ຜ	4	>	O	24	×	œ	×	>	ល	ρι	z	ŀ
	301		Icag	art	הם +	caa	.099 	ctt +	tta	tot	tga	aaa ++	rag) Lgc	cacc +	JCac	Į,	1 d C	393	tto	tgcagatttatcaacggcttttatcttgaaaatagtgccacgcagtctacaaattctggt +++	360
rd		ပ	æ	[t4	H	Z	O	ſz,	>+	ı	M	Z	Ŋ	K	E+	O	Ŋ	E	Z	ഗ	O	•

2/67

	420	•	480	1	540	ı	900	1	099	1	720	1	
	tta +	,	ata +	н	gtcc	ഗ	gaactggccagactatgctcacctaaccccaagagagttagcaagtgctggactc 	LI.	cacaggtattggtgaccaagtgcagtgcttttgttgtggtggaaaactgaaaat	z	accttgtgatcgtgcctggtcagaacacaggcgacactttcctaattgcttcttt	[t+	
	gcc	K	gat	Ω	aag	×	gga	o	aaa	×	tto	[z ₄	
	tt	[Le	gta	>	gatta	h	gct	«	ctg	្ន	tgc	U	
	tcat	H	ggttg +	>	aga -+-	αú	agtg	S	aaa +-	×	ctaatt	z	
	gat	Ω	cag	o	gct	æ	gca	Ø	gga	G	TCC!		
	aga:	or,	999	Ģ	Igaa	ធា	gtta	ı	rggt	O	Ctt	[EL	
	gaage +	S	gaactg +	E	tgaa -+	ធ	aga; -+-	Œ	rtgl -+-	ပ	a Ca + +	æ	
	9998	Ö	gage	¤	ttgt	U	88 G	α ;	ttg	O	gcg	œ	
kiap	toto	J	ו ו ו ה	h	gta	> -	00 1	۵	ctt	Ĭ1,	cag	ø.	
HUMAN xiap	actal -+	>	atct	H	gccat	X	ctaac	H	agtg -+	U	aca.	Ħ	
$\overline{\mathbb{Z}}$	333	Z	cta	>	ccctgo	ccctgo	4	CCT	J	gca	œ	3agaa +	M
I	tga 	网	aga	Ω			000	Δ.	tca	æ	lagt	>	g to
	aaagt -+	>	tgc	A . H	gaa	Z	tgo	4	acca	O	gcctg	3	
	,+	×	acat	Ħ	gag	£	actat	>	atga 	Q	atgo	4	
	gaatggtcagtacaaagttgaaaactatctgggaagcagagatcattttgcctta ++++++	>	ıgac	PSETHADYLLRTGOVVDI caccatatacccqaqqaaccctgccatgtattgtgaagaagctagattaaagtcc	000	<u>Ω</u>	cage	Ω	ttgg	O	a I C	ĸ	
		O ctga	tga		caté	> +	9900	C4	gta	H	gtg	Ω	
	atgg +-	Ö	cato		ccat	Н	actg +-	3	20 + 20 +	Ŋ	cttg +-	O	
	ıgaçı	z S) jgc	ል	aca(Ħ	aga	Z	aca	€		Ωı,	
	atcca	ø	gacag	æ	tcaga	Ω	tttca	O)	tacta	>-	tggga	្រ	
n		Н		Ω		ທ		Ĭ24		>		3	
<u>.</u>	361		421		481		541		601		661		
Ĺ		rd		๗		ø		ø		rd	•	·	

3/67

1080 1020 960 840 900 780 +1111 ctgttagaacagaagggacaagaatatataaaacaatattcatttaactcattcacttgag ttcccaaattcaacaaatcttccaagaaatccatccatggcagattatgaagcacggatc tatgetttaggtgaaggtgataaagtaaagtgettteaetgtggaggagggetaaetgat tggaagcccagtgaagacccttgggaacaacatgctaaatggtatccagggtgcaaatat gttttgggccggaatcttaatattcgaagtgaatctgatgctgtgagttctgataggaat tttacttttgggacatggatatactcagttaacaaggagcagcttgcaagagctggattt w D, J æ 民 S D Ö I O 릹 S ۵, H 4 G S L Ö Ω J > 工 U 3 O 4 ď × 回 I Ξ 4 2 [to × S S **HUMAN xiap** I Z U Ω, Z Θ O Н × S Z 回 > S K 民 ω × 3 Q, >4 p, O a J Z Ö 3 Ö Z H 团 × 臼 H H Z Q S Ü 民 S M Ö J 3 u FIG. 1C 901 961 1021 841 781 d đ đ Ø đ đ


4/67

	1140	•	1200	1	1260	•	1320	ı	1380	,	1440	•
FIG. 1D HUMAN xiap	gagtgtctggtaagaactactgagaaaacaccatcactaacta	ECLVRITEKTPSLTRRIDDT -	atcttccaaaatcctatggtacaagaagctatacgaatggggttcagtttcaaggacatt	I F Q N P M V Q E A I R M G F S F K D I	aagaaaataatggaggaaaaaattcagatatctgggagcaactataaatcacttgaggtt 	KKIMEEKIQISGSNYKSLEV-	ctggttgcagatctagtgaatgctcagaaagacagtatgcaagatgagtcaagtcagact	LVADEVNAQKDSMQDESSQT -	tcattacagaaagagattagtactgaagagcagctaaggcgcctgcaagaggagaagctt	S L Q K E I S T E E Q L R R L Q E E K L	tgcaaaatctgtatggatagaaatattgctatcgtttttgttccttgtggacatctagtc	CKICMDRNIAIVFVPCGHLV -
FIG	1081		1141		1201		1261		1321		1381	
		ત્વ	·	rd		M	•	ď		ď		ď

5/67

	ttact + 1500	! €+	gttct + 1560	ı	tccat + 1620	ı	atata + 1680	ł	tgtta + 1740	ı	tgtgt + 1800	•
HUMAN xiap	acttgtaaacaatgtgctgaagcagttgacaagtgtcccatgtgctacacagtcattact 	CKQCAEAVDКСРМСҮТVІ	ttcaagcaaaaaatttttatgtcttaatctaactctatagtaggcatgttatgttgttct 	K Q K I F M S *	tattaccctgattgaatgtgatgtgaactgactttaagtaatcaggattgaattccat 		tagcatttgctaccaagtaggaaaaaaatgtacatggcagtgttttagttggcaatata 		atctttgaatttcttgatttttcagggtattagctgtattatccatttttttt		tttaattgaaaccatagactaagaataagaagcatcatactataactgaacacaatgtgt 	
FIG. 1E	ac 1441	E+	tt 1501	E.	ta 1561	-	tag 1621		a1 1681 -		1741	
ш		๙		ď		ď		Ø		æ		Œ

١



```
2520
                                                                                     2280
                        2220
                                                                                                                                                     2340
                                                                                                                                                                                                                  2400
                                                                                                                                                                                                                                                                              2460
             nagatatgttaagtgtaaaatgcaagtggcnnnacactatgtatagtctgagccagatca
                                                                          aagtatgtatgttnttaatatgcatagaacnanagatttggaaagatatacaccaaactg
                                                                                                                                         gtnetetteetagggagetgtnttgttteecacecaceacetteectetgaacaaatge
HUMAN xiap
                                                                                                                                                                                                                                                                                                                                                                                                      2540
                                                                                                                                                                                                                                                                                                                                                                                           ctgagtgctggggcacttn
                                                                                                                                                                                                                                                                                                                                                                                                      2521
                           2161
                                                                                                                                                                                                                                               Ø
                                                                                                                                                                                                                                                                                                           đ
                                                                                                                                                                                                                                                                                                                                                                      đ
                                                                                                                       ď
                                                                                                                                                                                   Q
                                                        ಗ
```

360 180 240 300 120 9 **ACACGTTTGAACTGAAATACGACTTGTCATGTGAACTGTACCGAATGTCTACGTATTTCCA CITITICCIGCIGGGITCCIGICTCAGAAGGAGTCITGCICGIGCIGGITICTATTACA** atticattatgaacatagtagaaacagcatattcttatcaaatttgatgaaaagcgcca **TCCTTGAGATGTATCAGTATAGGATTTTAGGATCTCCATGTTGGAACTCTAAATGCATAGA** GCAAAGCCATGCACAAAACTACCTCCCTAGAGAAAGGCTAGTCCCTTTTCTTCCCCATTC aatggaaataatggaaatttttcattttggcttttcagcctagtattaaaactgataaaa ហ Œ, ທ Σ O Ч E 4 œ. Z æ ល 4 J J **HUMAN hiap-1** 凶 ທ ĵ, ပ ø, ທ W ហ u Z S Ω u U Z [Ŀ, 301 181 19 121 Ų U U O U υ

720 999 480 540 600 420 ŧ Σ H **ATTCAAACTCTCCATCAAATCCTGTAAACTCCAGAGCAAATCAAGAATTTTCTGCCTTGA** S H CACATTCCACACACTCATTACTTCCGGGTACAGAAACAGTGGATATTTCCGTGGCTCTT ď GTCTAAATTCCGTTAACAACTTGGAAGCTACCTCTCAGCCTACTTTTCCTTCTTCAGTAA CTGGTGTGAATGACAAGGTCAAATGCTTCTGTTGTGGCCTGATGCTGGATAACTGGAAAA GAGGAGACAGTCCTACTGAAAAGCATAAAAAGTTGTATCCTAGCTGCAGATTCGTTCAGA L O S > × O d Œ, G S 3 > E S œ Ŀ S Z H [te (z, A K p, (L) ... \rightarrow O Œ O K Ö E Σ ທ 4 z S ρ, Ļ ∢ Z Z 0 Q J 出 囟 **HUMAN hiap-1** S (c) U ********************* ហ Z × E H U O Z 4 Z [z, × Σ U 囟 I × ۵, Ω, L Z O > Z 回 × Z Ø ۵, H > Ω <u>ρ</u> I Q, >4 ທ H Z ហ S ហ Z ហ S Ω Z U O ល K FIG. 2B 541 601 199 481 361 421 U U U U υ U

	780	ı	840		006		960		1020		1080	,
		Ö		×				œ		ຸທ		Ö
	ZY.	н	ACCGA	٥,	AAATC	Z	AGCCC	4	201	4	TGG.	Ö
	CTAC	т х х	SGE	ш	AGA	ធ	13CC	4	SCT	a	TGA	۵
	TTCTGTCGCCAACAGATCTGGCACGAGCAGGCTTTTACTACATAG	>-	PATTG	3	TAT	н	CACA	x	TAACTGGCCCTCTAGTTCTAGTTAATCCTGAGCAGCTTGCAA	O ⁴	GTTTTTATTATGTGGGTAACAGTGATGATGTCAAATGCTTTTGCTGTGATGGTG	ບ
	CTT	[z ₄	S.	Z	ATT	[II4	GAC	H	TGA	凹	TTG 	ບ
	99	O	3AG	ເນ	X	D.	SCA.	O	1 1 1 1	Ω,	CHI	[z,
	, GC	g 4	ATT	ı	ATG.	ပ	CAT	×	TAA	Z	ATG	Ö
	CG2	œ	*	×	3	×	GAGCAT	ທ	AGT.	>	CAA	×
HUMAN hiap-1	200	K	GGTGGAAATTGAG	F A C G G K	BAGACATTTTCCCAAATGCCCATTTAT	ρι	ICI	ı	TCT	1	TGT	>
iğ Di	Ş	ם	ည်	Ö	i i	[24	FA	Z	IGT	>	IGA	Ω
MAR	GAT	Ω	13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	ပ	CA +	Œ	TTTCTAA	Ŋ	rAG.	ശ	rg. +	Ω
Ξ	AC.	E	TGCCTGT	4	3AG	œ	AGT	>	S i	ഗ	CAG	Ŋ
	Ü	O.	CTT	(z4	S. C.	J	'AC'	E	Ω Ω	Δ,	PA .	Z
	FTCG	S	SCTTGC	ပ	AGAACACCT		GATAC	>	55 1	3	3661	G
	CTC	L	ည်း	V A C	AGA	យ	AAG.	æ	TAA	Z	TGT	>
	LTI	[tr	AGT	>	GTC	S	TTC	Ø	CTT	[z4	TTA	>
	3AC	₽	CAG	œ	ATGCTAT	×	CAC	٤	ATT	ſz,	TTA	>-
	TT.	H	AGA.	Д	ည်း	4	AGA ++	Ω	AAC ++	₽	TTT +	(z.
	<u> </u>	Ω,	79	Ö	4		Ç	OI	3	×	99	Ö
	CATGGCCATTGACTTTTCTGTCGCCAACAGATCTGGCACGAGCAGGCTTTTACTACATAG ++++++	3	GACCTGGAGACAGAGTGGCTTGCTTTGCCTGTGGAAAATTGAGCAATTGGGAACCGA	G G	AGGATAATGCTATGTCAGAACACCTGAGACATTTTCCCAAATGCCCATTTATAGAAATC	2	AGCTTCAAGACACTTCAAGATACACAGTTTCTAATCTGAGCATGCAGACACATGCAGCCC	ы О	GCTTTAAAACATTCTTTAACTGGCCCTCTAGTGTTCTAGTTAATCCTGAGCAGCTTGCAA	ŗ.	GTGCGGGTTTTTATTATGTGGGTAACAGTGATGTGATGT	4
FIG. 2C	CJ 721		G/ 781 -		A 1 A		A 901 -		961 -		G 1021 -	
FIG		U		U		υ		U		U	• •	U

	1140) 		1200		,	1260))	t	1220	3	,	1380		ı	7	444	1
		• .	C			>			(c)			Z	8 †		H	GT.	-	IJ
	JOI 1		വ	CAG		ល	TGC		K	GAT		Σ	ACA		o	TGT		ר
	TI		(z.	AGC		æ	A P		z	CAT	 	Σ	A:		×	TCT	} } 	J
	37.0	_	X X F	ICA.		O	rGA	•	ចា	AAT	} 	н	3GT		L V K	IGA	} -	Ω
	CAA(AGT		0 > 0	AGA'		Ω	TGC) 	æ	CCT		L	CAA		z
	J J		4	ICA.			AGG		ဗ	AGA.	} !	Ω	PAG		ທ	AGT		>
	AGGTGTTGGGAATCTGGATGATCCATGGGTTCAACATGCCAAGTGGTTTCCAA		Ħ	GAGTACTTGATAAGAATTAAAGGACAGGAGTTCATCCGTCAAGTTCAAGCCAGTT		œ	NTCTACTTGAACAGCTGCTATCCACATCAGACAGCCCAGGAGATGAAAATGCAG		O.	TCAATTATCCATTTGGAACCTGGAGAAGACCATTCAGAAGATGCAATCATGATGA		យ	CCTGTGATTAATGCTGCCGTGGAAATGGGCTTTAGTAGAAGCCTGGTAAAACAGA		œ	ACT		ı
	S t	•	O	CAT	•	ы	CAG	•	ഗ	TTC	-	S)	TAG	•	ഗ	TAG		æ
-	GT		>	STT		ſe,	AGA(O.	CCA) 	æ	CHI		[e,	TTA	 	>
lap-	ATG(3 Q	GGA		闽	ATC	 	Ŋ	AGA		Ω	999		O	GAA		Z
Š	5	-	Ω,	ACA(-	O ³	CAC	; -	۴	AGA) - 	Гī	AAT		Σ	AGA	 - 	ា
HUMAN hiap-1	IGA.	 	Ω	AGG		Ö	ATC	 	Ω	TGG	i I !	ဗ	GGA		М	TGG) } 	O
Ī	AGA.		Ω	raa		×	SCT	l !	u u	ACC	 	ο.	CGT		A V	AAC]]]	۲
	TGG	 -	Ö	AAT	-	H	GCT	 - -		GGA] -	ជា	TGC		K	AGC	<u> </u>	æ
	ATC.] ;	ល	AAG	 - 	∝	ACA] }- }	ø	TTT]]]	J	TGC		æ	CCT		ı
	GGA	 	េ	GAT) ! 	Н	TGA) } .	(m)	ည	} }	x	TA		Z	AAT		×
	TTG) }	3	CTT] }	H	ACT		J	TAT	; 1 1	Н	GAT		V I V	A A		
	FIG.	} ! !	ပ	GTA	} 	>	TCT	- -	J	AAT	+ 	н	TGI	• 	>	GAG.	t 1 1	∝
	CAG.))	œ	rga		凹	ICA	•	Ħ	ATC		Ŋ	700	} }	ρι	TCA		O
	GACTCAGGTGTTGGGAATCTGGATGATCCATGGGTTCAACATGCCAAGTGGTTTCCAA	; ; ; ;	ı	GGTGTGAGTACTTGATAAGGATTAAAGGACAGGAGTTCATCCGTCAAGTTCAAGCCAGTT	 	ပ	ACCCTCATCTACTTGAACAGCTGCTATCCACATCAGACAGCCCCAGGAGATGAAATGCAG	1 1 1	D.		† † †	တ	ATACTCCTGTGATTAATGCTGCCGTGGAAATGGGCTTTAGTAGAAGCCTGGTAAAACAGA) 	Ę÷	CAGTTCAGAGAAAATCCTAGCAACTGGAGAGAATTATAGACTAGTCAATGATCTTGTGT	! ! !	>
FIG. 2D	_	1801		***************************************	777		6	1071	•	. (1921		127	1761		,	1961	
F		•	U			υ			O			U			U			U

	1500	1	1560	ı	1620	l Fa	۱ ۲ 1680		1740	į fai	1800	ا
	AGG	ា	ATT +	,J	AAG	闭	TGATA	H	CTC	M	AAG	C
	TG	ឲា	\$:	I	\$!	OI.	TT D	Ω	AAG -	Æ	ZAG.	. 6
	AA.	€	Ú.	O.	VTG.	ធ	GA	н	AG.	回	CAC	Ε
	AGC +	4	1111	ÇL,	TA	Z	S t	ı	Ω +	O	TCC	C
	AAG	æ	ACT	h	TAT	н	AGA	ធា	TCT	H	TAT	۲
	AGA	ធ	ည္ဟ	K	AAT	Н	AAG	æ	OTO:	လ	ATA	>
	GAG	æ	AAT	×	990	Ö	AGC	K	A :	Z	AA :	>
	GGA +	ш	TAG +	œ	13C	K	ACA - + +	a	CAG ++	ps;	CAT	1-
Ţ	AGA	回	GAA	Z	PAC	E٠	TTT	.	ATT	Ee,	SGA	6
hiap	3GA	ص	SAA	×	ACT	H	GTC	Ø	TGT	>	ACA	C
A V	AAG.	oc.	,	œ	ICT.	J	GAC	e	CAC	€·	3C.A.	c
HUMAN hiap-1	actcaatgcagaggtgaaataagggaaggaggagagagaaagagcaactgagg	H	AATCAAATGATTTATTAATCCGGAAGAATAGAATGGCACTTTTTCAACATT	н	GTGTAATTCCAATCCTGGATAGTCTACTAACTGCCGGAATTATTAATGAACAAG	Ø	atgttattaaacagacacagacgtctttacaagcaagaactgat	a	TAGTAAAAGGAAATATTGCAGCCACTGTATTCAGAAACTCTCTGCAAGAAGCTG	æ	TGTTATATGAGCATTTATTTGTGCAACAGGACATAAAATATATTCCCACAGAAG	2
I	164	阳	ATT	ם ניני	459 1	Ω	GAC	E E	13C	æ	ATT	£
	AGA.	Ω	ATT	H	Į Į	H	GAA	×	TAT +	н	TTT	ŗ.
	NGA.	ы	L	J	AAT	н	ACA	O	AAA.	Z	SCA.	=
	ည်	4	rGA	Ω	2	Ωι	FA	×	200		rga((a
	A.	Z	ZA S	Z	¥.	н	rAT	н	3	×	ATA	>
	CTC	H	¥4.	Ø	rgt)	>	ĘĘ †	>	KGT2	>	3TT2	نو
	TT	ı	GA	E	TG	Ü	GA	Ω	TL	L)	GTG	٥
	TAGACTTACTCAATGCAGAAGATGAAATAAGGGAAGAGAGAG	D	AAAAAGAATCAAATGATTTATTAATCCGGAAGAATAGAATGGCACTTTTTCAACATT	⋈	TGACTTGTGTAATTCCAATCCTGGATAGTCTACTAACTGCCGGAATTATTAATGAACAAG	E4	AACATGATGTTATTAAACAGAGACACAGACGTCTTTACAAGCAAG	Ħ	CGATTTTAGTAAAGGAAATATTGCAGCCACTGTATTCAGAAACTCTCTGCAAGAAGCTG	н	AAGCTGTGTTATATGAGCATTTATTTGTGCAACAGGACATAAAATATATTCCCACAGAAG	4
רומ. גר	1441		1501		1561		1621		1681		1741	
ב		U		U		υ		υ		υ		•

	rA -+ 1860	X I	AT -+ 1920	ט	TA -+ 1980	 	FAAT + 2040		TT -+ 2100	- CATA + 2160	1
	VTG.	ပ	AGTAT	>	3GGTA	O	FTA	·	LTA	CCA	
	YAC.	E	AGT	>	CAA(*	SAA		TTT	AAA :	
	YAG.	œ	rcr	ы	racaatc	Н	rta(A A	TATCT	
	ATGTTTCAGATCTACCAGTGGAAGAACAATTGCGGAGACTACCAGAAGAAGAACATGTA	M	aagtgtgtatggacaaagagtgtccatagtgtttattccttgtggtcatctagtagtat	р С С н г ∨	GCAAAGATTGTGCTCCTTCTTTAAGAAAGTGTCCTATTTGTAGGAGTACAATCAAGGGTA 	E	CAGTTCGTACATTTCTTTCATGAAGAAGAACCAAAACATCGTCTAAACTTTAGAATTAAT		TTATTAAATGTATTATAACTTTTAACTTTTATCCTAATTTGGTTTCCTTAAAATTTTTATT	tatttacaactcaaaaacattgttttgtgtaacatatttatatatgtgtaaccata	
	AGA	田	7GG	O	GAG	ល	TAA		100 111	TAT	
	ACC	ρ,	TTG	U	CCTATTTGTAGG	D.	AACATCGTC		GTT	PATA	
	ACT	J	CATAGTGTTTATTCCTT		TTG	U	ATC		rrrg +	\TT\ +	
y-	GAG	æ	TAT	Н	TAT	H	NA AC	(,	FAA	CAT	
HUMAN hiap-1	909	改	GTI	> 당	STCC	D Tr	3		ည်	rgtaacat -+	
Z Z	ATT	7	AGT	>	AAGTGT	ບ	PAG +		rta.	3TG1	
MΣ	ACA	O	CAT		A .	×	AAC		FF	TTC	
Image: section of the	AGA -	M	GTC	တ	AAG	œ	AAG		AAC	GFT	
	8 ‡	M	AGT +	>	TTT ++	,a	ATG +	*	TTT	ATT +	
	GT	>	AGA	២	PTCTT	ഗ	된 건	ທ	PAC	AAC .	
	Ö	۵	8	×	ည်	Ω,	ICT -	ы	TAT	\$:	
	Č.	ı	3GA	Ω	ည် <u>i</u>	K	ATT	(14	rat	Ž.	
	CAGATCTACC	Ω	ATC	E	DI +	· U	TAC:	E	VIG.	AC.	
	TTCA	Ø	TGI	υ	GAT	Ω	ည်	œ	3	S :	
	TTE	S	AAGTGT	>	\$:	N D	CAGTTC	>	ATT	7 T T T	
	ATC		A :		8		5	•	FF	TA	
FIG. 2F	1801		1861		1921		1981		2041	2101	
FIC		O		U		U		Ü		υ	U







FIG. 2H



SUBSTITUTE SHEET (RULE 26)

16/67

	09		120	ı	180	1	240	1	300	1	360	1
		• .		•								·
	17GG +		AAGAAAG		AGTT +		CATG	Σ	TAT	H	GATCTTGTCAGATTGGACAACAGCAACAAAAAAAAATGAAGTAT	>
	ATC.		AAG		CTT		ACT.		GAG	ഗ	SAA	×
	/GT/		STA.		YAT(1 C		IAA	×	AAT	X
	aagtagt +		CA		GGAAA		ICA		CAT	H	A T	×
	TX		AAACTGAGATAAATCCAGTA		3GTC		DI-C		AA.	z	AC :	ø
	Caaccccaaagagttgtgttct		TA		rgre		STAC		Ž i	O	A	×
	STS.		AG.		PPPIG		CAC		STAT	>	AAC +	Z
Ŋ	GT.		CTO		Ş		A I		Ď.	ß	7.AG	S
HUMAN hiap-2	AG.		Z. Z.		TT.		FATC		ည်	۵,	A A	z
N N	2		TGAATAT		rīg.		55		99	O	3AC	E
₹	CCC		GAZ		Ş İ		LTT		77)	Ωι	FTG	3
Œ	3		ည်		TA		IGT		ITT	[z,	AGA	Ω
	SAGTTACTA		Zi j		AGT		SA :		ACT	u	GIC	ເນ
	STT		ACTCATCCT		AAG.		TGT		AAG.	œ	CTT	ı
	AGAC		ACT		EAT :		TCA		25	ø	GAT	H
	AAS		SAT		CTA		ATT		CHO CHO	လ	CAC	€-
	CCTGAA		AGA(ATT -+-		GAA		TGC -+-	4	AGATAGCAC	ຜ
	ITA		CAC		E :		GAA		AAC 	H	AGA	Ω
	TTAGGTTACCTGAAAGAGTTACTACAACCCCAAAGAGTTGTGTTCTAAGTAGTATCTTGG		Taattcagagagatactcatcctacctgaatataaactgagataaatccagtaaagaaag		TGTAGTAAATTCTACATAAGAGTCTATCATTGATTTCTTTTTGTGGTGGAAATCTTAGTT ++++		CATGTGAAGAAATTTCATGTGAATGTTTTAGCTATCAAACAGTACTGTCACCTACTCATG		CACAAAACTGCCTCCCAAAGACTTTTCCCAGGTCCCTCGTATCAAAACATTAAGAGTATA	×	ATGGAAGATAGCACGATCTTGTCAGATTGGACAACAGCAACAACAAAAAAAA	ធ
3A			•				5 !		5 !	Œ		Σ
FIG. 3A	· ન		61		121		181		241		301	
		ď		ď	•	ø		๙		ď		đ

9 720 540 600 420 480 **AAATGCTTCTGTTGTGGCCTGATGCTGGATAACTGGAAACTAGGAGACAGTCCTATTCAA** aagcataaacagctatatcctagctgtagctttattcagaatctggtttcagctagtctg GTCTCAGAAAGGAGTCTTGCTCGTGCTGGTTTTTATTATACTGGTGTGAATGACAAGGTC **ACCTTGGAACATAGTAGCTTGTTCAGTGGTTCTTACTCCAGCCTTCCTCCAAACCCTCTT** GACTTTTCCTGTGAACTCTACAGAATGTCTACATATTCAACTTTCCCCGCCGGGGTGCCT ᆸ Q, > O ρı ល 노 S Z O S Z, Z W. S 工 م D, Ω ď, J ĵu, G Ö H Z [z, S 5 E **HUMAN hiap-2** 0 S S S × \succ 3 z ≯ E Ĺ, Z [t, 叱 S ຜ S G Ω \mathbf{z} O U D, Z Z, J ഗ ហ \mathbf{z} ĸ æ ഗ Œ, Ω, Æ, J E J Ö >4 J J S Z J U ഠ វ × S O O U 吖 S 二 × W ທ (z, 回 H I Œ, Ŋ S J Ω O FIG. 3B H 361 541 481 661 601 đ Q đ đ đ Ø **SUBSTITUTE SHEET (RULE 26)**

	AGAGCAGTTGAAGACATCTCTTCATCGAGGACTAACCCCTACAGTTATGCAATG	X	AGTACTGAAGAAGCCAGATTTCTTACCTACCATATGTGGCCATTAACTTTTTTGTCACCA	S S	TCAGAATTGGCAAGAGCTGGTTTTTATTATATAGGACCTGGAGATAGGGTAGCCTGCTTT	EL U	GCCTGTGGTGGGAAGCTCAGTAACTGGGAACCAAAGGATGATGCTATGTCAGAACACCGG	H	AGGCATTTTCCCAACTGTCCATTTTTGGAAAATTCTCTAGAAACTCTGAGGTTTAGCATT	s H	TCAAATCTGAGCATGCAGACATGCAGCTCGAATGAGAACATTTATGTACTGGCCATCT	
	CAGTT	S	TTT.	F	GTAG	A	TCAG	ល ធា	AGGT	ir.	TACTO	3
	CTAC	≯	AACT	€-	TAGG	œ	TATG	Σ	TCTG	h	TATG	×
	CCC	D.	GTGGCCATT	ı	AGA	Ω	FEG	¥	AAC	H	ATT	[t.
	TA.	Z	ည်း	ρı	TGC	Ċ	TG.	Ω	AGA -+-	យ	AAC	. E4
HUMAN hiap-2	rtcatcgaggactaac	٤٠	GTO	3	ACC	ים	655	Ω	TCT	ı	TCGAATGAGAACATT	æ
his	GAC	æ	TAT	Σ	'AGG	Ø	AS.	×	TTC	ល	AAT	Z
MA	ATC +	ഗ	D D	**	TAT	H	ACC	Д	AAA +	Z	TCG	œ
∃ H	TT.	Ø	Č.	>+	TT	×	999	回	669	ជា	AGC	æ
	SC FC	ຜ	TAC	ۥ	TT	>	CTG	3	TTI	J	TGC	A
	CA	Н	TTTCTTACCTACCAT	H	TTT	(EL	TAT	Z	CCATT	ŢĿ	ACA	. 33
	AGAC	Ω	SAT	(Le	TGC	O	CAC	S	77.00	щ	GAC	E
	rrg.	凹	5	œ	3AGC	4	GCJ	ı	CTG	U	S	Q
	AG	>	AGG	K	AAG	ద	GAA	×	TCCCAACT	z	CAT	E
	AGC	A	AGA	េ	TTGGCAAGAGCTGGTTTTTATTATATAGGACCTGGAGA	4	TGG ++	Q	700	Ω	GAG	Ŋ
	TAC	æ	TG	ធា	ATT	ч	TGG	Ö	TTT 	Œ	TCT	J
45	AATTCT	ល	TAC	64	AGA	ω	GCCTGT	O A	AGGCAT	R	AAA	Z
30		Z		ທ		ß		4	-	æ		
FIG. 3C	721		781		841		901		961		1021	
		๙		ಹ		ď		ぱ		Ø		Ø

	1140	1	1200		1260	1	1320	1	1380	•	1440	•
FIG. 3D HUMAN hiap-2	AGTGTTCCAGTTCAGCCTGAGCTTGCAAGTGCTGGTTTTTATTATGTGGGTCGCAAT		GATGATGTCAAATGCTTTGGTTGTGATGGTGGCTTGAGGTGTTGGGAATCTGGAGATGAT	D D V K C F G C D G G L R C W E S G D D	CCATGGGTAGAACATGCCAAGTGGTTTCCAAGGTGTGAGTTCTTGATACGAATGAAAGGC		CAAGAGTTTGTTGATGAGATTCAAGGTAGATATCCTCATCTTCTTGAACAGCTGTTGTCA	Q E F V D E I Q G R Y P H L L E Q L L S	ACTICAGATACCACTGGAGAAGAAATGCTGACCCACCAATTATTCATTTGGACCTGGA	TSDTTGEENADPPIIHFGPG	GAAAGTTCTTCAGAAGATGCTGTCATGATGAATACACCCTGTGGTTAAATCTGCCTTGGAA	ESSSEDAV M M N T P V V K S A L E
Ĭ	1081		1141		1201		1261		1321		1381	
		, ø		๙		๙		๗		๙		ď
					CHECTI	T. T.	CHEET	. /	- 001			

20/67

	1500	i	1560	1	1620		1680		1740		1800)) 1
FIG. 3E HUMAN hiap-2	ATGGGCTTTAATAGAGACCTGGTGAAACAACAGTTCTAAGTAAAATCCTGACAACTGGA	MGFNRDLVKQTVLSKILTTG -	GAGAACTATAAAACAGTTAATGATATTGTGTCAGCACTTCTTAATGCTGAAGATGAAAAA 1501+++1	ENYKTVNDIVSALLNAEDEK-	AGAGAAGAGGAAAAAAAAAAGCTGAAGAAATGGCATCAGATGATTTGTCATTAATT	REEKKOAEEMASDULSLI-	CGGAAGAACAGAATGGCTCTTTCAACAATTGACATGTGTGTTCCTATCCTGGATAAT	RKNRMALFQQLTCVLPILDN-	CTTTTAAAGGCCAATGTAATTAATAAACAGGAACATGATATTATTAAACAAAAACAGG	L L K A N V I N K Q E H D I I K Q K T Q -	ATACCTTTACAAGCGAGAGAACTGATTGATACCATTTGGGTTAAAGGAAATGCTGCGGCC	IPLQARELIDTIW VK GNAAA
		ಹ		ø		ø		ಡ		ಡ		ಡ

FIG. 3G	39	HUMAN hiap-2	
, •	2161	AICIAAAGIAAAAGGGAAIIAIGAGIIIIICAAIIAGIAACAIICAIGIICIGC	2220
rd			1
···	2221	TTTGGTACTAATAATCTTGTTTCTGAAAGATGGTATCATATATTTAATCTTAATCTGTT	2280
ø			1
• • •	2281	TATTTACAAGGGAAGATTTATGTTTGGTGAACTATATTAGTATGTAT	2340
๗			•
	2341	AGTAGCGTCXCTGCTTGTTATGCATCATTTCAGGAGTTACTGGATTTGTTGTTCTTTCAG	2400
æ			ı
	2401	AAAGCTTTGAAXACTAAATTATAGTGTAGAAAAGAACTGGAAACCAGGAACTCTGGAGTT	2460
୍			•
	2461	CATCAGAGTTATGGTGCCGAATTGTCTTTGGTGCTTTTCACTTGTGTTTTTAAAATAAGGA	2520
ø			•
	2521	TTTTTCTCTTATTTCTCCCCCTAGTTTGTGAGAAACATCTCAATAAAGTGCTTTAAAAAG	2580
ಡ			.

	GACACTCTGCTGGGGGGGGGCGCCCTCCTCGGGAACCGTCGCCCCCCCC	GCGGCGCTTAGTTAGGACTGGAGTGCTTGGCGCGAAAAGGTGGACAAGTCCTATTTTCCA	•	GAGAAGATGACTTTTAACAGTTTTGAAGGAACTAGAACTTTTGTACTTGCAGACACCAAT	, S E O	AAGGATGAAGAATTTGTAGAAGAGTTTAATAGATTAAAAACATTTGCTAACTTCCCAAGT	(i)	AGTAGTCCTGTTTCAGCATCAACATTGGCGCGAGCTGGGTTTCTTTATACCGGTGAAGGA	រ ២ ធ	GACACCGTGCAATGTTTCAGTTGTCATGCGGCAATAGATAG	S Q D
	GGGAA + -	AGTO		rrg(4	CTA	Z	ATA(€4	AGT.	>-
) TC	- K		FAC	h	AAAAACATTTGC +	æ	rtt	>	360	ø
	Ö	7997		rī.	>	AT	ŢŦ	rac:	J	3AT	3
	p ACCTCC	69		11.	ĬĿ,	¥ t	E	3GT	[t+	ATA(œ
	dap GAC	4		PAG	£ +	A .	×	TG	O	PAG2	۵
	SE S	600		TAC	œ	ATT	J	AGG	K	A .	H
	MOUSE xiap rccrccggac)) +		PAAC +	F	Taatagatt +	æ)) (œ	99	4
	CTC	TTG		98		Ž.	Z	999	4	73	4
	ည္ဟ	7GC		TGA	យ	GTT	<u>(r</u>	ATT	.ı	ភ្ជ	×
	၁၁၅	GAG		TTT	Œ	AGA	凶	AAC.	H	TTC	ပ
	990	TAGGACTGGAGTGCTTGGCGCGAAAAGGTGGA(CAG	Ŋ	TGTAGAAGAGT	(c)	ATC	ທ	5	S
	99	GGA		TA.	Z	TGT	>	AGC	4	TTT	(z,
	999	TTA		TTT	(z.	ATT	ĵz,	TTC	Ŋ	ATG	ပ
	GCT -+-	TAG -+-		GAC	€→	AAGAAT	ω	TGT -+-	>	₹ 1	O
	ICT 	GCT		GAT	E	7G.	M	70C	ρı	CGT	>
	CAC	Ö i		GAA		GGA	N D	TAG	ß	CAC	64
_	GAC	ပ္ပ	•				×		Ŋ	S :	Ω
FIG. 4A	H	61		121		181		241		301	
		d	๗		đ		a		ં ત		ં વ

	420	1	480	4	540	1	900		099	•	720	
	GCTGTTGGAAGACACAGGAGAATATCCCCAAATTGCAGATTTATCAATGGTTTTTTATTTT 	Í24	GAAAATGGTGCTGCACAGTCTACAAATCCTGGTATCCAAAATGGCCAGTACAAATCTGAA	យ	AACTGTGTGGGAAATAGAAATCCTTTTGCCCCTGACAGGCCACCTGAGACTCATGCTGAT	Ω	TATCTCTTGAGAACTGGACAGGTTGTAGATATTTCAGACACCATATACCCGAGGAACCCT	Q,	GCCATGTGTGGAGGAAGCCAGATTGAAGTCATTTCAGAACTGGCCGGACTATGCTCAT	æ	TTAACCCCCAGAGAGTTAGCTAGTGCTGGCCTCTACTACACAGGGGCTGATGATCAAGTG	>
	E i	>-	TA!	Ŋ	LTG.	æ	20	Z	LTG(A	II.	O
	TT	[Li	3	×	Ş	Ħ	CCCGAGGAAC	œ	ET.	>	T.	۵
	13G 1 + 1	Ö	GTA +	>	GACTC	H	22 †	O.	GGA 	Ω A	TGA	۵
	CAA	Z	නු <u> </u>	ø	TGA	េ	TATA	>			ည်	4
	TTGCAGATTTATCAATGGTTTTT	н	GTATCCAAAATGGCCAGTACAAA	0 N G Q Y	GCCCCTGACAGGCCACCTGA	Ω,	CAT		ET C	N O	90	U
	ATT.	(tr	¥ ÷	Z)) (М	CAGACACC	H	3440	Z	CAC.	€
	. AG	æ	3	o	8	æ	AGA(Ω	5	O	TAC	>4
	ŢŢ ŎŢ	U	rat(ļН	ľĜĀ	Ω	Į.	တ	VET.	(z,	TAC	>-
ciap	CCAAA	z	9	I D	Si	ىم	GATATTT	н.	ST C	S	ST.	٦.
MOUSE xiap		Q,	rccrg	ω	Į Į	4	KGA:	IQAAÖD	AAAG	×	9	_o
ö	JJC.	S	A I	Z		ĬĿ,	GTA	>	TTG		CCT	<
Σ	AT.	н	SAC.	H	ည	Ω,	£5:	>	AGA	я г ж	AGT	S
	AGA:	æ	STCT.	N)	AAATCCT	Z	GGACAGGT	œ	AGCC	Ø	GCT	4
	AGG	æ	CAC	ø	'AG	民	9	Ö	GA.	Ю	TTA	니
	CAC	Ħ	ည	Æ	AAAT	z	S :	E	G. S.	M	GAG	ជ
	AGA	æ	Ω +	Ø	7555 +-	o	AG.	R.	AGT	S	AGA	吐
	3	O		Ġ	GTG	>	TTG	H	IGT.		22.2	Ωı
	GTT	>	AAT	z	TGT	U	Cid	,ı	CATG	5) V V	E
	J. J.	>	GAA	ы М	AACTGT	N O	TATCTC	r K	$\tilde{\mathcal{O}}$	N N	ŢĀ	1
	361		421		481	_	541	•	_	P.		
4B	36		4		4		20)	601		661)
FIG. 4B		Ø	٠	๙		Ø		ď		æ		ø

	780	ı	840	1 0	006	1	096	1	1020	ı	1080	,
	TTTTGTGGGGAAAACTGAAAATTGGGAACCCTGTGATCGTGCCTGGTCA	s S	GAACACAGGAGACACTTTCCCAATTGCTTTTTGGGCCCGGAACGTTAATGTTCGA	&	AGTGAATCTGGTGTGAGTTCTGATAGGAATTTCCCCAAATTCAACAACTCTCCAAGAAAT	Z	CCAGCCATGGCAGAATATGAAGCACGGATCGTTACTTTTGGAACATGGATATACTCAGTT	>	AACAAGGAGCAGCTTGCAAGAGCTGGATTTTATGCTTTAGGTGAAGGCGATAAAGTGAAG	×	TGCTTCCACTGTGGAGGGCTCACGGATTGGAAGCCAAGTGAAGACCCCTGGGACCAG	ø
	7G	3	GIT	>	AGA	œ	TCA	တ	GTG	>	GAC	_ ۵
)))	A	AAT	Z	CCA	۵.	TAC	≯	AA :	×	136	3
	CGT + +	œ	GTT	· >	ICT ++-	ഗ	ATA -+-	н	GAT	Ω	000	D,
	GAT	Ω	AAC	Z	AAC	z	13GG	3	ည္ဟ	O	GAC	Ω
	TGT		999	硂	ACA	Ę	ACA	E-1	8	凹	GAA	ГE
	ACCC -+	C) D4	000	. ტ	TCA	· v	GGA +	G	1561	ပ	LAG1	ဟ
	GAA	ធ	TTCCCAATTGCTTTTTTGTTTTGGGC		AAT	z	Lite	Ĩz,	TT	u	ည	Q
MOUSE xiap	7760	3	GTT	>	Ö	C4	5	E-4	ည်	FYAL	3AA(×
JSE	BAAAAT	Z	PTT	[E4	FFF	Œ	GE	>	rta • - •	> -	ITG F	3
ğ	3	×	111	(Zr)	SAAS	Z	3AT(н	ATT	(L4	SGA.	Ω
_	CTO	ı	TTGC	ပ	rage :	. 🕰	Š.	C	rgg.	O	P C	E
	3	×	A.	z	rga	Ω	20	4	AGC	æ	GCT	H
	36G.	U	Ω	Δ,	I'I'C		TGA.	េ	AAG.	∝.	AGG	O
	ည်း	v	TT	Ĺ	GAG	໌ທ	ATA	>	13G	4	AGG	ပ
	rtG)	ပ	ACA(**	rgr	S A S	AGA	ഥ	GCT	L	TGG	O
	TTTTGTGGGGGAAA.	ပ	3AG	R H F	rGG	b	360	æ	3C.A	œ	CTG	ပ
		(z4	:AG	æ	ATC	S	CAT	×	GGA	េ	უ :	X
	CAATGC	U	SCA CA	×	rga	回	AGC	4	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	×	CTT	Œ
O	CAATGC	ں ص	GA					P A			F F	Ö
FIG. 4C	721		Ç	18/	,	1 \$0	901		961))	1021	
		ď		๙		ø		ಡ		ď		Ø

WO 02/026968 PCT/CA01/01379

					27/6	57					
	1500	1	1560	1	1620	•	1680	•	1740		7 -
FIG. 4E MOUSE xiap	CAGCTAAGGCGCCTACAAGAGGAGAAGCTTTCCAAAATCTGTATGGATAGAAATATTGCT	O L R R L Q E E K L S K I C M D R N I A	ATCGTTTTTTCCTTGTGGACATCTGGCCACTTGTAAACAGTGTGCAGAAGCAGTTGAC	IVFFPCGHLATCKQCAEAVD	AAATGTCCCATGTGCTACACCGTCATTACGTTCAACCAAAAATTTTTATGTCTTAGTGG	KCPMCYTVITFNOKIFMS*	GGCACCACATGTTATGTTCTTCTTGCTCTAATTGAATGTGTAATGGGAGCGAACTTTAAG	のできます。 のである。これのこのできます。 ままれる これの	1681	AGCACT	1741
-		ಥ		Ø		ಗ		ď		đ	Ø



	1860	1920	, t)) 	2040	2100
4F MOUSE xiap	TAGCCAGTGTTTTACTCGATTGAAACCTTAGACAGAGAAGCATTTTATAGCTTTTTCACAT	GTATATTGGTAGTACACTGACTTGATTTCTATATGTAAGTGAATTCATCACCTGCATGTT	TCATGCCTTTTGCATAAGCAAATGGAGTGTTCTGTATAAGCATGGAGATGTGATG	GAATCTGCCCAATGACTTTAATTGGCTTATTGTAAACACGGAAAGAACTGCCCCACGC		CTGGGAGGATAAAGATTGTTTTAGATGCTCACTTCTGTGTTTTAGGATTCTGCCCATTTA
71 <u>6.</u> 47	1801	1861	1001		1981	2041
		ø	ď	Ø	rd	

	Ĭ	FIG. 5A	⋖								¥.	M-hiap-1	_									
	•	GAATTCCGGGAGACCTACACCCCCGGAGATCAGAGGTCATTGCTGGCGTTCAGAGCCTAG	ĎĮ:	ეეე ემ	GAG	:ACC	TA(ZACC	Ω Ω	,GG7	GA1	CAC	AGG	T.	TTG	CTC	ഉ	TTC	AGA	ညည	TAG	ć
	rd	GAAGTGGGCTGCGGTATCAGCCTAGCAGTAAAACCGACCAGAAGCCATGCACAAAACTAC	TG	360	+-+		CAT(+ 	, CI	VGC.	GT2	AAA	, D	ACC	. AGA	AGG	CAT	- - -	CAA	AAC	-+++++++) 9
	61	ATCCCCAG	; ; ;	AGA	++	AG2	*CT	+ IGT(TTC	T S	ည်	. E	S	÷ CT	.4C	'ATC	-+-	ATG	GTT	AGAAAGACTIGICCCTICCCCTCCTGTCATCTCACCAIGAACAIGGTICAA	120
ď	121	;	l !	1 1 1	+			+		į	Ĭ	1	i		+		Σ	<u> </u>	Σ	>	O A W N W	180
	; (_	₹GC	gac	TT.	ľCT?	JOS.	CTTTCTAGCCAAGCTGATGAAGAGTGCTGACACCTTTGA	3CT(3AT(3AA(3AG1	ည်	GA GA	34C(TI	GAC	TTC	AAG	TAT	CTTTCTAGCCAAGCTGATGAAGAGTGCTGACACCTTTGAGTTGAAGTATGAC	240
Ø	181	<u>.</u>	S		<u> </u>	i ii	4	<u> </u>	J	Σ	×	က	Æ	Q	- E-1	[E4	团	· H	×	⋈	Д	, 1 1
		TTTTCCTG	ıcc	TGI	GAC	3CT	GTA	מכפי	ATT(JTC(CAC	3TA)	T T	ີ່ປູ	TTJ	ນິ	'AGC	.GG7	GTJ	ີ່ວິ	TGAGCTGTACCGATTGTCCACGTATTCAGCTTTTCCCAGGGGAGTTCCTGTG	ć
ď	241	1 1 [24	S C		+ E	H	; > -	ELYRLSTYSAFPRGV	H	S	EH	<u>.</u>	တ	4	<u>;</u> + E4	Ωı	R	ָ ט	>	<u> </u>	+	0 0 1
	1		GAP	AGC	3AG	ICT	ວອອ	TCG	ĪĠĊ	ĪĞĞ	CIT	rta(CTA(JAC.	1991	ည်	AA	rga(AA(GT(TCAGAAAGGAGTCTGGCTCGTGCTTTTTACTACACTGGTGCCAATGACAAGGTCAAG	360
Ø	301	S	M	_	+ s	ļ	A	SLARAGFYYTGANDK	4	ש	[E4	}	≯	E	. უ	Æ	z	Ω	×	>	出)))
	•		TTC	ŢĞ(TG	TGG	CCT	GAT	GCI	AGA	CAA	CIG	3AA.	ACA.	AGG	3GA	CAG	ğ	AT	3GA(TGCTTCTGCTGTGGCTTGGTAGACAACTGGAAACAAGGGGACAGTCCCATGGAGAAG	420
Ø	361	່ວ	E	່ບ	+ U	ָם י	-1	<u> </u>	1	Q	Z	3	×	Ø	ש	Д	တ	<u>α</u>	Σ	四	C G L M L D N W K Q G D S P M E K) 1

	FIG. 5B	ָּיָלָ מַּ	2000	ت ت ن	ς.π.Σ.	ن	של ה קבו	ָרָ בָּי	-N α	M-hiap-1 Aacmmo	-1 TGT	A C A	VAC	ևևև	A	į. Į	J.	ָּאַ מָּרְ	֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֖֓֞֟֞֓֓֓֞֟֝֓֞֓֓֞֝֓֞֝֓֞֝֓֞֝֓֓֝֓֞֝֓֡֝֝֓֡֝	5B M-hiap-1 กษณรษาจากการการการการการการการการการการการการกา	
421 H R K	1 02	1	R R	+ 17	>1	i L	က	ט	Z	E4	>	O	E	+ 13	z	ρ,	*	Z	S	LYPSCNFVQTLNPANSL	480
	Æ	ည္	TAG	TCC	TCG	S S S	ric.	ľĊŢ.	ונכני	TTC	CAC.	999	GAT(ZAG(ZAC	ZAT(, , ,	LT.	3AG	GAAGCTAGTCCTCGGCCTTCTTCCTTCCACGGCGATGAGCACCATGCCTTTGAGCTTT	2
- EB - EB - T D B	1	. 4	S	і Д	i R	i Di	S	H	Д	i S	! ! [+! }-	4	Σ	+ W	E	Σ	μ	J	S	E A S P R P S L P S T A M S T M P L S F) # n i
GC2 541		AAG	GCAAGTTCTGAGAATACTGGCTATTTCAGTGGCTCTTACTCGAG	TGA -+-	GAA	TAC	TGG(CTA	LTT 	CAG	TGG +	CTC	TTA	CHC -+-	GAG	CTTTCCCTCA	ΩΩ + t	. G	AGAC	GCAAGTTCTGAGAATACTGGCTATTTCAGTGGCTCTTACTCGAGCTTTCCCTCAGACCCT	009
₹		ט ל	S S A	ы <u>С</u>	z, ()	T	יט ק ב	א לט א	ų L	מ ב	9 E	v TTT	I GAG	SAC.	v AAG	r Tag	r CTA	າ ປິ	CTT	A S S E N I G I F S G S I S S E F S D F GTGRACTTGAGGATTGTCCTGCTTTGAGCACAAGTCCCTACCACTTTGCA	ı
601				+		1	+	1		B B B	+	1		1 + 1	† 	\ 	+	t t	1 1 1	+ + + + + + + + + + + + + + + + + + + +	099
>		z	[±4	않	Ø	Z	Ö	Ω	U	ρ,	4	니	တ	Ę-I	တ	Д	Ħ	Ħ	[tu	æ	1
	C	GAZ	CAC	'AGA	GAA	CAGAGAAGGCCAGATTACTCACCTATGAAACATGGCCATTGTCTTT	CAG	ATT	ACT	CAC	CTA	TGA	AAC	ATG	225	ATT	GIC	TTT	TCTG	ATGAACACAGAGAAGGCCAGATTACTCACCTATGAAACATGGCCATTGTCTTTTCTGTCA	720
W		H N W	H	+ E4	×	4	<u> </u>	1	H	H	>	田	E	×	Д	H	က	ſĿι		വ	1
ນີ້	ri .	CCAGCAA	AAA.	וטפע	ວອອງ	CAA	CAAAGCAGG	AGG	CII	CTA	TACTA	CAT	AGG	ACC	TGG	AGA	TAG	AGT	999	AGCTGGCCAAAGCAGGCTTCTACTACATAGGACCTGGAGATAGAGTGGCCTGC	780
Ռ	1	P A K		1	1	×	Æ	b	<u>L</u>		> 1	H	Ü	. _{Сч}	Ö	Ω	ᇟ		Ø	Ü	

1020 960 840 900 CACAGTGATGATGTCAAGTGTTTATGCTGTGGTGGGGCTGAGGTGCTGGGAATCTGGA GATGACCCCTGGGTGGAACATGCCAAGTGGTTTCCAAGGTGTGAGTACTTGCTCAGAATC AAAGGCCAAGAATTTGTCAGCCAAGTTCAAGCTGGCTATCCTCATCTACTTGAGCAGCTA CCTTCTAGTGCACTAGTTCATTCCCAGGAACTTGCAAGTGCGGGCTTTTATTATACAGGA ACTGTCTCTAACCTGAGCATGCAGACACACGCAGCCCGTATTAGAACATTCTCTAACTGG CAGAGGCATTTCCCCAGCTGTCCGTTCTTAAAAGACTTGGGTCAGTCTGCTTCGAGATAC TTTGCGTGCGATGGGAAACTGAGCAACTGGGAACGTAAGGATGATGCTATGTCAGAGCAC Ö Σ Ü a U × Ö Ü 吆 M-hiap-1 ᆸ M U 臼 Ö S Д S ·H S 耳 Σ Ы 凶 ഗ ¥ Ы Ü Ø S U FIG. 5C 1141 196 901 1021 1081 841 781 Q đ đ Q Ø Ø Ø




רו <u>ה</u> . טט	Š								M-hiap-1	3p-1											
(TT	TTATCTAC	TAC	GIC	AGA	CTC	gg.	AGA	AGA.	rgac	JAA1	ຼີ່ວ	GAC	25	(CC)	ATC	GTC.	CAT	rtti	GTCAGACTCCCCAGAAGATGAGAATGCAGACGCAGCAATCGTGCATTTTGGC	0
1201	i a	מ	<u> </u>	+ W	D	S	, С	田	D	ы	z	Æ	Ω	Æ	Æ	Н	>	ж	Įž,	SDSPEDENADAAIVHFG	1440
9	ູ່ບ	CCTGGAGA	AGA		TTC	3GGA	AGA	TGT	CGT	CATO	3AT(3AG(PAC .	υ Σ	GTC	GII	CAA	AGC2	AGCC	AAGITCGGAAGAIGTCGTCAIGAIGAGCACGCCTGTGGTTAAAGCAGCCTTG	0,00
1261	р	ָ ט	i i i	+ W	S	臼	А	D V Q	>	Σ	Σ	တ	E→	- Δι	>	>	×	Æ	A	SSEDVVMMSTPVVKAAL	0 4 1
6		AAT	999	CII	Ş	FTAG	GAG	CCT	GGT	GAG	ACA.	GAC	GT	ICA F	GTG	SCA CA	3AT(CT	ည္ဟ	GAAATGGGCTTCAGTAGGAGCCTGGTGAGACAGACGGTTCAGTGGCAGATCCTGGCCACT	. 0
1321	口口	Σ	ט	1 + Eu !	S	24	S	l l	r d	RQTVQWQIL	a	€ ⊣	>	0	3	ø	├ 	ıд	A	FSRSLVRQTVQWQILAT	0000
4 (9 9	TGA	GAA	CTA	CAC	3GAC		CAG	TGA	CCT	CGT	TAT	PGG(CHE	ACT	CGA	rg G	AGA	AGA(GGTGAGAACTACAGGACCGTCAGTGACCTCGTTATAGGCTTACTCGATGCAGAAGACGAG	7
1381	ן ט	i i i	Z	 	K	Ħ	>	S	D	TVSDLVIGLLD		ы	· છ	1	J	Д	4	凶	Ω	YRTVSDLVIGLLDAEDE	7 4 4 0 -
	•	GAG	AGP	/GG7	YGC	4GAT	rgg2	GCA	CGC	GGG	CGA	GGA	GGA	GGA	GIC	AGA'	IGA	ICT	AGC	ATGAGAGAGGAGCAGAGCAGGCGGCCGAGGAGGAGGAGTCAGATGATCTAGCACTA	7 0
1 4 4 T	Σ	N N	B	· - 回		Σ	凶	Ø	4	R	i Pi	四	ធ	· 四 ·	တ	SDDLA	ıΩ	ı	4	17))
i L	-	300:	GA	VGPA	ACA.	AAA.	ැ යෙ	1 GC1	LLL	CCA	ACA	TTT	GAC	GTG	TGT	GAC	ACC	AAT	GCT	ATCCGGAAGAACAAATGGTGCTTTTCCAACATTTGACGTGTGTGACACACAATGCTGTAT	i i
T05T	Н	E	×	Z	X		>	J	[I4	VLFQ	Ħ	ı	H	ບ	C V T	H	ļ Ω _i	E W	I. Y	} ≯	M V L F Q H L T C V T P M L Y -

	FIG. 5E	2 上	_								M-hiap-1	ap-1										
			CCI	CCI	AAG	TGC	'AAG	Sec	CAT	CAC	TGA	ACA(3GA(3TG(AA.	ເວີລີ	GIO	AA.	SE SE	BAA	TGCCTCCTAAGTGCAAGGGCCATCACTGAACAGGAGTGCAATGCTGTGAAACAGAAACCA	(
Ø	1561	ני	1	-+	+ 0		R	+ A	¦	H	Ю	ď	<u> </u>	ບ	; +	Æ	>	<u>+</u> ×	O	×	ARAITEQECNAVKOKP	1620
	,		CAC	CIT	AC.	MGC	ZAAG	SES.	ACT	GAT	TGA	TAC'	rgt	TL	AGC:	AAA	rge7	AAA(AC.	1GC	CACACCTTACAAGCAAGCACACTGATTGATACTGTGTTAGCAAAAGGAAACACTGCAGCA	000
Ø	1621	- •	<u>-</u>	H T L Q A	+ 0	4	STLIDTULAKGNT	+ !		: : H	LIDI	. [V L	ı,	Ą	×	ט	z	[⊷	4	A A	000
	•		ĊŢ	ACCTCATT	CAC	JAA.	ACTC	S S	TCG	GGA	AAT	TGA	ָ ט	IGC	GTT	AATTGACCCTGCGTTATACAGAGATAT	AG.	GA.	rati	ATT'	CAGAAACTCCCTTCGGGAAATTGACCCTGCGTTATACAGAGATATATTTGTG	7
Ø	1881	_	S		+ &	z	တ	SLRE	24	· 闰		O I	<u>р</u>	Æ	A L Y	÷	DC;	D	H	Ĺ.	F A	O .
			ACA	CAACAGGA	ICA1	TTAC	3GAG	TCI	TCCC	CAC	AGA	AGATGAC	CAT	IG	ATTGCAGCTC	ICI	TACC	VAT(3GA.	AGA	CATTAGGAGTCTTCCCACAGATGACATTGCAGCTCTACCAATGGAAGAACAG	0
đ	1/47 1		O	D D D	!	ద	တ	ı				I O O		Æ	4				田	回		000
	, ,		ည်	TTGCGGCC	ပ္ပ	ວວຼາ	7992	AGG2	CAC	AAT	GTG	TAA	AGT	GIG	TAT	GGA(CGC	AGA(GT	ATC	CCTCCCGGAGGACAGAATGTGTAAAGTGTGTATGGACCGAGAGGTATCCATC	0
đ	T087		×	L R P	, + 1	Δ,	岡	Д	DK	Σ	U	×	>	υ	Σ	Д	æ	ы	>	ι O	E V S I	
	,		ĞT	GTGTTCAT	ITC(CT	TCCCTGTGGCCATCTGGTCGTGTG	2005	LT CT	GGT	CGI	GTG	CAA	AGA	CIG	CGC	ກີດ	CTC.	ľĊŢ	GAG	TCCCTGTGGCCATCTGGTGTGCAAAGACTGCGCTCCCTCTCTGAGGAAG	6
Ø	1991		<u> </u>	V F I	D ₁	ບ	ტ	Ħ	ı	>	Δ	ט	×	A	U	A	ρı	တ	1	24	X	CAPSLRK -

5 5 -	M-hiap-1	
,	AGGGACCATCAAGGGCACAGTGCGCACATTTCTCTCTGCAACAA	0
1921	CPICRGTIKGTVRTFLS*	1 7 8 0 1 7 8 0
	CTAATGGTCCATGGCTGCAACTTCAGCCAGGAAGTTCACTGTCACTCCCAGTTCCAT	
1981		2040
.2041		2100
1	GAAAAACT) ;
2101		2160
•	TICCITAA	(
2161	TACTACCTGCATCTAAAGTATTCATATTCATATATTCAGATGTCATGAGAGAGA	2220
2221		2280
1	TGTTCTTG)
2281	+	2340
	GGCTAGAATCCATGAACCAAGCTGCAAAGATCTCACGCTAAATAAGGCGGAAAGATTTGG	
2341		2400
	AGAAACGAAAGGAAATTCTTTCCTGTCCAATGTATACTCTTCAGACTAATGACCTCTTCC	
2401	+	2460
	TATCAAGCCTTCTA	
2461	2474	

	FIG	FIG. 6A		,					-	M-hi	M-hiap-2										
٠	,	CTGTGGTGGAGATCTATTGTCCAAGTGGTGAGAACTTCATCTGGAAGTTTAAGCGGTCA	GA(3AT	CTA	TTG	TCC	AAG	TGG	TGA	GAA	ACT	ICA	TCT	3GAJ	\GT	TA	AGCC	GTC		
	rd	GAAATACTATTACTACTCATGGACAAAACTGTCTCCCAGAGACTCGCCCAAGGTACCTTA	+ AT	IAC	TAC	TCA	TGG	ACA	AAA	L L	TCT	l D	AGA	GAC	, S	i i	AAG	JTA(ATTACTACTCATGGACAAAACTGTCTCCCAGAGACTCGCCCAAGGTACCTTA	00 + €	
	61	CDCCCDADADCTTAAACGTATAATGGAGAAGAGCACAATCTTGTCAAATTGGACAAAGG	+-+	CTT		CGT	+ ATA		GAG	-+-	AGC	ACA	+ ATC	TTG	TCA.	A		A	AACTTAAACGTATAATGGAGAGAGAGCACAATCTTGTCAAATTGGACAAAAGA	+ 120	0
	121		†	+			+			+	1	1	+) 1					+	+ 180	0
Q								Σ	स्र	×	ഗ	Ę	н	'n	co 	ا ا	' 2'	EKSTILSNWTK	M M	1	
	•	GAGCGAAGAAAAATGAAGTTTGACTTTTCGTGTGAACTCTACCGAATGTCTACATATTC	3AA	AAA	ATC	:AAG	TT	GAC	TT	נדכ	TGI	GAA	CTC	TAC	CGA	ATG	ICT	ACA'	rat7		
Д	181	S E		×	Σ	×	+ E4	۵	E4	ပ	ט	Б	† - 7]	.			! !	M K F D F S C E L Y R M S T Y	K M K F D F S C E L Y R M S T Y S	+ 240	0
		AGCTTTTCCCAGGGGAGTTCCTGTCTCAGAGAGGAGTCTGGCTCGTGCTTGGCTTTTATTA	Ö,	AGG	799;	AGTJ	ည် .	GTC	CIC	AGAC	AGG	AGT	CTC	GCT	CGT	gCI	395	TTT	[AT]		(
Д	74T	AF	† ! ! (2)	æ	_U	>	- Д-	>	တ	四	24	S	J	GVPVSERSLARA	\$	1	!	1 [zr 1	+	-+ 300 Y -	>
	(GTG	AAJ	rga(CAA	\GT(:AA(3TG	CTT(TGC	TGI	999	CTG	ATG	TTG	GAT	AAC	TGG/		(
Q	301	ם ב	† ! ! >	Z	Ω	X V	>	×	ບ	<u> </u>	ບ	ບ	ט	KVKCFCCGLMLD	Σ	, ,,]		Z	CGLMLDNWK	-+ 360 K -	>
	,	ACAAGGGGACAGTCCTGTTGAAAGCACAGATTCTATCCCAGCTGCAGCTTTGTACA	GAC	.ag	ונכ בכילו	rgttgaaa	[GA.	AAA(SCA	CAG	CS	TT	TA	S	AGC	TGC	AGC	TTT	GTA((
Q	7 D F	361	٠ ١	S P	ì	>	VEKH	×	Ħ		O	[L ₁	→	RQFYPS	S	ט	S	1 [II.	PSCSFVQ	-+ 420 Q -	5

	בי סם	0								Σ	M-hiap-2	p-2										
	(_	TCT	GCI	TTC	AGC) [3]	TCT	SCA A	GTC	TCC.	ATC	FAAC	AA.	rat(3TC	rac	rgr	GAA	TITCAGCCAGICTGCAGICTCCAICIAAGAATAIGICICCTGIGAAAAGIAG	IAG	0
Д	421	7 7 H		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	S	4	တ	, 	0	တ	Д	C)	×	z	Σ	S	i Cu	<u>+</u> >	X	SASLOSPSKNMSPVKSR	+ K) (0 t [†] 1
	•		ATTTGCAC	.AC	TTC	GTC	:ACC	CCICI	GG	ACG	GGAACGAGGTGGCATTCACTCCAACCTGTGCTC	TGG(CAL	ICA	CTC	CAA	CCT	GTG	CTC	ŢĀ	ეეე	
۵	4. 1	•	1 4 F		S	တ		- LJ	阳	24	Ŋ	ט	Н	Ħ	ဟ	Z	1	ט	S	. ທ	+ 1	0 1
	1	_	TAA	ITT	TAC	3AG		1991	AAG?	CTT	CTC	ATC	A.A.G.	3AT	GGA.	ICC	CIG	CAG	CTA	TCTTAATTCTAGAGCAGTGGAAGACTTCTCATCAAGGATGGAT	CAT	0
Ω	541				2	A A	>	<u> 1</u>	Д	<u> </u>	VEDFSS	တ	æ	Σ	<u> </u>	<u>.</u>	ָט	÷ 0		A	R M D P C S Y A M	009
	,	· -	TAC	ZAG2	VAC.	AGG	ğ	AT	<u> </u>	TAC	TTA	CAG	IAT(GIG	229	TTI	AAG	TTT	TCI	GIC	GAGTACAGAAGAGGCCAGATTTCTTACTTACAGTATGTGGCCTTTAAGTTTTCTGTCACC	Q V
Д	601		ST	i H	<u> </u>	A	i ez	<u> </u> [24	i i	E	E A R F L T Y S M W P L S F	S	Σ	3	L CL	-	S	ት [፲٠ ት	<u> </u>	S	+ d S	099
	,	•	'AGA	SG	166	Š	3AG	STG.	3CTJ	CT	TTA	CAT	AGG(300	TGG	AGA	CAG	GGT	099	CTG	AGCAGAGCTGGCTGGCTTCTATTACATAGGGCCTGGAGACAGGGTGGCCTGTTT	
Ω	T 9 9		ш	E L	Æ	æ	A	ט	E4	≯	×	-	U	<u>a</u>	ָ שׁ	D	2	; 	4	ט	AELARAGFYIIGPGDRVACF	07/
	, t		CIC	3TG(JTG.	. 3G. A	A.A.C.	rga(30.50	CTO	GGA	ACC	AAA	GGA	TTA	TGC	TAI	GIC	AG?	GCP	TGCCTGTGGGAAACTGAGCAACTGGGAACCAAAGGATTATGCTATGTCAGAGCACCG	
Ω	17/		υ !	ט	ט	×	H	S	Z	3	M	D.	×	۵	, _{>}	4	Σ	S	េយ	<u> </u>	A C G G K L S N W E P K D Y A M S E H R	08/



1500 1260 1320 1380 1200 GGAAATGGGCTTCAGTAGGAGCCTGGTGAGACAGACGGTTCAGCGGCAGATCCTGGCCAC TGGTGAGAACTACAGGACCGTCAATGATATTGTCTCAGTACTTTTGAATGCTGAAGATGA GAGAAGAGAAGAGAAAGGAAAGACAGACTGAAGAGATGGCATCAGGTGACTTATCACT CACTTCAGACACCCCAGGAGAAAAATGCTGACCCTACAGAGACAGTGGTGCATTTTGG CCCTGGAGAAAGTTCGAAAGATGTCGTCATGATGAGCACGCCTGTGGTTAAAGCAGCCTT 回 ഗ K H U ທ [-d H p, O 团 H H 囟 ᆸ ഗ Д S M-hiap-2 0 团 Σ O Σ A; Z 民 ᆸ 团 P S M K Σ U 团 R S ഗ ρı, 回 Z ល Н 又 囟 O 回 吆 召 U 山 S FIG. 6D 1201 1261 1321 1381 1141 Д Д Д Д Д

SUBSTITUTE SHEET (RULE 26)

	FIG. 6E	9E									M-hiap-2	3p-2										
	, U	_	TCI	TCI	TG.	1 66(CE	3TG1	TAAT	TAC	AAA	ACA	GGA	ACA.	IGA	TAT	TAT	TAG	ACA	GAA	TAATCTTCTTGAGGCCAGTGTAATTACAAAACAGGAACATGATATTATTAGACAGAAAAC	
Ω	TOCT	מ	1		凶	4	လ	>	H	H	X	ď	团	 EE		, H	! !	+ 24	O	×	EASVITKQEHDIIRQKT	726U
	i i	-	GAI	'ACC	ĞŢ.	ľAC	AAG(CAAC	AGE	GCI	TAT	TGA	CAC	CGT	TTT	AGT	CAA	999	AAA	TGC	acagatacccttacaagcaagagcttattgacacgttttagtcaagggaaatgctgc	
Q	T9CT	o	1 H		1	0	K	PG	(A)	<u>.</u>	† !	ם	! [————————————————————————————————————		<u> </u>	>	×	+ U	Z	Æ	A R E L I D T V L V K G N A A	1620
	, ,	•	CAN	CAI	Ç.	ICA.	AAA.	ACIC	CTC	GAA	999	AAT	TGA	CTC	CAC	GTT	ATA	TGA	AAA	CTJ	AGCCAACATCTTCAAAAACTCTGAAGGGAATTGACTCCACGTTATATGAAAACTTATT	
٩	1621		Z	! ! H	Eu +	×	Z	ဟ	h	×	ט		D	S	: E4	-	>-	+ 四 + 四	Z	h	ANIFKNSLKGIDSTLYENLF	1680 -
	•		GGA	AAA	GA	YTA.	IGA	AGT?	\TA]	TCC	AAC	AGA	AGA	CGT	TIC	AGG	CTI	GIC	ATI	GG.	_ TGTGGAAAAAAATATGAAGTATATTCCAACAGAAGACGTTTCAGGCTTGTCATTGGAAGA	
Д	1681		四	×	2	Σ	×	; >1	; ; ⊢ !	ы	+ 1 E+ 1	回	0	>	, v	ט	1	+ - S		回	VEKNMKYIPTEDVSGLSLEE	1740
	ţ		GTI	ည	igac	3AT.	TAC	AAG?	AAG2	ACG	AAC	TTG	CAA	AGT	GTG	TAT	gg)	CAG.	AGP	GGJ	GCAGTTGCGGAGATTACAAGAAGAACTTGCAAAGTGTGTATGGACAGAGGTTTC	
Q	T 7 / T		17 0	R	ez t	1	Ø	凹	囮	×	H	U	×		ט	Σ	Д	+ E	田田	>	Q L R R L Q E E R T C K V C M D R E V S	0081
		_	TATTGTGT"	ĞŢ	Ž.	ľľĆ	CGT	3TG	3TC	\TC1	'AGT	AGT	CTG	ව්	GGA	ATG	TGC	ပ္ပ	TIC	Ę,	TCATTCCGTGTCATCTAGTAGTCTGCCAGGAATGTGCCCCTTCTCTAAG	
ą	1801		>	<u>Du</u>	; H	D ₁	ט	t to	二	J		>	ပ	O	<u> </u>	U	A	<u> </u>	တ	H	I V F I P C G H L V V C Q E C A P S L R	1860

2340 2280 2040 2100 2160 2220 1920 1980 **rccagtctgggaaataaggaggaatctgctggtaaaaatttgctggatgtgagaa**t **AGATGAAAGTGTTTCGGGTGGGGGGGGTGTCAGTGTAGTGTGTGCAGGGATGTATGCAG** 3AAGGGTAGCATTGTATATTTAAGCTTAGTCTGTTGCAAGGGAAGGTCTATGCTGTTGAG GAAGAATGGTCTGAAAGTATTGTTGGACATCAGAAGCTGTCAGAACAAAGAATGAACTAC CTTCTTGGGATTTGGGAAATTTGGGGAAGCTTTGGAATCCAGTGATGTGGAGCTCAGAAA **ICCTGGAACCAGTGACTCTGGTACTCAGTAGATAGGGTACCCTGTACTTCTTGGTGCTTT IGATTTCAGCTCTTCAGCAGGACATTCTACTCTTTCAAGATTAGTAATCTTGCTTTAT** GAAGTGCCCCATCTGCAGGGGGACAATCAAGGGGACTGTGCGCACATTTCTCTCATGAGT ഗ L [L E 吆 Þ ы r M-hiap-2 × **!-1** H Ç œ SCCAAACACTGTGTAG U Н Д U 노 FIG. 6F 2341 2401 1981 2041 2101 2161 2221 2281 1921 1861

Д

41/67

FIG. 7A





FIG. 7B



SUBSTITUTE SHEET (RULE 26)

42/67

FIG. 7C





FIG. 7D



SUBSTITUTE SHEET (RULE 26)

43/67

FIG. 7E




FIG. 7F 0.6 0.5 Total protein (mg/ ml) 0.4 0.3 0.2 0.1 90 Mock co. **D**5 E5 G5 Be 9 A5 CS 5 A6 **B**2 呈

SUBSTITUTE SHEET (RULE 26)

asOligo well ID

44/67

FIG. 7G





FIG. 7H



SUBSTITUTE SHEET (RULE 26)

45/67

FIG. 71





FIG. 7J



46/67

FIG. 7K






FIG. 7L



SUBSTITUTE SHEET (RULE 26)










FIG. 8C



48/67



SUBSTITUTE SHEET (RULE 26)

49/67

FIG. 9B







FIG. 9C











SUBSTITUTE SHEET (RULE 26)


52/67








SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

55/67



SUBSTITUTE SHEET (RULE 26)

56/67

FIG. 15A

1 TTGCAGGTAC TTAGAATTTT TCCTGAGCCA CCCTCTAGAG GGCAGTGTTA 51 CATATATATC TGTAATTATC CAGTTACAAC AAAAAAAGGG CTCTCATTCA 101 TGCATGAAAA TCAGAAATAT TTCATACTCT TAAAGAACAC ATTGGAACCA 151 ATATTATGAT TAAAACATAT TTTGCTAAGC AAAGAGATAT TAAAAATTAA 201 TTCATTAACA TTCTGAACAT TTTTTAACTT GTAAAAACAA CTTTGATGCC 251 TTGAATATA AATGATTCAT TATAACAATT ATGCATAGAT TTTAATAATC 301 TGCATATTTT ATGCTTTCAT GTTTTTCCTA ATTAATGATT TGACATGGTT 351 AATAATTATA ATATATTCTG CATCACAGTT TACATATTTA TGTAAAATAA 401 GCATTTAAAA ATTATTAGTT TTATTCTGCC TGCTTAAATA TTACTTTCCT 451 CAAAAAGAGA AAACAAAAAT GCTAGATTTT ACTTTATGAC TTGAATGATG 501 TGGTAATGTC GAACTCTAGT ATTTAGAATT AGAATGTTTC TTAGCGGTCG 551 TGTAGTTATT TTTATGTCAT AAGTGGATAA TTTGTTAGCT CCTATAACAA 601 AAGTCTGTTG CTTGTGTTTC ACATTTTGGA TTTCCTAATA TAATGTTCTC 651 TTTTTAGAAA AGGTGGACAA GTCCTATTTT CAAGAGAAGA TGACTTTTAA 701 CAGTTTTGAA GGATCTAAAA CTTGTGTACC TGCAGACATC AATAAGGAAG 751 AAGAATTTGT AGAAGAGTTT AATAGATTAA AAACTTTTGC TAATTTTCCA 801 AGTGGTAGTC CTGTTTCAGC ATCAACACTG GCACGAGCAG GGTTTCTTTA 851 TACTGGTGAA GGAGATACCG TGCGGTGCTT TAGTTGTCAT GCAGCTGTAG 901 ATAGATGGCA ATATGGAGAC TCAGCAGTTG GAAGACACAG GAAAGTATCC 951 CCAAATTGCA GATTTATCAA CGGCTTTTAT CTTGAAAATA GTGCCACGCA

57/67

FIG. 15B

- 1001 GTCTACAAAT TCTGGTATCC AGAATGGTCA GTACAAAGTT GAAAACTATC
 1051 TGGGAAGCAG AGATCATTTT GCCTTAGACA GGCCATCTGA GACACATGCA
 1101 GACTATCTTT TGAGAACTGG GCAGGTTGTA GATATATCAG ACACCATATA
 1151 CCCGAGGAAC CCTGCCATGT ATTGTGAAGA AGCTAGATTA AAGTCCTTTC
 1201 AGAACTGGCC AGACTATGCT CACCTAACCC CAAGAGAGTT AGCAAGTGCT
 1251 GGACTCTACT ACACAGGTAT TGGTGACCAA GTGCAGTGCT TTTGTTGTGG
 1301 TGGAAAACTG AAAAATTGGG AACCTTGTGA TCGTGCCTGG TCAGAACACA
 1351 GGCGACACTT TCCTAATTGC TTCTTTGTTT TGGGCCGGAA TCTTAATATT
 1401 CGAAGTGAAT CTGATGCTGT GAGTTCTGAT AGGAATTTCC CAAATTCAAC
 1451 AAATCTTCCA AGAAATCCAT CCATGGCAGA TTATGAAGCA CGGATCTTTA
 1501 CTTTTGGGAC ATGGATATAC TCAGTTAACA AGGAGCAGCT TGCAAGAGCT
 1551 GGATTTTATG CTTTAGGTGA AGGTGATAAAA GTAAAAGTGCT TTCACTGTGG
 1601 AGGAGGGCTA ACTGATTGGA AGCCCAGTGA AGACCCTTGG
- 1651 CTAAATGGTA TCCAGGGTGC AAATATCTGT TAGAACAGAA GGGACAAGAA
 - 1701 TATATAAACA ATATTCATTT AACTCATTCA CTTGAGGAGT GTCTGGTAAG
 - 1751 AACTACTGAG AAAACACCAT CACTAACTAG AAGAATTGAT GATACCATCT
 - 1801 TCCAAAATCC TATGGTACAA GAAGCTATAC GAATGGGGTT CAGTTTCAAG
- 1851 GACATTAAGA AAATAATGGA GGAAAAAATT CAGATATCTG GGAGCAACTA
- 1901 TAAATCACTT GAGGTTCTGG TTGCAGATCT AGTGAATGCT CAGAAAGACA
 - 1951 GTATGCAAGA TGAGTCAAGT CAGACTTCAT TACAGAAAGA GATTAGTACT
- 2001 GAAGAGCAGC TAAGGCGCCT GCAAGAGGAG AAGCTTTGCA AAATCTGTAT

58/67

FIG. 15C

2051 GGATAGAAAT ATTGCTATCG TTTTTGTTCC TTGTGGACAT CTAGTCACTT 2101 GTAAACAATG TGCTGAAGCA GTTGACAAGT GTCCCATGTG CTACACAGTC 2151 ATTACTTTCA AGCAAAAAT TTTTATGTCT TAATCTAACT CTATAGTAGG 2201 CATGTTATGT TGTTCTTATT ACCCTGATTG AATGTGTGAT GTGAACTGAC 2251 TTTAAGTAAT CAGGATTGAA TTCCATTAGC ATTTGCTACC AAGTAGGAAA 2301 AAAAATGTAC ATGGCAGTGT TTTAGTTGGC AATATAATCT TTGAATTTCT 2351 TGATTTTCA GGGTATTAGC TGTATTATCC ATTTTTTTA CTGTTATTTA 2401 ATTGAAACCA TAGACTAAGA ATAAGAAGCA TCATACTATA ACTGAACACA 2451 ATGTGTATTC ATAGTATACT GATTTAATTT CTAAGTGTAA GTGAATTAAT 2501 CATCTGGATT TTTTATTCTT TTCAGATAGG CTTAACAAAT GGAGCTTTCT 2551 GTATATAAAT GTGGAGATTA GAGTTAATCT CCCCAATCAC ATAATTTGTT 2601 TTGTGTGAAA AAGGAATAAA TTGTTCCATG CTGGTGGAAA GATAGAGATT 2651 GTTTTTAGAG GTTGGTTGTT GTGTTTTAGG ATTCTGTCCA TTTTCTTTTA 2701 AAGTTATAAA CACGTACTTG TGCGAATTAT TTTTTTAAAG TGATTTGCCA 2751 TTTTTGAAAG CGTATTTAAT GATAGAATAC TATCGAGCCA ACATGTACTG 2801 ACATGGAAAG ATGTCAAAGA TATGTTAAGT GTAAAATGCA AGTGGCÀAAA

2851 CACTATGTAT AGTCTGAGCC AGATCAAAGT ATGTATGTTT TTAATATGCA
2901 TAGAACAAAA GATTTGGAAA GATATACACC AAACTGTTAA ATGTGGTTTC
2951 TCTTCGGGGA GGGGGGGATT GGGGGGAGGGG CCCCATAGGG GTTTTATAGG

59/67

FIG. 16A

- 1 TTGCTCTGTC ACCCAGTTTG GAGTGCAGTT ATGCAGTCTC ACACTGCAAG
- 51 CTCTGCCTCA TGGGCTCAAG TGAACCTCCT GCCTCAGCCT CTCAAGTAGC
- 101 TGGGACCACA GGCAGGTGCC ACCATGTCTG GCTAATTTTT
 GAGTTTCTTT
- 151 GTAGAGATGG TGTTTTGCCA AGTCACCCAG TTTGAGGCTG GTCTCAAACA
- 201 CCTGGGCTCA AGCAATCCAT CTACCTCAGC CTCCCAAAGT GCTGGGATTA
- 251 CAGGAGTGAG CCATGGCATG AGGCCTTGTG GGGTGTCTCT
 TTTAAATGAA
- 301 AGCATACTCT GTTTACGTAT TTGATATGAA GGAATATCCT TCCTTTCCAC
- 351 AAAGACAAAA ATTATCCTAT TTTTCTCAAA ACATATGTCC
 TTTTTCTCTA
- 401 CTTTTCATTT TTGTTACTTT TGATGGACAC ATGTGTTACA
 TTGATTTCAC
- 451 TTTCTCATAA TTCTGCTGTA AGAAAAACAA TAGTGCCAGT TCAATGACAA
- 501 ATAGCAACAG TCTGTTATTG CTAGACTGTT ACTGTTAGTG GAGACTACCA
- 551 GAACAGTCAG TCCCAGTGTC AGGGAATCAA AGAGAACATG
 TTCCCTCTCT
- 601 AAAGGGCACA GCTGCTGCTC AGCTTTAGCT GATTGCTGCC CTGCAGGACT
- 651 ATAGGCCCAG TGTTGCTAGA TCTTTTGATG TTTCAAGAGA AGCTTGGAAT
- 701 CTAGAATGTG ATGGGAAGTC TCTTACATTT AAACATGTTG GCAATTAATG

60/67

FIG. 16B

- 751 GTAAGATTTA AAAATACTGT GGTCCAAGAA AAAAATGGAT TTGGAAACTG
- 801 GATTAAATTC AAATGAGGCA TGCAGATTAA TCTACAGCAT GGTACAATGT
- 851 GAATTTTCTG GTTTCTTTAA TTGCACTGTA ATTAGGTAAG ATGTTAGCTT
- 901 TGGGGAAGCT AAGTGCAGAG TATGCAGAAA CTATTATTTT
 TGTAAGTTTT
- 951 CTCTAAGTAT AAATAAATTT CAAAATAAAA ATAAAAACTT AGTAAAGAAC
- 1001 TATAATGCAA TTCTATGTAA GCCAAACATA ATATGTCTTC CAGTTTGAAA
- 1051 CCTCTGGGTT TTATTTTATT TTATTTTATT TTTGAGACAG
 AGTCTTGCTG
- 1101 TGTCACCCAG GCTGGAGTGT AGTGGCACTA TTTCGGCCCA CTGCAACCTC
- 1151 CACCTCCCAG GCTCAAATGA TTCTCCTGCC TCAGCCTCCG
 GAGTAGCTGG
- 1201 GATTACAGGC GCGTACCACC ACACCCAGCT AATTTTTGTA
- 1251 AGATGGGGTT TCACCATTTT GGCCAGGCTG GTTTTGAACT CCTGACCTCA
- 1301 AGTGATCCAC TTGTCTTGGC CTCCCAAAAT GCTGGGATTA CAGGCGTGAG
- 1351 CCACTGCACC AGGCAGAGGC CTCTGTTTTT TATCTCTTTT
 TGGCCTCTAC
- 1401 AGTGCCTAGT AAAGCACCTG ATACATGGTA AACGATCAGT AATTACTAGT
- 1451 ACTCTATTTT GGAGAAAATG ATTTTTTAAA AAGTCATTGT GTTCCATCCA

FIG. 16C 61/67

- 1501 TGAGTCGTTT GAGTTTTAAA ACTGTCTTTT TGTTTGTTTT
 TGAACAGGTT
- 1551 TACAAAGGAG GAAAACGACT TCTTCTAGAT TTTTTTTCA GTTTCTTCTA
- 1601 TAAATCAAAA CATCTCAAAA TGGAGACCTA AAATCCTTAA AGGGACTTAG
- 1651 TCTAATCTCG GGAGGTAGTT TTGTGCATGG GTAAACAAAT TAAGTATTAA
- 1701 CTGGTGTTTT ACTATCCAAA GAATGCTAAT TTTATAAACA TGATCGAGTT
- 1751 ATATAAGGTA TACCATAATG AGTTTGATTT TGAATTTGAT
 TTGTGGAAAT
- 1801 AAAGGAAAAG TGATTCTAGC TGGGGCATAT TGTTAAAGCA
- 1851 AGTTGGCCAG GCAGTCTCCT ACTGGCACAT TCTCCCATTA
 TGTAGAATAG
- 1901 AAATAGTACC TGTGTTTGGG AAAGATTTTA AAATGAGTGA CAGTTATTTG
- 1951 GAACAAAGAG CTAATAATCA ATCCACTGCA AATTAAAGAA ACATGCAGAT
- 2001 GAAAGTTTTG ACACATTAAA ATACTTCTAC AGTGACAAAG AAAAATCAAG
- 2051 AACAAAGCTT TTTGATATGT GCAACAAATT TAGAGGAAGT AAAAAGATAA
- 2101 ATGTGATGAT TGGTCAAGAA ATTATCCAGT TATTTACAAG GCCACTGATA
- 2151 TTTTAAACGT CCAAAAGTTT GTTTAAATGG GCTGTTACCG CTGAGAATGA
- 2201 TGAGGATGAG AATGATGGTT GAAGGTTACA TTTTAGGAAA TGAAGAAACT
 - 2251 TAGAAAATTA ATATAAAGAC AGTGATGAAT ACAAAGAAGA

62/67

FIG. 16D

TTTTTATAAC

2301 AATGTGTAAA ATTTTTGGCC AGGGAAAGGA ATATTGAAGT TAGATACAAT

2351 TACTTACCTT TGAGGGAAAT AATTGTTGGT AATGAGATGT GATGTTTCTC

2401 CTGCCACCTG GAAACAAAGC ATTGAAGTCT GCAGTTGAAA AGCCCAACGT

2451 CTGTGAGATC CAGGAAACCA TGCTTGCAAA CCACTGGTAA

2501 AAAAAAAAA AAAGCCACAG TGACTTGCTT ATTGGTCATT
GCTAGTATTA

2551 TCGACTCAGA ACCTCTTTAC TAATGGCTAG TAAATCATAA TTGAGAAATT

2601 CTGAATTTTG ACAAGGTCTC TGCTGTTGAA ATGGTAAATT

2651 TTTGTCATGA TAAATTCTGG TTCAAGGTAT GCTATCCATG AAATAATTTC

2701 TGACCAAAAC TAAATTGATG CAATTTGATT ATCCATCTTA
GCCTACAGAT

2751 GGCATCTGGT AACTTTTGAC TGTTTTAAAA AATAAATCCA CTATCAGAGT

2801 AGATTTGATG TTGGCTTCAG AAACATTTAG AAAAACAAAA GTTCAAAAAT

2851 GTTTTCAGGA GGTGATAAGT TGAATAACTC TACAATGTTA GTTCTTTGAG

2901 GGGGACAAAA AATTTAAAAT CTTTGAAAGG TCTTATTTTA CAGCCATATC

2951 TAAATTATCT TAAGAAAATT TTTAACAAAG GGAATGAAAT ATATATCATG

3001 ATTCTGTTTT TCCAAAAGTA ACCTGAATAT AGCAATGAAG
TTCAGTTTTG

63/67

FIG. 16E

- 3051 TTATTGGTAG TTTGGGCAGA GTCTCTTTTT GCAGCACCTG TTGTCTACCA
- 3101 TAATTACAGA GGACATTTCC ATGTTCTAGC CAAGTATACT ATTAGAATAA
- 3151 AAAAACTTAA CATTGAGTTG CTTCAACAGC ATGAAACTGA GTCCAAAAGA
- 3201 CCAAATGAAC AAACACATTA ATCTCTGATT ATTTATTTTA AATAGAATAT
- 3251 TTAATTGTGT AAGATCTAAT AGTATCATTA TACTTAAGCA ATCATATTCC
- 3301 TGATGATCTA TGGGAAATAA CTATTATTTA ATTAATATTG AAACCAGGTT
- 3351 TTAAGATGTG TTAGCCAGTC CTGTTACTAG TAAATCTCTT TATTTGGAGA
- 3401 GAAATTTTAG ATTGTTTTGT TCTCCTTATT AGAAGGATTG TAGAAAGAAA
- 3451 AAAATGACTA ATTGGAGAAA AATTGGGGAT ATATCATATT TCACTGAATT
- 3501 CAAAATGTCT TCAGTTGTAA ATCTTACCAT TATTTTACGT ACCTCTAAGA
- 3551 AATAAAAGTG CTTCTAATTA AAATATGATG TCATTAATTA TGAAATACTT
- 3601 CTTGATAACA GAAGTTTTAA AATAGCCATC TTAGAATCAG TGAAATATGG
- 3651 TAATGTATTA TTTTCCTCCT TTGAGTNAGG TCTTGTGCTT
 TTTNTTCCTG
- 3701 GCCACTAAAT NTCACCATNT CCAANAAGCA AANTAAACCT ATTCTGAATA
- 3751 TTTTTGCTGT GAAACACTTG NCAGCAGAGC TTTCCCNCCA TGNNAGAAGC

64/67

FIG. 16F

- 3801 TTCATGAGTC ACACATTACA TCTTTGGGTT GATTGAATGC CACTGAAACA
- 3851 TTTCTAGTAG CCTGGAGNAG TTGACCTACC TGTGGAGATG CCTGCCATTA
- 3901 AATGGCATCC TGATGGCTTA ATACACATCA CTCTTCTGTG NAGGGTTTTA
- 3951 ATTTTCAACA CAGCTTACTC TGTAGCATCA TGTTTACATT GTATGTATAA
- 4001 AGATTATACN AAGGTGCAAT TGTGTATTTC TTCCTTAAAA TGTATCAGTA
- 4051 TAGGATTTAG AATCTCCATG TTGAAACTCT AAATGCATAG AAATAAAAAT
- 4101 AATAAAAAT TTTTCATTTT GGCTTTTCAG CCTAGTATTA AAACTGATAA
- 4151 AAGCAAAGCC ATGCACAAAA CTACCTCCCT AGAGAAAGGC TAGTCCCTTT
- 4201 TCTTCCCCAT TCATTTCATT ATGAACATAG TAGAAAACAG CATATTCTTA
- 4251 TCAAATTTGA TGAAAAGCGC CAACACGTTT GAACTGAAAT ACGACTTGTC
- 4301 ATGTGAACTG TACCGAATGT CTACGTATTC CACTTTTCCT GCTGGGGTTC
- 4351 CTGTCTCAGA AAGGAGTCTT GCTCGTGCTG GTTTCTATTA CACTGGTGTG
- 4401 AATGACAAGG TCAAATGCTT CTGTTGTGGC CTGATGCTGG ATAACTGGAA
- 4451 AAGAGGAGAC AGTCCTACTG AAAAGCATAA AAAGTTGTAT CCTAGCTGCA
- 4501 GATTCGTTCA GAGTCTAAAT TCCGTTAACA ACTTGGAAGC TACCTCTCAG
 - 4551 CCTACTTTTC CTTCTTCAGT AACACATTCC ACACACTCAT

65/67

FIG. 16G

TACTTCCGGG

4601 TACAGAAAAC AGTGGATATT TCCGTGGCTC TTATTCAAAC TCTCCATCAA

4651 ATCCTGTAAA CTCCAGAGCA AATCAAGAAT TTTCTGCCTT GATGAGAAGT

4701 TCCTACCCCT GTCCAATGAA TAACGAAAAT GCCAGATTAC
TTACTTTTCA

4751 GACATGGCCA TTGACTTTTC TGTCGCCAAC AGATCTGGCA CGAGCAGGCT

4801 TTTACTACAT AGGACCTGGA GACAGAGTGG CTTGCTTTGC CTGTGGTGGA

4851 AAATTGAGCA ATTGGGAACC GAAGGATAAT GCTATGTCAG
AACACCTGAG

4901 ACATTTTCCC AAATGCCCAT TTATAGAAAA TCAGCTTCAA GACACTTCAA

4951 GATACACAGT TTCTAATCTG AGCATGCAGA CACATGCAGC CCGCTTTAAA

5001 ACATTCTTTA ACTGGCCCTC TAGTGTTCTA GTTAATCCTG AGCAGCTTGC

5051 AAGTGCGGGT TTTTATTATG TGGGTAACAG TGATGATGTC AAATGCTTTT

5101 GCTGTGATGG TGGACTCAGG TGTTGGGAAT CTGGAGATGA
TCCATGGGTT

5151 CAACATGCCA AGTGGTTTCC AAGGTGTGAG TACTTGATAA GAATTAAAGG

5201 ACAGGAGTTC ATCCGTCAAG TTCAAGCCAG TTACCCTCAT CTACTTGAAC

5251 AGCTGCTATC CACATCAGAC AGCCCAGGAG ATGAAAATGC AGAGTCATCA

5301 ATTATCCATT TTGAACCTGG AGAAGACCAT TCAGAAGATG CAATCATGAT

66/67

FIG. 16H

- 5351 GAATACTCCT GTGATTAATG CTGCCGTGGA AATGGGCTTT AGTAGAAGCC
- 5401 TGGTAAAACA GACAGTTCAG AGAAAAATCC TAGCAACTGG AGAGAATTAT
- 5451 AGACTAGTCA ATGATCTTGT GTTAGACTTA CTCAATGCAG AAGATGAAAT
- 5501 AAGGGAAGAG GAGAGAAAA GAGCAACTGA GGAAAAAGAA TCAAATGATT
- 5551 TATTATTAAT CCGGAAGAAT AGAATGGCAC TTTTTCAACA TTTGACTTGT
- 5601 GTAATTCCAA TCCTGGATAG TCTACTAACT GCCGGAATTA TTAATGAACA
- 5651 AGAACATGAT GTTATTAAAC AGAAGACACA GACGTCTTTA CAAGCAAGAG
- 5701 AACTGATTGA TACGATTTTA GTAAAAGGAA ATATTGCAGC CACTGTATTC
- 5751 AGAAACTCTC TGCAAGAAGC TGAAGCTGTG TTATATGAGC ATTTATTTGT
- 5801 GCAACAGGAC ATAAAATATA TTCCCACAGA AGATGTTTCA GATCTACCAG
- 5851 TGGAAGAACA ATTGCGGAGA CTACAAGAAG AAAGAACATG TAAAGTGTGT
- 5901 ATGGACAAAG AAGTGTCCAT AGTGTTTATT CCTTGTGGTC ATCTAGTAGT
- 5951 ATGCAAAGAT TGTGCTCCTT CTTTAAGAAA GTGTCCTATT TGTAGGAGTA
- 6001 CAATCAAGGG TACAGTTCGT ACATTTCTTT CATGAAGAAG AACCAAAACA
- 6051 TCGTCTAAAC TTTAGAATTA ATTTATTAAA TGTATTATAA CTTTAACTTT

67/67

FIG. 161

- 6101 TATCCTAATT TGGTTTCCTT AAAATTTTTA TTTATTTACA
 ACTCAAAAAA
- 6151 CATTGTTTTG TGTAACATAT TTATATATGT ATCTAAACCA TATGAACATA
- 6201 TATTTTTAG AAACTAAGAG AATGATAGGC TTTTGTTCTT
 ATGAACGAAA
- 6251 AAGAGGTAGC ACTACAAACA CAATATTCAA TCAAAATTTC AGCATTATTG
- 6301 AAATTGTAAG TGAAGTAAAA CTTAAGATAT TTGAGTTAAC CTTTAAGAAT
- 6351 TTTAAATATT TTGGCATTGT ACTAATACCG GGAACATGAA GCCAGGTGTG
- 6401 GTGGTATGTG CCTGTAGTCC CAGGCTGAGG CAAGAGAATT ACTTGAGCCC
- 6451 AGGAGTTTGA ATCCATCCTG GGCAGCATAC TGAGACCCTG
- 6501 CAAACAGAAC AAAAACAAAA CACCAGGGAC ACATTTCTCT
 GTCTTTTTTG
- 6551 ATCAGTGTCC TATACATCGA AGGTGTGCAT ATATGTTGAA TCACATTTTA
- 6601 GGGACATGGT GTTTTTATAA AGAATTCTGT GAGAAAAAAT TTAATAAAGC
 - 6651 AACCAAAAA AAAAAAAA

SEQUENCE LISTING

<110>	University of Ottawa Aegera Therapeutics, Inc.	
	Antisense IAP Nucleic Acids and Uses reof	
<130>	07891/025WO1	
	US 09/672,717 2000-09-28	
<160>	231	
<170>	FastSEQ for Windows Version 4.0	
<210><211><212><213>	19	
<220> <223>	based on Homo sapiens	
<400> aaaatt		.9
<210><211><211><212><213>	19	
<220> <223>	based on Homo sapiens	
<400> tctaga		.9
<210> <211> <212> <213>	19	
<220> <223>	based on Homo sapiens	
<400> cagata		و.
<210> <211> <212> <213>	19	
<220>	based on Home sanions	

<400> 4 tgagagccct ttttttgtt	19
<210> 5 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 5 agtatgaaat atttctgat	19
<210> 6 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 6 attggttcca atgtgttct	19
<210> 7 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 7 ttagcaaaat atgttttaa	19
<210> 8 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 8 tgaattaatt tttaatatc	19
<210> 9 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 9 attcaaggca tcaaagttg	19

PCT/CA01/01379

WO 02/026968

<210> 10 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens .	
<400> 10 gtcaaatcat taattagga	19
<210> 11 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 11 aatatgtaaa ctgtgatgc	19
<210> 12 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 12 gcagaataaa actaataat	19
<210> 13 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 13 gaaagtaata tttaagcag	19
<210> 14 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 14 ttaccacatc attcaagtc	19
<210> 15 <211> 19 <212> DNA	

<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 15 ctaaatacta gagttcgac	19
<210> 16 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 16 acacgaccgc taagaaaca	19
<210> 17 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 17 tatccactta tgacataaa	19
<210> 18 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 18 gttataggag ctaacaaat	19
<210> 19 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 19 aatgtgaaac acaagcaac	19
<210> 20 <211> 19 <212> DNA <213> Artificial Sequence	
<220>	

<223> based on Homo sapiens	
<400> 20 acattatatt aggaaatcc	19
<210> 21 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 21 cttgtccacc ttttctaaa	19
<210> 22 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 22 atcttctctt gaaaatagg	19
<210> 23 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	•
<400> 23 ccttcaaaac tgttaaaag	19
<210> 24 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 24 atgtctgcag gtacacaag	19
<210> 25 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 25	

atctattaaa ctcttctac	19
<210> 26 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 26 acaggactac cacttggaa	19
<210> 27 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 27 tgccagtgtt gatgctgaa	19
<210> 28 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 28 gtataaagaa accctgctc	19
<210> 29 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 29 cgcacggtat ctccttcac	19
<210> 30 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 30 ctacagctgc atgacaact	19
<210> 31	

<211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 31 gctgagtete catattgce	19
<210> 32 <211> 19 <212> DNA <213> Artificial Sequence	•
<220> <223> based on Homo sapiens	
<400> 32 atactttcct gtgtcttcc	19
<210> 33 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 33 gataaatctg caatttggg	19
<210> 34 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 34 ttgtagactg cgtggcact	19
<210> 35 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 35 accattctgg ataccagaa	19
<210> 36 <211> 19 <212> DNA <213> Artificial Sequence	

<220> <223> based on Homo sapiens	
<400> 36 agttttcaac tttgtactg	19
<210> 37 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 37 . atgatetetg etteceaga	19
<210> 38 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 38 agatggcctg tctaaggca	19
<210> 39 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 39 · agttctcaaa agatagtct	19
<210> 40 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 40 gtgtctgata tatctacaa	19
<210> 41 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	

<400> 41 tcgggtatat ggtgtctga	19
<210> 42 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 42 cagggttcct cgggtatat	19
<210> 43 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 43 gcttcttcac aatacatgg	19
<210> 44 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 44 ggccagttct gaaaggact	19
<210> 45 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 45 gctaactctc ttggggtta	19
<210> 46 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 46 gtgtagtaga gtccagcac	1 9

<210> 47 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 47 aagcactgca cttggtcac	19
<210> 48 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 48 ttcagttttc caccacaac	19
<210> 49 <211> 19 <212> DNA <213> Artificial Sequence	
<220> . <223> based on Homo sapiens	
<400> 49 acgatcacaa ggttcccaa	19
<210> 50 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 50 tcgcctgtgt tctgaccag	19
<210> 51 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 51 ccggcccaaa acaaagaag	19
<210> 52 <211> 19 <212> DNA	

<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 52 . gattcacttc gaatattaa	19
<210> 53 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 53 tatcagaact cacagcatc	19
<210> 54 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 54 ggaagatttg ttgaatttg	19
<210> 55 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 55 tetgecatgg atggattte	19
<210> 56 <211> 19 <212> DNA <213> Artificial Sequence	
<pre><<220> <223> based on Homo sapiens</pre>	
<400> 56 aagtaaagat ccgtgcttc	19
<210> 57 <211> 19 <212> DNA <213> Artificial Sequence	
<220>	

<223> based on Homo sapiens	
<400> 57 ctgagtatat ccatgtccc	19
<210> 58 <211> 19	
<212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 58	
gcaagctgct ccttgttaa	19
<210> 59	
<211> 19 <212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 59	
aaagcataaa atccagctc	19
<210> 60	
<211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 60 gaaagcactt tactttatc	19
<210> 61 <211> 19	
<212> DNA	
<213> Artificial Sequence	•
<220>	
<223> based on Homo sapiens	
<400> 61	
actgggcttc caatcagtt	19
<210> 62	
<211> 19 <212> DNA	
<212> DNA <213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 62	

gttgttccca agggtcttc	. 19
<210> 63 <211> 19 <212> DNA <213> Artificial Sequence	,
<220> <223> based on Homo sapiens	
<400> 63 accetggata ccatttage	19
<210> 64 <211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 64 tgttctaaca gatatttgc	19
<210> 65 <211> 19 <212> DNA	
<213> Artificial Sequence <220>	
<223> based on Homo sapiens <400> 65	
tatatattct tgtcccttc <210> 66	19
<211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 66 agttaaatga atattgttt	19
<210> 67 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 67 gacactcctc aagtgaatg	19
<210> 68	

<211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 68	10
<210> 69	19
<211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 69 gttagtgatg gtgttttct	19
<210> 70	
<211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 70 agatggtatc atcaattct	19
<210> 71	
<211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 71	19
tgtaccatag gattttgga	19
<210> 72 <211> 19	
<212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 72 ccccattcgt atagcttct	19
<210> 73 <211> 19	
<212> DNA	
<213> Artificial Sequence	

<220> <223> based on Homo sapiens	
<400> 73 attattttct taatgtcct	19
<210> 74 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 74 caagtgattt atagttgct	19
<210> 75 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 75 tagatetgea accagaace	19
<210> 76 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	·
<400> 76 catcttgcat actgtcttt	19
<210> 77 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 77 ccttagctgc tcttcagta	19
<210> 78 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	

<400> 78 aagcttctcc tcttgcagg	19
<210> 79 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 79 atatttctat ccatacaga	19
<210> 80 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 80 ctagatgtcc acaaggaac	19
<210> 81 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 81 agcacattgt ttacaagtg	19
<210> 82 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 82 agcacatggg acacttgtc	19
<210> 83 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 83 cttgaaagta atgactgtg	19

<210> 84 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 84 cctactatag agttagatt	19
<210> 85 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 85 attcaatcag ggtaataag	19
<210> 86 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 86 aagtcagttc acatcacac	19
<210> 87 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 87 cagtaaaaaa aatggataa	19
<210> 88 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 88 ttcagttata gtatgatgc	19
<210> 89 <211> 19 <212> DNA	

<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 89 tacacttaga aattaaatc	19
<210> 90 <211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 90 tetetatett tecaceage	19
<210> 91 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 91 agaatcctaa aacacaaca	19
<210> 92 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 92 attcgcacaa gtacgtgtt	19
<210> 93 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 93 tgtcagtaca tgttggctc	19
<210> 94 <211> 19 <212> DNA <213> Artificial Sequence	
<220>	

<223> based on Homo sapiens	
<400> 94 acatagtgtt ttgccactt	19
<210> 95 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 95 ctttgatctg gctcagact	19
<210> 96 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 96 gaaaccacat ttaacagtt	19
<210> 97 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 97 tcatttgagc ctgggaggu	19
<210> 98 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 98 cggaggctga ggcaggaga	19
<210> 99 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<100 \ 99	

ggtgtggtgg tacgcgcct	19
<210> 100 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 100 acccatgcac aaaactacc	19
<210> 101 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 101 agaatgtgcc agtaggaga	19
<210> 102 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 102 tctcacagac gttgggctt	19
<210> 103 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 103 ccagtggttt gcaagcatg	19
<210> 104 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 104 gaaatttagt ggccaggaa	19
<210> 105	

-011- 10	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 105	
agaaatacac aattgcacc	
242 426	
<210> 106	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
1000-	
<220>	
<223> based on Homo sapiens	
-400× 106	
<400> 106	
tactgataca ttttaagga	
<210 107	
<210> 107 <211> 19	
<211> 19 <212> DNA	
<213> Artificial Sequence	
-210 VICILICIAL Deduction	
<220>	
<223> based on Homo sapiens	
-220- Daboa oil nome bapaons	
<400> 107	
ttcaacatgg agattctaa	
<210> 108	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 108	
atttctatgc atttagagt	
.010: 100	
<210> 109	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220	
<220>	
<223> based on Homo sapiens	
<400> 109	
aatactaggc tgaaaagcc	
<210> 110	
<211> 19	
recommendation of the second o	
<212> DNA	
<212> DNA <213> Artificial Sequence	

<220> <223> based on Homo sapiens	
<400> 110 ggctttgctt ttatcagtt	19
<210> 111 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 111 tctagggagg tagttttgt	19
<210> 112 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 112 gggaagaaaa gggactagc	19
<210> 113 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 113 gttcataatg aaatgaatg	19
<210> 114 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 114 ataagaatat gctgttttc	19
<210> 115 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	

<400> 115 ttcaaacgtg ttggcgctt	19
<210> 116 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 116 atgacaagtc gtatttcag	19
<210> 117 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 117 aagtggaata cgtagacat	19
<210> 118 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 118 agacaggaac cccagcagg	19
<210> 119 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	·
<400> 119 cgagcaagac tcctttctg	19
<210> 120 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 120 agtgtaatag aaaccagca	19

<210> 121 <211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 121 tgacettgtc attcacace	19
<210> 122 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 122 ttatccagca tcaggccac	19
<210> 123 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 123 actgtctcct cttttccag	19
<210> 124 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 124 ttttatgctt ttcagtagg	19
<210> 125 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 125 acgaatctgc agctaggat	19
<210> 126 <211> 19 <212> DNA	

<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 126 caagttgtta acggaattt	19
<210> 127 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 127 taggctgaga ggtagcttc	19
<210> 128 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 128 gttactgaag aaggaaaag	19
<210> 129 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 129 gaatgagtgt gtggaatgt	19
<210> 130 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 130 tgttttctgt acccggaag	19
<210> 131 <211> 19 <212> DNA <213> Artificial Sequence	
<220>	

<223> based on Homo sapiens	
<400> 131 gagccacgga aatatccac	19
<210> 132 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 132 tgatggagag tttgaataa	19
<210> 133 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 133 gatttgctct ggagtttac	19
<210> 134 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 134 ggcagaaaat tcttgattt	19
<210> 135 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 135 ggacaggggt aggaacttc	19
<210> 136 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 136	

gcattttcgt tattcattg	19
<210> 137	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 137	
ctgaaaagta agtaatctg	19
<210> 138	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 138	
ggcgacagaa aagtcaatg	19
<210> 139	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 139	
ccactctgtc tccaggtcc	19
<210> 140	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 140	
ccaccacagg caaagcaag	19
,	
<210> 141	
<211> 19 <212> DNA	
<212> DNA <213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 141	19
tteggttecc aattgetea	19
<210> 142	

<211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 142 ttctgacata gcattatcc	19
<210> 143 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 143 tgggaaaatg tctcaggtg	19
<210> 144 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 144 tataaatggg catttggga	19
<210> 145 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 145 tgtcttgaag ctgattttc	19
<210> 146 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 146 gaaactgtgt atcttgaag	19
<210> 147 <211> 19 <212> DNA <213> Artificial Sequence	

<220> <223> based on Homo sapiens	
<400> 147 tgtctgcatg ctcagatta	19
<210> 148 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 148 gaatgtttta aagcgggct	19
<210> 149 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 149 cactagaggg ccagttaaa	19
<210> 150 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 150 ccgcacttgc aagctgctc	19
<210> 151 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 151 catcatcact gttacccac	19
<210> 152 <211> 19 <212> DNA <213> Artificial Sequence	
<220>	

<400> 152	
ccaccatcac agcaaaagc	19
ceaceaceae ageaaaage	
(210) 153	
<210> 153	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 153	
tccagattcc caacacctg	19
-	
<210> 154	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
1213/ Artificial Sequence	
<220>	
<223> based on Homo sapiens	
400 454	
<400> 154	
cccatggatc atctccaga	19
<210> 155	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 155	
aaccacttgg catgttgaa	19
adecaecegg caegeegad	
<210> 156	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 156	
caagtactca caccttgga	19
<210> 157	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
-	
<220>	
<223> based on Homo sapiens	
<400> 157	
	19
cctgtccttt aattcttat	19

PCT/CA01/01379

WO 02/026968

<210> 158 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 158 tgaacttgac ggatgaact	19
<210> 159 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 159 tagatgaggg taactggct	19
<210> 160 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 160 tggatagcag ctgttcaag	19
<210> 161 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 161 cattttcatc tcctgggct	19
<210> 162 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 162 tggataattg atgactctg	19
<210> 163	

<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 163 gtcttctcca ggttcaaaa	19
<210> 164 <211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 164 tattcatcat gattgcatc	19
<210> 165 <211> 19	
<212> DNA <213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 165 catttccacg gcagcatta	19
<210> 166 <211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 166 ccaggcttct actaaagcc	19
<210> 167 <211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 167 gctaggattt ttctctgaa	19
<210> 168	
<211> 19 <212> DNA -213> Artificial Segments	
<213> Artificial Sequence	
<220>	

<223> based on Homo sapiens	
<400> 168 tctataattc tctccagtt	19
<210> 169 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 169 acacaagatc attgactag	19
<210> 170 <211> 19 <212> DNA <213> Artificial Sequence .	
<220> <223> based on Homo sapiens	
<400> 170 tctgcattga gtaagtcta	19
<210> 171 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 171 ctcttccctt atttcatct	19
<210> 172 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 172 tcctcagttg ctctttctc	19
<210> 173 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
-400- 173	

gccattctat tcttccgga	19
<210> 174 <211> 19 <212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 174 agtcaaatgt tgaaaaagt	19
<210> 175 <211> 19	
<212> DNA <213> Artificial Sequence	
<220> . <223> based on Homo sapiens	
<400> 175 ccaggattgg aattacaca	19
<210> 176	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 176 attccggcag ttagtagac	19
<210> 177	
<211> 19	
<212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 177	•
taacatcatg ttcttgttc	19
<210> 178	
<211> 19 <212> DNA	
<212> DNA <213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 178	
gtctgtgtct tctgtttaa	19
<210> 179	

<211> 19	
<212> DNA	
<213> Artificial Sequence	
.000	
<220>	
<223> based on Homo sapiens	
<400> 179	
ttctcttgct tgtaaagac	19
coccetyce egeadagae	
<210> 180	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
•	
<220>	
<223> based on Homo sapiens	
<400> 180	10
ctaaaatcgt atcaatcag	19
.010. 101	
<210> 181	
<211> 19 <212> DNA	
<212> DNA <213> Artificial Sequence	
\Zi3> Altificial Sequence	
<220>	
<223> based on Homo sapiens	
<400> 181	
ggctgcaata tttcctttt	19
<210> 182	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> based on Homo sapiens	
122) Daboa on nome baptons	
<400> 182	
gagagtttct gaatacagt	19
<210> 183	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
000	
<220>	
<223> based on Homo sapiens	
<400> 183	
acagetteag ettettgea	19
<210> 184	
<211> 19	
<212> DNA	
<213> Artificial Sequence	

<220> <223> based on Homo sapiens	
<400> 184 aaataaatgc tcatataac	19
<210> 185 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 185 gaaacatctt ctgtgggaa	19
<210> 186 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 186 gttcttccac tggtagatc	19
<210> 187 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 187 cttcttgtag tctccgcaa	19
<210> 188 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 188 ttgtccatac acactttac	19
<210> 189 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	

<400> 189 aaccaaatta ggataaaag	19
<210> 190 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 190 atgttcatat ggtttagat	19
<210> 191 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 191 taagttttac ttcacttac	19
<210> 192 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 192 atgttcccgg tattagtac	19
<210> 193 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 193 gggctcaagt aattotott	19
<210> 194 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 194 gcccaggatg gattcaaac	19

<210><211><212><213>	19	
<220> <223>	based on Homo sapiens	
<221> <222> <223>		
<221> <222> <223>		
<400> yagaag		19
<210> <211> <212> <213>	18	
<220> <223>	based on Homo sapiens	
<222>	misc_feature 1,17,18 y=u or t	
<400> ygtgct		18
<210><211><212><213>	20	
<220> <223>	based on Homo sapiens	
<400> tctgc		20
<210><211><211><212><213>	18	
<220> <223>	based on Homo sapiens	
<400> gaaag	198 gaaag cgcaaccg	18
<210> <211>		

<212> DNA <213> Artific	rial Sequence	
<220> <223> based o	on Homo sapiens	
<400> 199 agccagatga cg	yaccccata gaggaacata	30
<210> 200 <211> 21 <212> DNA		
<213> Artific	ial Sequence	
<220> <223> based o	on Homo sapiens	
<400> 200 tggagatgat cc	catgggttc a	21
<210> 201 <211> 29 <212> DNA <213> Artific	cial Sequence	
<220> <223> based o	on Homo sapiens	
<400> 201 gaactcctgt co	ctttaattc ttatcaagt	29
<210> 202 <211> 27 <212> DNA <213> Artific	cial Sequence	
<220> <223> based o	on Homo sapiens	
<400> 202 ctcacacctt gg	gaaaccact tggcatg	27
<210> 203 <211> 27		
<212> DNA <213> Artific	cial Sequence	
<220> <223> based o	on Homo sapiens	
<400> 203 ggtgataaag ta	aaagtgctt tcactgt	27
<210> 204 <211> 28 <212> DNA	cial Semence	

<220> <223> based on Homo sapiens	
<400> 204 tcagtagttc ttaccagaca ctcctcaa	28
<210> 205 <211> 34 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 205 caacatgcta aatggtatcc agggtgcaaa tatc	34
<210> 206 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 206 gaaggtgaag gtcggagtc	19
<210> 207 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 207 gaagatggtg atgggattc	19
<210> 208 <211> 20 <212> DNA <213> Artificial Sequence	
<220> <223> based on Homo sapiens	
<400> 208 caagetteee gtteteagee	20
<210> 209 <211> 19 <212> DNA <213> Artificial Sequence	
<220> <221> modified_base <222> 1.17	

```
<223> y= cm
 <221> modified_base
 <222> 3,18
 <223> y=gm
 <221> modified_base
 <222> 19
 <223> y=um
 <223> based on Homo sapiens
 <400> 209
 yayagatttc atttaayyy
                                                                     19
 <210> 210
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> modified_base
 <222> 1,18
 <223> y=cm
<221> modified_base
 <222> 2,17
 <223> y=um
<223> based on Homo sapiens
<400> 210
yyacgctcgc catcgtyya
                                                                    19
<210> 211
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
<221> modified_base
<222> 3,18
<223> y=cm
<221> modified_base
<222> 1,17
<223> y=um
<221> modified_base
<222> 2,16
<223> y=gm
<223> based on Homo sapiens
<400> 211
yycccaagaa tactagyya
                                                                   19
<210> 212
```

<211> <212> <213>		
	modified_base 1,17,18 y=um	
<221> <222> <223>		
<223>	based on Homo sapiens	
<400> yaagci		19
<210><211><211><212><213>	19	
<220> <223>	based on Homo sapiens	
<400> aaggg		19
<210> <211> <212> <213>	19	
<220> <223>	based on Homo sapiens	
<400> agagg	214 acgga gtcggaggc	19
<210><211><212><213>	19	
<220> <223>	based on Homo sapiens	
<400> cggag	215 cgtga ggatggaga	19
<210><211><212><213>	68	
<220> <223>	based on Homo sapiens	

```
<221> VARIANT
 <222> 1-3,6,9,10,14,15,18-20,24, 30,32,33,35,37,40, 42-47, 49-51,
 53-57, 59-62, 64,66
 <223> Xaa=any amino acid
 <221> VARIANT
 <222> 13, 16,17
 <223> Xaa=any amino acid or is absent
 <400> 216
 Xaa Xaa Xaa Arg Leu Xaa Thr Phe Xaa Xaa Trp Pro Xaa Xaa Xaa Xaa
 1
                                 10
 Xaa Xaa Xaa Xaa Leu Ala Xaa Ala Gly Phe Tyr Tyr Xaa Gly Xaa
           20
                              25
                                                30
 Xaa Asp Xaa Val Xaa Cys Phe Xaa Cys Xaa Xaa Xaa Xaa Xaa Trp
                         40
                                           45
 Xaa Xaa Xaa Asp Xaa Xaa Xaa Xaa His Xaa Xaa Xaa Xaa Pro Xaa
  50
                     55
                                       60
 Cys Xaa Phe Val
 65
<210> 217
 <211> 46
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> based on Homo sapiens
 <221> VARIANT
 <222> 2-7,9-11,17-21,23,25, 30-32,34-35, 38-42 ,45
 <223> Xaa=any amino acid
 <221> VARIANT
 <222> 8
 <223> Xaa=Glu or Asp
 <221> VARIANT
 <222> 14,22
 <223> Xaa=Val or Ile
 <400> 217
 1
               5
                                 10
                                                    15
 Xaa Xaa Xaa Xaa Xaa Xaa Phe Xaa Pro Cys Gly His Xaa Xaa Xaa
            20
                              25
 Cys Xaa Xaa Cys Ala Xaa Xaa Xaa Xaa Cys Pro Xaa Cys
             . 40
 <210> 218
 <211> 2540
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
```

```
<222> (1)...(2540)
<223> n=a,t,c, or g
<400> 218
gaaaaggtgg acaagtccta ttttcaagag aagatgactt ttaacagttt tgaaggatct 60
aaaacttgtg tacctgcaga catcaataag gaagaagaat ttgtagaaga gtttaataga 120
ttaaaaactt ttgctaattt tccaagtggt agtcctgttt cagcatcaac actggcacga 180
gcagggtttc tttatactgg tgaaggagat accgtgcggt gctttagttq tcatqcaqct 240
gtagatagat ggcaatatgg agactcagca gttggaagac acaggaaagt atccccaaat 300
tgcagattta tcaacggctt ttatcttgaa aatagtgcca cgcagtctac aaattctggt 360
atccagaatg gtcagtacaa agttgaaaac tatctgggaa gcagagatca ttttgcctta 420
gacaggccat ctgagacaca tgcagactat cttttgagaa ctgggcaggt tgtagatata 480
teagacacea tataceegag gaaceetgee atgtattgtg aagaagetag attaaagtee 540
tttcagaact ggccagacta tgctcaccta accccaagag agttagcaag tgctggactc 600
tactacacag gtattggtga ccaagtgcag tgcttttgtt gtggtggaaa actgaaaaat 660
tgggaacctt gtgatcgtgc ctggtcagaa cacaggcgac actttcctaa ttgcttcttt 720
gttttgggcc ggaatcttaa tattcgaagt gaatctgatg ctgtgagttc tgataggaat 780
ttcccaaatt caacaaatct tccaagaaat ccatccatgg cagattatga agcacggatc 840
tttacttttg ggacatggat atactcagtt aacaaggagc agcttgcaag agctggattt 900
tatgctttag gtgaaggtga taaagtaaag tgctttcact gtggaggagg gctaactgat 960
tggaagccca gtgaagaccc ttgggaacaa catgctaaat ggtatccagg gtgcaaatat 1020
ctgttagaac agaagggaca agaatatata aacaatattc atttaactca ttcacttgag 1080
gagtgtctgg taagaactac tgagaaaaca ccatcactaa ctagaagaat tgatgatacc 1140
atcttccaaa atcctatggt acaagaagct atacgaatgg ggttcagttt caaggacatt 1200
aagaaaataa tggaggaaaa aattcagata tctgggagca actataaatc acttgaggtt 1260
ctggttgcag atctagtgaa tgctcagaaa gacagtatgc aagatgagtc aagtcagact 1320
tcattacaga aagagattag tactgaagag cagctaaggc gcctgcaaga ggagaagctt 1380
tgcaaaatct gtatggatag aaatattgct atcgtttttg ttccttgtgg acatctagtc 1440
acttgtaaac aatgtgctga agcagttgac aagtgtccca tgtgctacac agtcattact 1500
ttcaagcaaa aaatttttat gtcttaatct aactctatag taggcatgtt atgttgttct 1560
tattaccctg attgaatgtg tgatgtgaac tgactttaag taatcaggat tgaattccat 1620
tagcatttgc taccaagtag gaaaaaaat gtacatggca gtgttttagt tggcaatata 1680
atctttgaat ttcttgattt ttcagggtat tagctgtatt atccattttt tttactgtta 1740
tttaattgaa accatagact aagaataaga agcatcatac tataactgaa cacaatgtgt 1800
attcatagta tactgattta atttctaagt gtaagtgaat taatcatctg gattttttat 1860
tcttttcaga taggcttaac aaatggagct ttctgtatat aaatgtggag attagagtta 1920
atctccccaa tcacataatt tgttttgtgt gaaaaaggaa taaattgttc catgctggtg 1980
gaaagataga gattgttttt agaggttggt tgttgtgttt taggattctg tccattttct 2040
tgtaaaggga taaacacgga cgtgtgcgaa atatgtttgt aaagtgattt gccattgttg 2100
aaagcgtatt taatgataga atactatcga gccaacatgt actgacatgg aaagatgtca 2160
gagatatgtt aagtgtaaaa tgcaagtggc gggacactat gtatagtctg agccagatca 2220
aagtatgtat gttgttaata tgcatagaac gagagatttg gaaagatata caccaaactg 2280
ttaaatgtgg tttctcttcg gggagggggg gattggggga ggggccccag aggggtttta 2340
gaggggcctt ttcactttcg actttttca ttttgttctg ttcggatttt ttataagtat 2400
gtagaccccg aagggtttta tgggaactaa catcagtaac ctaacccccg tgactatcct 2460
gtgctcttcc tagggagctg tgttgtttcc cacccaccac ccttccctct gaacaaatgc 2520
ctgagtgctg gggcactttn
<210> 219
<211> 497
<212> PRT
<213> Homo sapiens
<400> 219
Met Thr Phe Asn Ser Phe Glu Gly Ser Lys Thr Cys Val Pro Ala Asp
                                   10
Ile Asn Lys Glu Glu Glu Phe Val Glu Glu Phe Asn Arg Leu Lys Thr
            20
                                25
```

Phe Ala Asn Phe Pro Ser Gly Ser Pro Val Ser Ala Ser Thr Leu Ala 40 Arg Ala Gly Phe Leu Tyr Thr Gly Glu Gly Asp Thr Val Arg Cys Phe Ser Cys His Ala Ala Val Asp Arg Trp Gln Tyr Gly Asp Ser Ala Val Gly Arg His Arg Lys Val Ser Pro Asn Cys Arg Phe Ile Asn Gly Phe 85 90 Tyr Leu Glu Asn Ser Ala Thr Gln Ser Thr Asn Ser Gly Ile Gln Asn 100 105 Gly Gln Tyr Lys Val Glu Asn Tyr Leu Gly Ser Arg Asp His Phe Ala 120 Leu Asp Arg Pro Ser Glu Thr His Ala Asp Tyr Leu Leu Arg Thr Gly 135 140 Gln Val Val Asp Ile Ser Asp Thr Ile Tyr Pro Arg Asn Pro Ala Met 150 155 Tyr Cys Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr 165 170 Ala His Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr 180 185 Gly Ile Gly Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys 195 200 205 Asn Trp Glu Pro Cys Asp Arg Ala Trp Ser Glu His Arg Arg His Phe 215 Pro Asn Cys Phe Phe Val Leu Gly Arg Asn Leu Asn Ile Arg Ser Glu 230 235 Ser Asp Ala Val Ser Ser Asp Arg Asn Phe Pro Asn Ser Thr Asn Leu 245 250-Pro Arg Asn Pro Ser Met Ala Asp Tyr Glu Ala Arg Ile Phe Thr Phe 260 265 Gly Thr Trp Ile Tyr Ser Val Asn Lys Glu Gln Leu Ala Arg Ala Gly 275 280 Phe Tyr Ala Leu Gly Glu Gly Asp Lys Val Lys Cys Phe His Cys Gly 295 300 Gly Gly Leu Thr Asp Trp Lys Pro Ser Glu Asp Pro Trp Glu Gln His 310 315 Ala Lys Trp Tyr Pro Gly Cys Lys Tyr Leu Leu Glu Gln Lys Gly Gln 325 330 Glu Tyr Ile Asn Asn Ile His Leu Thr His Ser Leu Glu Glu Cys Leu 340 345 Val Arg Thr Thr Glu Lys Thr Pro Ser Leu Thr Arg Arg Ile Asp Asp 355 360 Thr Ile Phe Gln Asn Pro Met Val Gln Glu Ala Ile Arg Met Gly Phe 375 380 Ser Phe Lys Asp Ile Lys Lys Ile Met Glu Glu Lys Ile Gln Ile Ser 390 395 Gly Ser Asn Tyr Lys Ser Leu Glu Val Leu Val Ala Asp Leu Val Asn 405 410 Ala Gln Lys Asp Ser Met Gln Asp Glu Ser Ser Gln Thr Ser Leu Gln 425 Lys Glu Ile Ser Thr Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Lys 440 Leu Cys Lys Ile Cys Met Asp Arg Asn Ile Ala Ile Val Phe Val Pro 455 Cys Gly His Leu Val Thr Cys Lys Gln Cys Ala Glu Ala Val Asp Lys 470 475 Cys Pro Met Cys Tyr Thr Val Ile Thr Phe Lys Gln Lys Ile Phe Met 485 490

Ser

```
<210> 220
<211> 2676
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> (1)...(2676)
<223> n=a,t,c, or g
<400> 220
tccttgagat gtatcagtat aggatttagg atctccatgt tggaactcta aatgcataga 60
aatggaaata atggaaattt ttcattttgg cttttcagcc tagtattaaa actgataaaa 120
gcaaagccat gcacaaaact acctccctag agaaaggcta gtcccttttc ttccccattc 180
atttcattat gaacatagta gaaaacagca tattcttatc aaatttgatg aaaagcgcca 240
acacgtttga actgaaatac gacttgtcat gtgaactgta ccgaatgtct acgtattcca 300
cttttcctgc tggggttcct gtctcagaaa ggagtcttgc tcgtgctggt ttctattaca 360
ctggtgtgaa tgacaaggtc aaatgcttct gttgtggcct gatgctggat aactggaaaa 420
gaggagacag tcctactgaa aagcataaaa agttgtatcc tagctgcaga ttcgttcaga 480
gtctaaattc cgttaacaac ttggaagcta cctctcagcc tacttttcct tcttcagtaa 540
cacattccac acactcatta cttccgggta cagaaaacag tggatatttc cgtggctctt 600
attcaaactc tccatcaaat cctgtaaact ccagagcaaa tcaagaattt tctgccttga 660
tgagaagttc ctacccctgt ccaatgaata acgaaaatgc cagattactt acttttcaga 720
catggccatt gacttttctg tcgccaacag atctggcacg agcaggcttt tactacatag 780
gacctggaga cagagtggct tgctttgcct gtggtggaaa attgagcaat tgggaaccga 840
aggataatgc tatgtcagaa cacctgagac attttcccaa atgcccattt atagaaaatc 900
agetteaaga caetteaaga tacacagttt etaatetgag catgeagaca catgeagece 960
getttaaaac attetttaac tggeeeteta gtgttetagt taateetgag cagettgcaa 1020
gtgcgggttt ttattatgtg ggtaacagtg atgatgtcaa atgcttttgc tgtgatggtg 1080
gactcaggtg ttgggaatct ggagatgatc catgggttca acatgccaag tggtttccaa 1140
ggtgtgagta cttgataaga attaaaggac aggagttcat ccgtcaagtt caagccagtt 1200
acceteatet acttgaacag etgetateea cateagacag eccaggagat gaaaatgeag 1260
agtcatcaat tatccatttg gaacctggag aagaccattc agaagatgca atcatgatga 1320
atactcctgt gattaatgct gccgtggaaa tgggctttag tagaagcctg gtaaaacaga 1380
cagttcagag aaaaatccta gcaactggag agaattatag actagtcaat gatcttgtgt 1440
tagacttact caatgcagaa gatgaaataa gggaagagga gagagaaaga gcaactgagg 1500
aaaaagaatc aaatgattta ttattaatcc ggaagaatag aatggcactt tttcaacatt 1560
tgacttgtgt aattccaatc ctggatagtc tactaactgc cggaattatt aatgaacaag 1620
aacatgatgt tattaaacag aagacacaga cgtctttaca agcaagagaa ctgattgata 1680
cgattttagt aaaaggaaat attgcagcca ctgtattcag aaactctctg caagaagctg 1740
aagctgtgtt atatgagcat ttatttgtgc aacaggacat aaaatatatt cccacagaag 1800
atgtttcaga tctaccagtg gaagaacaat tgcggagact accagaagaa agaacatgta 1860
aagtgtgtat ggacaaagaa gtgtccatag tgtttattcc ttgtggtcat ctagtagtat 1920
gcaaagattg tgctccttct ttaagaaagt gtcctatttg taggagtaca atcaagggta 1980
cagttcgtac atttcttca tgaagaagaa ccaaaacatc gtctaaactt tagaattaat 2040
ttattaaatg tattataact ttaactttta tcctaatttg gtttccttaa aatttttatt 2100
tatttacaac tcaaaaaaca ttgttttgtg taacatattt atatatgtat ctaaaccata 2160
tgaacatata ttttttagaa actaagagaa tgataggctt ttgttcttat gaacgaaaaa 2220
gaggtagcac tacaaacaca atattcaatc caaatttcag cattattgaa attgtaagtg 2280
aagtaaaact taagatattt gagttaacct ttaagaattt taaatatttt ggcattgtac 2340
taataccggg aacatgaagc caggtgtggt ggtatgtacc tgtagtccca ggctgaggca 2400
agagaattac ttgagcccag gagtttgaat ccatcctggg cagcatactg agaccctgcc 2460
tttaaaaaacn aacagnacca aanccaaaca ccagggacac atttctctgt cttttttgat 2520
cagtgtccta tacatcgaag gtgtgcatat atgttgaatc acattttagg gacatggtgt 2580
```

<210> 221

ttttataaag aattctgtga gnaaaattt aataaagcaa ccaaattact cttaaaaaaa 2640 aaaaaaaaaa aaaaactcg aggggcccgt accaat 2676

<211> 604 <212> PRT <213> Homo sapiens <400> 221 Met Asn Ile Val Glu Asn Ser Ile Phe Leu Ser Asn Leu Met Lys Ser 10 Ala Asn Thr Phe Glu Leu Lys Tyr Asp Leu Ser Cys Glu Leu Tyr Arg Met Ser Thr Tyr Ser Thr Phe Pro Ala Gly Val Pro Val Ser Glu Arg 35 40 Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys Val 55 Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Arg Gly Asp 70 Ser Pro Thr Glu Lys His Lys Lys Leu Tyr Pro Ser Cys Arg Phe Val 90 85 Gln Ser Leu Asn Ser Val Asn Asn Leu Glu Ala Thr Ser Gln Pro Thr 105 Phe Pro Ser Ser Val Thr His Ser Thr His Ser Leu Leu Pro Gly Thr 125 115 120 Glu Asn Ser Gly Tyr Phe Arg Gly Ser Tyr Ser Asn Ser Pro Ser Asn 135 140 Pro Val Asn Ser Arg Ala Asn Gln Glu Phe Ser Ala Leu Met Arg Ser 155 150 Ser Tyr Pro Cys Pro Met Asn Asn Glu Asn Ala Arg Leu Leu Thr Phe 170 175 165 Gln Thr Trp Pro Leu Thr Phe Leu Ser Pro Thr Asp Leu Ala Arg Ala 185 190 180 Gly Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys 205 200 195 Gly Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Asn Ala Met Ser Glu 220 210 215 His Leu Arg His Phe Pro Lys Cys Pro Phe Ile Glu Asn Gln Leu Gln 230 235 Asp Thr Ser Arg Tyr Thr Val Ser Asn Leu Ser Met Gln Thr His Ala 250 245 Ala Arg Phe Lys Thr Phe Phe Asn Trp Pro Ser Ser Val Leu Val Asn 265 260 Pro Glu Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Gly Asn Ser Asp 280 285 Asp Val Lys Cys Phe Cys Cys Asp Gly Gly Leu Arg Cys Trp Glu Ser 300 295 Gly Asp Asp Pro Trp Val Gln His Ala Lys Trp Phe Pro Arg Cys Glu 310 315 Tyr Leu Ile Arg Ile Lys Gly Gln Glu Phe Ile Arg Gln Val Gln Ala 330 335 Ser Tyr Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Ser Pro 345 340 Gly Asp Glu Asn Ala Glu Ser Ser Ile Ile His Leu Glu Pro Gly Glu 365 360 Asp His Ser Glu Asp Ala Ile Met Met Asn Thr Pro Val Ile Asn Ala 375 380 Ala Val Glu Met Gly Phe Ser Arg Ser Leu Val Lys Gln Thr Val Gln

```
395
                    390
385
Arg Lys Ile Leu Ala Thr Gly Glu Asn Tyr Arg Leu Val Asn Asp Leu
                                   410
Val Leu Asp Leu Leu Asn Ala Glu Asp Glu Ile Arg Glu Glu Glu Arg
                                425
            420
Glu Arg Ala Thr Glu Glu Lys Glu Ser Asn Asp Leu Leu Leu Ile Arg
                                                445
                            440
Lys Asn Arg Met Ala Leu Phe Gln His Leu Thr Cys Val Ile Pro Ile
                                            460
                       455
Leu Asp Ser Leu Leu Thr Ala Gly Ile Ile Asn Glu Gln Glu His Asp
                                        475
                    470
Val Ile Lys Gln Lys Thr Gln Thr Ser Leu Gln Ala Arg Glu Leu Ile
                                    490
                485
Asp Thr Ile Leu Val Lys Gly Asn Ile Ala Ala Thr Val Phe Arg Asn
                                505
            500
Ser Leu Gln Glu Ala Glu Ala Val Leu Tyr Glu His Leu Phe Val Gln
                                                525
                            520
Gln Asp Ile Lys Tyr Ile Pro Thr Glu Asp Val Ser Asp Leu Pro Val
                        535
                                            540
    530
Glu Glu Gln Leu Arg Arg Leu Pro Glu Glu Arg Thr Cys Lys Val Cys
                    550
                                        555
Met Asp Lys Glu Val Ser Ile Val Phe Ile Pro Cys Gly His Leu Val
                                                         575
                565
                                    570
Val Cys Lys Asp Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg
                                                     590
            580
                                585
Ser Thr Ile Lys Gly Thr Val Arg Thr Phe Leu Ser
                            600
        595
<210> 222
<211> 2580
<212> DNA
<213> Homo sapiens
<221> misc_feature
<222> (1)...(2580)
<223> n=a,t,c, or g
<400> 222
ttaggttacc tgaaagagtt actacaaccc caaagagttg tgttctaagt agtatcttgg 60
taattcagag agatactcat cctacctgaa tataaactga gataaatcca gtaaagaaag 120
tgtagtaaat tctacataag agtctatcat tgatttcttt ttgtggtgga aatcttagtt 180
catgtgaaga aatttcatgt gaatgtttta gctatcaaac agtactgtca cctactcatg 240
cacaaaactg cctcccaaag acttttccca ggtccctcgt atcaaaacat taagagtata 300
atggaagata gcacgatctt gtcagattgg acaaacagca acaaacaaaa aatgaagtat 360
gacttttcct gtgaactcta cagaatgtct acatattcaa ctttccccgc cggggtgcct 420
gtctcagaaa ggagtcttgc tcgtgctggt ttttattata ctggtgtgaa tgacaaggtc 480
aaatgettet gttgtggeet gatgetggat aactggaaac taggagacag teetatteaa 540
aagcataaac agctatatcc tagctgtagc tttattcaga atctggtttc agctagtctg 600
ggatccacct ctaagaatac gtctccaatg agaaacagtt ttgcacattc attatctccc 660
accttggaac atagtagett gttcagtggt tettaeteca geetteetee aaaccetett 720
aattctagag cagttgaaga catctcttca tcgaggacta acccctacag ttatgcaatg 780
agtactgaag aagccagatt tottacotac catatgtggc cattaacttt titgtcacca 840
tcagaattgg caagagctgg tttttattat ataggacctg gagatagggt agcctgcttt 900
gcctgtggtg ggaagctcag taactgggaa ccaaaggatg atgctatgtc agaacaccgg 960
aggeatttte ceaactgtee atttttggaa aattetetag aaactetgag gtttageatt 1020
tcaaatetga gcatgcagac acatgcaget cgaatgagaa catttatgta etggccatet 1080
```

```
agtgttccag ttcagcctga gcagcttgca agtgctggtt tttattatgt gggtcgcaat 1140
gatgatgtca aatgctttgg ttgtgatggt ggcttgaggt gttgggaatc tggagatgat 1200
ccatgggtag aacatgccaa gtggtttcca aggtgtgagt tcttgatacg aatgaaaggc 1260
caagagtttg ttgatgagat tcaaggtaga tatcctcatc ttcttgaaca gctgttgtca 1320
acttcagata ccactggaga agaaaatgct gacccaccaa ttattcattt tggacctgga 1380
gaaagttett cagaagatge tgteatgatg aatacacetg tggttaaate tgeettggaa 1440
atgggcttta atagagacct ggtgaaacaa acagttctaa gtaaaatcct gacaactgga 1500
gagaactata aaacagttaa tgatattgtg tcagcacttc ttaatgctga agatgaaaaa 1560
agagaagagg agaaggaaaa acaagctgaa gaaatggcat cagatgattt gtcattaatt 1620
cggaagaaca gaatggctct ctttcaacaa ttgacatgtg tgcttcctat cctggataat 1680
cttttaaagg ccaatgtaat taataaacag gaacatgata ttattaaaca aaaaacacag 1740
atacctttac aagcgagaga actgattgat accatttggg ttaaaggaaa tgctgcggcc 1800
aacatcttca aaaactgtct aaaagaaatt gactctacat tgtataagaa cttatttgtg 1860
gataagaata tgaagtatat tccaacagaa gatgtttcag gtctgtcact ggaagaacaa 1920
ttgaggaggt tgcaagaaga acgaacttgt aaagtgtgta tggacaaaga agtttctgtt 1980
gtatttattc cttgtggtca tctggtagta tgccaggaat gtgccccttc tctaagaaaa 2040
tgccctattt gcaggggtat aatcaagggt actgttcgta catttctctc ttaaagaaaa 2100
atagtetata tittaaeetg cataaaaagg tetttaaaat attgttgaae aettgaagee 2160
atctaaagta aaaagggaat tatgagtttt tcaattagta acattcatgt tctagtctgc 2220
tttggtacta ataatcttgt ttctgaaaag atggtatcat atatttaatc ttaatctgtt 2280
tatttacaag ggaagattta tgtttggtga actatattag tatgtatgtg tacctaaggg 2340
agtagcgtcn ctgcttgtta tgcatcattt caggagttac tggatttgtt gttctttcag 2400
aaagetttga anactaaatt atagtgtaga aaagaactgg aaaccaggaa etetggagtt 2460
catcagagtt atggtgccga attgtctttg gtgcttttca cttgtgtttt aaaataagga 2520
tttttctctt atttctcccc ctagtttgtg agaaacatct caataaagtg ctttaaaaag 2580
<210> 223
<211> 618
<212> PRT
<213> Homo sapiens
<400> 223
Met His Lys Thr Ala Ser Gln Arg Leu Phe Pro Gly Pro Ser Tyr Gln
                                    10
Asn Ile Lys Ser Ile Met Glu Asp Ser Thr Ile Leu Ser Asp Trp Thr
            20
                                25
Asn Ser Asn Lys Gln Lys Met Lys Tyr Asp Phe Ser Cys Glu Leu Tyr
        35
Arg Met Ser Thr Tyr Ser Thr Phe Pro Ala Gly Val Pro Val Ser Glu
                        55
Arg Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys
65
                    70
Val Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Leu Gly
                85
                                    90
Asp Ser Pro Ile Gln Lys His Lys Gln Leu Tyr Pro Ser Cys Ser Phe
                                105
Ile Gln Asn Leu Val Ser Ala Ser Leu Gly Ser Thr Ser Lys Asn Thr
        115
                            120
                                                125
Ser Pro Met Arg Asn Ser Phe Ala His Ser Leu Ser Pro Thr Leu Glu
    130
                        135
                                            140
His Ser Ser Leu Phe Ser Gly Ser Tyr Ser Ser Leu Pro Pro Asn Pro
                    150
                                        155
Leu Asn Ser Arg Ala Val Glu Asp Ile Ser Ser Ser Arg Thr Asn Pro
                165
                                    170
                                                        175
Tyr Ser Tyr Ala Met Ser Thr Glu Glu Ala Arg Phe Leu Thr Tyr His
                                185
```

Met Trp Pro Leu Thr Phe Leu Ser Pro Ser Glu Leu Ala Arg Ala Gly

```
200
      195
Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys Gly
                     215
                                       220
Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Asp Ala Met Ser Glu His
              230 .235
Arg Arg His Phe Pro Asn Cys Pro Phe Leu Glu Asn Ser Leu Glu Thr
                      250
             245
Leu Arg Phe Ser Ile Ser Asn Leu Ser Met Gln Thr His Ala Ala Arg
          260
                           265
Met Arg Thr Phe Met Tyr Trp Pro Ser Ser Val Pro Val Gln Pro Glu
                                        285
       275
                        280
Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Gly Arg Asn Asp Asp Val
                    295
                                       300
Lys Cys Phe Gly Cys Asp Gly Gly Leu Arg Cys Trp Glu Ser Gly Asp
                310
                                   315
Asp Pro Trp Val Glu His Ala Lys Trp Phe Pro Arg Cys Glu Phe Leu
              325
                               330
Ile Arg Met Lys Gly Gln Glu Phe Val Asp Glu Ile Gln Gly Arg Tyr
       340
                345
Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Thr Thr Gly Glu
 355 360
Glu Asn Ala Asp Pro Pro Ile Ile His Phe Gly Pro Gly Glu Ser Ser
                     375
                                       380
Ser Glu Asp Ala Val Met Met Asn Thr Pro Val Val Lys Ser Ala Leu
                 390
                                  395
Glu Met Gly Phe Asn Arg Asp Leu Val Lys Gln Thr Val Leu Ser Lys
              405
                               410
Ile Leu Thr Thr Gly Glu Asn Tyr Lys Thr Val Asn Asp Ile Val Ser
          420
                            425
                                              430
Ala Leu Leu Asn Ala Glu Asp Glu Lys Arg Glu Glu Glu Lys Glu Lys
   435
                        440
Gln Ala Glu Glu Met Ala Ser Asp Asp Leu Ser Leu Ile Arg Lys Asn
                              460
                     455
Arg Met Ala Leu Phe Gln Gln Leu Thr Cys Val Leu Pro Ile Leu Asp
                  470
                                   475
Asn Leu Leu Lys Ala Asn Val Ile Asn Lys Gln Glu His Asp Ile Ile
             485
                               490
Lys Gln Lys Thr Gln Ile Pro Leu Gln Ala Arg Glu Leu Ile Asp Thr
                            505
          500
Ile Trp Val Lys Gly Asn Ala Ala Ala Asn Ile Phe Lys Asn Cys Leu
                                        525
                        520
Lys Glu Ile Asp Ser Thr Leu Tyr Lys Asn Leu Phe Val Asp Lys Asn
                   535
Met Lys Tyr Ile Pro Thr Glu Asp Val Ser Gly Leu Ser Leu Glu Glu
         550
                           555
Gln Leu Arg Arg Leu Gln Glu Glu Arg Thr Cys Lys Val Cys Met Asp
                                 570
Lys Glu Val Ser Val Val Phe Ile Pro Cys Gly His Leu Val Val Cys
                             585
Gln Glu Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg Gly Ile
       595 600
Ile Lys Gly Thr Val Arg Thr Phe Leu Ser
```

<210> 224

<211> 2100

<212> DNA

<213> Mus musculus

<400> 224 gacactetge tgggeggegg geegeeetee teegggaeet eeeeteggga aeegtegeee 60 gcggcgctta gttaggactg gagtgcttgg cgcgaaaagg tggacaagtc ctattttcca 120 gagaagatga cttttaacag ttttgaagga actagaactt ttgtacttgc agacaccaat 180 aaggatgaag aatttgtaga agagtttaat agattaaaaa catttgctaa cttcccaagt 240 agtagtcctg tttcagcatc aacattggcg cgagctgggt ttctttatac cggtgaagga 300 gacaccgtgc aatgtttcag ttgtcatgcg gcaatagata gatggcagta tggagactca 360 gctgttggaa gacacaggag aatatcccca aattgcagat ttatcaatgg tttttatttt 420 gaaaatggtg ctgcacagtc tacaaatcct ggtatccaaa atggccagta caaatctgaa 480 aactgtgtgg gaaatagaaa teettttgee eetgacagge caeetgagae teatgetgat 540 tatctcttga gaactggaca ggttgtagat atttcagaca ccatataccc gaggaaccct 600 gccatgtgta gtgaagaagc cagattgaag tcatttcaga actggccgga ctatgctcat 660 ttaaccccca gagagttagc tagtgctggc ctctactaca caggggctga tgatcaagtg 720 caatgetttt gttgtggggg aaaactgaaa aattgggaac cetgtgatcg tgcctggtca 780 gaacacagga gacactttcc caattgcttt tttgttttgg gccggaacgt taatgttcga 840 agtgaatctg gtgtgagttc tgataggaat ttcccaaatt caacaaactc tccaagaaat 900 ccagccatgg cagaatatga agcacggatc gttacttttg gaacatggat atactcagtt 960 aacaaggagc agcttgcaag agctggattt tatgctttag gtgaaggcga taaagtgaag 1020 tgcttccact gtggaggagg gctcacggat tggaagccaa gtgaagaccc ctgggaccag 1080 catgctaagt gctacccagg gtgcaaatac ctattggatg agaaggggca agaatatata 1140 aataatattc atttaaccca tccacttgag gaatctttgg gaagaactgc tgaaaaaaca 1200 ccaccgctaa ctaaaaaaat cgatgatacc atcttccaga atcctatggt gcaagaagct 1260 atacgaatgg gatttagctt caaggacctt aagaaaacaa tggaagaaaa aatccaaaca 1320 tccgggagca gctatctatc acttgaggtc ctgattgcag atcttgtgag tgctcagaaa 1380 gataatacgg aggatgagtc aagtcaaact tcattgcaga aagacattag tactgaagag 1440 cagctaaggc gcctacaaga ggagaagctt tccaaaatct gtatggatag aaatattgct 1500 atcgtttttt ttccttgtgg acatctggcc acttgtaaac agtgtgcaga agcagttgac 1560 aaatgtccca tgtgctacac cgtcattacg ttcaaccaaa aaatttttat gtcttagtgg 1620 ggcaccacat gttatgttct tcttgctcta attgaatgtg taatgggagc gaactttaag 1680 taatcctgca tttgcattcc attagcatcc tgctgtttcc aaatggagac caatgctaac 1740 agcactgttt ccgtctaaac attcaatttc tggatctttc gagttatcag ctgtatcatt 1800 tagccagtgt tttactcgat tgaaacctta gacagagaag cattttatag cttttcacat 1860 gtatattggt agtacactga cttgatttct atatgtaagt gaattcatca cctgcatgtt 1920 tcatgccttt tgcataagct taacaaatgg agtgttctgt ataagcatgg agatgtgatg 1980 gaatctgccc aatgacttta attggcttat tgtaaacacg gaaagaactg ccccacgctg 2040 ctgggaggat aaagattgtt ttagatgctc acttctgtgt tttaggattc tgcccattta 2100

<210> 225

<211> 496

<212> PRT

<213> Mus musculus

<400> 225

 Met
 Thr
 Phe
 Asn
 Ser
 Phe
 Glu
 Gly
 Thr
 Arg
 Thr
 Phe
 Val
 Leu
 Ala
 Asp

 Thr
 Asn
 Lys
 Asp
 Glu
 Glu
 Phe
 Val
 Glu
 Phe
 Asn
 Arg
 Leu
 Lys
 Thr

 Phe
 Ala
 Asn
 Phe
 Pro
 Ser
 Ser
 Ser
 Pro
 Val
 Ser
 Ala
 Ser
 Thr
 Leu
 Ala
 ```
Tyr Phe Glu Asn Gly Ala Ala Gln Ser Thr Asn Pro Gly Ile Gln Asn
                            105
                                              110
         100
Gly Gln Tyr Lys Ser Glu Asn Cys Val Gly Asn Arg Asn Pro Phe Ala
                        120
                                         125
      115
Pro Asp Arg Pro Pro Glu Thr His Ala Asp Tyr Leu Leu Arg Thr Gly
                               . 140
          135
Gln Val Val Asp Ile Ser Asp Thr Ile Tyr Pro Arg Asn Pro Ala Met
         150 155
Cys Ser Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr
             165
                               170
                                                 175
Ala His Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr
                          185
         180
Gly Ala Asp Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys
                        200
                                          205
Asn Trp Glu Pro Cys Asp Arg Ala Trp Ser Glu His Arg Arg His Phe
                     215
                                       220
Pro Asn Cys Phe Phe Val Leu Gly Arg Asn Val Asn Val Arg Ser Glu
                            235
                230
Ser Gly Val Ser Ser Asp Arg Asn Phe Pro Asn Ser Thr Asn Ser Pro
                             250
                                                 255
             245
Arg Asn Pro Ala Met Ala Glu Tyr Glu Ala Arg Ile Val Thr Phe Gly
                         265
          260
Thr Trp Ile Tyr Ser Val Asn Lys Glu Gln Leu Ala Arg Ala Gly Phe
                        280
                                         285
       275
Tyr Ala Leu Gly Glu Gly Asp Lys Val Lys Cys Phe His Cys Gly Gly
                   295
                                      300
Gly Leu Thr Asp Trp Lys Pro Ser Glu Asp Pro Trp Asp Gln His Ala
                310
                                   315
Lys Cys Tyr Pro Gly Cys Lys Tyr Leu Leu Asp Glu Lys Gly Gln Glu
                            330
                                                 335
              325
Tyr Ile Asn Asn Ile His Leu Thr His Pro Leu Glu Glu Ser Leu Gly
        340 345
Arg Thr Ala Glu Lys Thr Pro Pro Leu Thr Lys Lys Ile Asp Asp Thr
    355 360
                                365
Ile Phe Gln Asn Pro Met Val Gln Glu Ala Ile Arg Met Gly Phe Ser
   370 375
                                       380
Phe Lys Asp Leu Lys Lys Thr Met Glu Glu Lys Ile Gln Thr Ser Gly
               390
                                   395
Ser Ser Tyr Leu Ser Leu Glu Val Leu Ile Ala Asp Leu Val Ser Ala
                               410
                                                 415
             405
Gln Lys Asp Asn Thr Glu Asp Glu Ser Ser Gln Thr Ser Leu Gln Lys
                                             430
          420
                            425
Asp Ile Ser Thr Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Lys Leu
    435
                       440
                                          445
Ser Lys Ile Cys Met Asp Arg Asn Ile Ala Ile Val Phe Phe Pro Cys
  450 455
                                       460
Gly His Leu Ala Thr Cys Lys Gln Cys Ala Glu Ala Val Asp Lys Cys
        470
                         475
Pro Met Cys Tyr Thr Val Ile Thr Phe Asn Gln Lys Ile Phe Met Ser
                                490
              485
```

<210> 226

<211> 2474

<212> DNA

<213> Mus musculus

<400> 226

```
gaattccggg agacctacac ccccggagat cagaggtcat tgctggcgtt cagagcctag 60
gaagtgggct gcggtatcag cctagcagta aaaccgacca gaagccatgc acaaaactac 120
atccccagag aaagacttgt cocttcccct coctgtcate teaccatgaa catggttcaa 180
gacagegeet ttetageeaa getgatgaag agtgetgaca eetttgagtt gaagtatgae 240
ttttcctgtg agctgtaccg attgtccacg tattcagctt ttcccagggg agttcctgtg 300
tcagaaagga gtctggctcg tgctggcttt tactacactg gtgccaatga caaggtcaag 360
tgcttctgct gtggcctgat gctagacaac tggaaacaag gggacagtcc catggagaag 420
cacagaaagt tgtaccccag ctgcaacttt gtacagactt tgaatccagc caacagtctg 480
gaagctagte cleggeette tetteettee acggegatga geaccatgee tittgagettt 540
gcaagttctg agaatactgg ctatttcagt ggctcttact cgagctttcc ctcagaccct 600
gtgaacttcc gagcaaatca agattgtcct gctttgagca caagtcccta ccactttgca 660
atgaacacag agaaggccag attactcacc tatgaaacat ggccattgtc ttttctgtca 720
ccagcaaagc tggccaaagc aggcttctac tacataggac ctggagatag agtggcctgc 780
tttgcgtgcg atgggaaact gagcaactgg gaacgtaagg atgatgctat gtcagagcac 840
cagaggeatt tececagetg teegttetta aaagaettgg gteagtetge ttegagatae 900
actgtctcta acctgagcat gcagacacac gcagcccgta ttagaacatt ctctaactgg 960
ccttctagtg cactagttca ttcccaggaa cttgcaagtg cgggctttta ttatacagga 1020
cacagtgatg atgtcaagtg tttatgctgt gatggtgggc tgaggtgctg ggaatctgga 1080
gatgacccct gggtggaaca tgccaagtgg tttccaaggt gtgagtactt gctcagaatc 1140
aaaggccaag aatttgtcag ccaagttcaa gctggctatc ctcatctact tgagcagcta 1200
ttatctacgt cagactcccc agaagatgag aatgcagacg cagcaatcgt gcattttggc 1260
cctggagaaa gttcggaaga tgtcgtcatg atgagcacgc ctgtggttaa agcagccttg 1320
gaaatgggct tcagtaggag cctggtgaga cagacggttc agtggcagat cctggccact 1380
ggtgagaact acaggaccgt cagtgacctc gttataggct tactcgatgc agaagacgag 1440
atgagagagg agcagatgga gcaggcggcc gaggaggagg agtcagatga tctagcacta 1500
atcoggaaga acaaaatggt gcttttccaa catttgacgt gtgtgacacc aatgctgtat 1560
tgcctcctaa gtgcaagggc catcactgaa caggagtgca atgctgtgaa acagaaacca 1620
cacaccttac aagcaagcac actgattgat actgtgttag caaaaggaaa cactgcagca 1680
acctcattca gaaactccct tcgggaaatt gaccctgcgt tatacagaga tatatttgtg 1740
caacaggaca ttaggagtct tcccacagat gacattgcag ctctaccaat ggaagaacag 1800
ttgcggcccc tcccggagga cagaatgtgt aaagtgtgta tggaccgaga ggtatccatc 1860
gtgttcattc cctgtggcca tctggtcgtg tgcaaagact gcgctccctc tctgaggaag 1920
tgtcccatct gtagagggac catcaagggc acagtgcgca catttctctc ctgaacaaga 1980
ctaatggtcc atggctgcaa cttcagccag gaggaagttc actgtcactc ccagttccat 2040
teggaacttg aggecageet ggatageaeg agacacegee aaacacacaa atataaacat 2100
gaaaaacttt tgtctgaagt caagaatgaa tgaattactt atataataat tttaattggt 2160
ttccttaaaa gtgctatttg ttcccaactc agaaaattgt tttctgtaaa catatttaca 2220
tactacctgc atctaaagta ttcatatatt catatattca gatgtcatga gagagggttt 2280
tgttcttgtt cctgaaaagc tggtttatca tctgatcagc atatactgcg caacgggcag 2340
ggctagaatc catgaaccaa gctgcaaaga tctcacgcta aataaggcgg aaagatttgg 2400
agaaacgaaa ggaaattott tootgtocaa tgtatactot toagactaat gacctottoo 2460
                                                                   2474
tatcaagcct tcta
```

<210> 227

<211> 602

<212> PRT

<213> Mus musculus

<400> 227

 Met
 Asn
 Met
 Val
 Gln
 Asp
 Ser
 Ala
 Phe
 Leu
 Ala
 Lys
 Leu
 Met
 Lys
 Ser
 Ala
 Phe
 Leu
 Ala
 Phe
 Leu
 Lys
 Tyr
 Asp
 Phe
 Ser
 Cys
 Glu
 Leu
 Tyr
 Arg

 Ala
 Asp
 Ala
 Phe
 Pro
 Arg
 Gly
 Val
 Pro
 Val
 Ser
 Glu
 Arg
 Ala
 Arg
 Arg
 Ala
 Arg
65					70					75					80
Ser	Pro	Met	Glu	Lys 85	His	Arg	Lys	Leu	Tyr 90	Pro	Ser	Cys	Asn	Phe 95	Val
			100					105					Arg 110		
		115					120					125	Ala		
	130					135					140		Pro		
145					150					155			Ser		160
				165					170				Leu	175	
			180					185					Ala 190		
		195					200					205	Phe		
_	210					215					220		Met		
225					230					235			Leu		240
				245					250				Thr	255	
			260					265					Leu 270		
		275					280					285	His		
	290					295					300		Trp		
305					310					315			Arg		320
_				325					330				Val Asp	335	
_	_		340					345					350 Pro		
		355					360					365	Lys		
	370					375					380		Val		
385					390					395			Asp		400
				405					410				Gln	415	
			420					425					430 . Ile		
		435	i				440					445			
	450)				455					460		Cys		
465	_				470	i				475					480
	_			485	;				490)			Ile	495	
			500)				505	i				Asn 510		
		515	5				520					525			
Ile	arç	g Sei	Lev	Pro	Thr	Asp) Asp) Ile	Ala	Ala	. Leu	Pro	Met	Glu	Glu

540

535

530

```
Gln Leu Arg Pro Leu Pro Glu Asp Arg Met Cys Lys Val Cys Met Asp
                    550
                                        555
Arg Glu Val Ser Ile Val Phe Ile Pro Cys Gly His Leu Val Val Cys
                565
                                    570
                                                        575
Lys Asp Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg Gly Thr
            580
                                585
                                                    590
Ile Lys Gly Thr Val Arg Thr Phe Leu Ser
<210> 228
<211> 2416
<212> DNA
<213> Mus musculus
<400> 228
ctgtggtgga gatctattgt ccaagtggtg agaaacttca tctggaagtt taagcggtca 60
gaaatactat tactactcat ggacaaaact gtctcccaga gactcgccca aggtacctta 120
cacccaaaaa cttaaacgta taatggagaa gagcacaatc ttgtcaaatt ggacaaagga 180
gagcgaagaa aaaatgaagt ttgacttttc gtgtgaactc taccgaatgt ctacatattc 240
agcttttccc aggggagttc ctgtctcaga gaggagtctg gctcgtgctg gcttttatta 300
tacaggtgtg aatgacaaag tcaagtgctt ctgctgtggc ctgatgttgg ataactggaa 360
acaaggggac agtcctgttg aaaagcacag acagttctat cccagctgca gctttgtaca 420
gactotgott toagocagto tgcagtotoc atotaagaat atgtotootg tgaaaagtag 480
atttgcacat tcgtcacctc tggaacgagg tggcattcac tccaacctgt gctctagccc 540
tettaattet agageagtgg aagaettete ateaaggatg gateeetgea getatgeeat 600
gagtacagaa gaggccagat ttcttactta cagtatgtgg cctttaagtt ttctgtcacc 660
agcagagctg gccagagctg gcttctatta catagggcct ggagacaggg tggcctgttt 720
tgcctgtggt gggaaactga gcaactggga accaaaggat tatgctatgt cagagcaccg 780
cagacatttt ccccactgtc catttctgga aaatacttca gaaacacaga ggtttagtat 840
atcaaatcta agtatgcaga cacactctgc tcgattgagg acatttctgt actggccacc 900
tagtgttcct gttcagcccg agcagcttgc aagtgctgga ttctattacg tggatcgcaa 960
tgatgatgtc aagtgccttt gttgtgatgg tggcttgaga tgttgggaac ctggagatga 1020
cccctggata gaacacgcca aatggtttcc aaggtgtgag ttcttgatac ggatgaaggg 1080
teaggagttt gttgatgaga tteaagetag atateeteat ettettgage agetgttgte 1140
cacttcagac accccaggag aagaaaatgc tgaccctaca gagacagtgg tgcattttgg 1200
ccctggagaa agttcgaaag atgtcgtcat gatgagcacg cctgtggtta aagcagcctt 1260
ggaaatgggc ttcagtagga gcctggtgag acagacggtt cagcggcaga tcctggccac 1320
tggtgagaac tacaggaccg tcaatgatat tgtctcagta cttttgaatg ctgaagatga 1380
gagaagagaa gaggagaagg aaagacagac tgaagagatg gcatcaggtg acttatcact 1440
gattcggaag aatagaatgg ccctctttca acagttgaca catgtccttc ctatcctgga 1500
taatcttctt gaggccagtg taattacaaa acaggaacat gatattatta gacagaaaac 1560
acagataccc ttacaagcaa gagagcttat tgacaccgtt ttagtcaagg gaaatgctgc 1620
agccaacatc ttcaaaaact ctctgaaggg aattgactcc acgttatatg aaaacttatt 1680
tgtggaaaag aatatgaagt atattccaac agaagacgtt tcaggcttgt cattggaaga 1740
gcagttgcgg agattacaag aagaacgaac ttgcaaagtg tgtatggaca gagaggtttc 1800
tattgtgttc attccgtgtg gtcatctagt agtctgccag gaatgtgccc cttctctaag 1860
gaagtgcccc atctgcaggg ggacaatcaa ggggactgtg cgcacatttc tctcatgagt 1920
gaagaatggt ctgaaagtat tgttggacat cagaagctgt cagaacaaag aatgaactac 1980
tgatttcagc tcttcagcag gacattctac tctctttcaa gattagtaat cttgctttat 2040
gaagggtagc attgtatatt taagcttagt ctgttgcaag ggaaggtcta tgctgttgag 2100
ctacaggact gtgtctgttc cagagcagga gttgggatgc ttgctgtatg tccttcagga 2160
cttcttggga tttgggaatt tggggaaagc tttggaatcc agtgatgtgg agctcagaaa 2220
tectggaace agtgactetg gtactcagta gatagggtac cetgtactte ttggtgettt 2280
tccagtctgg gaaataagga ggaatctgct gctggtaaaa atttgctgga tgtgagaaat 2340
agatgaaagt gtttcgggtg ggggcgtgca tcagtgtagt gtgtgcaggg atgtatgcag 2400
gccaaacact gtgtag
                                                                  2416
```

<210> 229 <211> 591

<212> PRT <213> Mus musculus <400> 229 Met Glu Lys Ser Thr Ile Leu Ser Asn Trp Thr Lys Glu Ser Glu Glu Lys Met Lys Phe Asp Phe Ser Cys Glu Leu Tyr Arg Met Ser Thr Tyr Ser Ala Phe Pro Arg Gly Val Pro Val Ser Glu Arg Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys Val Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Gln Gly Asp Ser Pro Val Glu Lys His Arg Gln Phe Tyr Pro Ser Cys Ser Phe Val Gln Thr Leu Leu Ser Ala Ser Leu Gln Ser Pro Ser Lys Asn Met Ser Pro Val Lys Ser Arg Phe Ala His Ser Ser Pro Leu Glu Arg Gly Gly Ile His Ser Asn Leu Cys Ser Ser Pro Leu Asn Ser Arg Ala Val Glu Asp Phe Ser Ser Arg Met Asp Pro Cys Ser Tyr Ala Met Ser Thr Glu Glu Ala Arg Phe Leu Thr Tyr Ser Met Trp Pro Leu Ser Phe Leu Ser Pro Ala Glu Leu Ala Arg Ala Gly Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys Gly Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Tyr Ala Met Ser Glu His Arg Arg His Phe Pro His Cys Pro Phe Leu Glu Asn Thr Ser Glu Thr Gln Arg Phe Ser Ile Ser Asn Leu Ser Met Gln Thr His Ser Ala Arg Leu Arg Thr Phe Leu Tyr Trp Pro Pro Ser Val Pro Val Gln Pro Glu Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Asp Arg Asn Asp Asp Val Lys Cys Leu Cys Cys Asp Gly Gly Leu Arg Cys Trp Glu Pro Gly Asp Asp Pro Trp Ile Glu His Ala Lys Trp Phe Pro Arg Cys Glu Phe Leu Ile Arg Met Lys Gly Gln Glu Phe Val Asp Glu Ile Gln Ala Arg Tyr Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Thr Pro Gly Glu Glu Asn Ala Asp Pro Thr Glu Thr Val Val His Phe Gly Pro Gly Glu Ser Ser Lys Asp Val Val Met Met Ser Thr Pro Val Val Lys Ala Ala Leu Glu Met Gly Phe Ser Arg Ser Leu Val Arg Gln Thr Val Gln Arg Gln Ile Leu Ala Thr Gly Glu Asn Tyr Arg Thr Val Asn Asp Ile Val Ser Val Leu Leu Asn Ala Glu Asp Glu Arg Arg Glu

```
Glu Glu Lys Glu Arg Gln Thr Glu Glu Met Ala Ser Gly Asp Leu Ser
                               425
                                                   430
           420
Leu Ile Arg Lys Asn Arg Met Ala Leu Phe Gln Gln Leu Thr His Val
                           440
                                               445
Leu Pro Ile Leu Asp Asn Leu Leu Glu Ala Ser Val Ile Thr Lys Gln
                       455
Glu His Asp Ile Ile Arg Gln Lys Thr Gln Ile Pro Leu Gln Ala Arg
                    470
                                       475
Glu Leu Ile Asp Thr Val Leu Val Lys Gly Asn Ala Ala Ala Asn Ile
                485
                                   490
Phe Lys Asn Ser Leu Lys Gly Ile Asp Ser Thr Leu Tyr Glu Asn Leu
                                                   510
                               505
           500
Phe Val Glu Lys Asn Met Lys Tyr Ile Pro Thr Glu Asp Val Ser Gly
                           520
                                             · 525
        515
Leu Ser Leu Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Arg Thr Cys
                       535
                                           540
    530
Lys Val Cys Met Asp Arg Glu Val Ser Ile Val Phe Ile Pro Cys Gly
                    550
                                       555
His Leu Val Val Cys Gln Glu Cys Ala Pro Ser Leu Arg Lys Cys Pro
                565
                                   570
Ile Cys Arg Gly Thr Ile Lys Gly Thr Val Arg Thr Phe Leu Ser
            580
                               585
<210> 230
<211> 6669
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> (1)...(6669)
<223> n=a,t,c, or g
<400> 230
ttgctctgtc acccagtttg gagtgcagtt atgcagtctc acactgcaag ctctgcctca 60
tgggctcaag tgaacctcct gcctcagcct ctcaagtagc tgggaccaca ggcaggtgcc 120
accatgtctg gctaattttt gagtttcttt gtagagatgg tgttttgcca agtcacccag 180
tttgaggetg gteteaaaca eetgggetea ageaateeat etaceteage eteceaaagt 240
gctgggatta caggagtgag ccatggcatg aggccttgtg gggtgtctct tttaaatgaa 300
agcatactct gtttacgtat ttgatatgaa ggaatatcct tcctttccac aaagacaaaa 360
attatectat tttteteaaa acatatgtee ttttteteta etttteattt ttgttaettt 420
tgatggacac atgtgttaca ttgatttcac tttctcataa ttctgctgta agaaaaacaa 480
tagtgccagt tcaatgacaa atagcaacag tctgttattg ctagactgtt actgttagtg 540
gagactacca gaacagtcag tcccagtgtc agggaatcaa agagaacatg ttccctctct 600
aaagggcaca gctgctgctc agctttagct gattgctgcc ctgcaggact ataggcccag 660
tgttgctaga tcttttgatg tttcaagaga agcttggaat ctagaatgtg atgggaagtc 720
tcttacattt aaacatgttg gcaattaatg gtaagattta aaaatactgt ggtccaagaa 780
aaaaatggat ttggaaactg gattaaattc aaatgaggca tgcagattaa tctacagcat 840
ggtacaatgt gaattttctg gtttctttaa ttgcactgta attaggtaag atgttagctt 900
tggggaaget aagtgeagag tatgeagaaa etattatttt tgtaagtttt etetaagtat 960
aaataaattt caaaataaaa ataaaaactt agtaaagaac tataatgcaa ttctatgtaa 1020
tttgagacag agtcttgctg tgtcacccag gctggagtgt agtggcacta tttcggccca 1140
ctgcaacctc cacctcccag gctcaaatga ttctcctgcc tcagcctccg gagtagctgg 1200
gattacaggc gcgtaccacc acacccagct aatttttgta tttttagtag agatggggtt 1260
tcaccatttt ggccaggctg gttttgaact cctgacctca agtgatccac ttgtcttggc 1320
ctcccaaaat gctgggatta caggcgtgag ccactgcacc aggcagaggc ctctgttttt 1380
```

tatetetttt tggeetetae agtgeetagt aaageacetg atacatggta aaegateagt 1440 aattactagt actctatttt ggagaaaatg attttttaaa aagtcattgt gttccatcca 1500 tgaqtcqttt gaqttttaaa actgtctttt tgtttgtttt tgaacaggtt tacaaaggag 1560 gaaaacgact tcttctagat ttttttttca gtttcttcta taaatcaaaa catctcaaaa 1620 tggagaccta aaatccttaa agggacttag tctaatctcg ggaggtagtt ttgtgcatgg 1680 gtaaacaaat taagtattaa ctggtgtttt actatccaaa gaatgctaat tttataaaca 1740 tgatcgagtt atataaggta taccataatg agtttgattt tgaatttgat ttgtggaaat 1800 aaaggaaaag tgattctagc tggggcatat tgttaaagca tttttttcag agttggccag 1860 gcagtctcct actggcacat tctcccatta tgtagaatag aaatagtacc tgtgtttggg 1920 aaagatttta aaatgagtga cagttatttg gaacaaagag ctaataatca atccactgca 1980 aattaaagaa acatgcagat gaaagttttg acacattaaa atacttctac agtgacaaag 2040 aaaaatcaag aacaaagctt tttgatatgt gcaacaaatt tagaggaagt aaaaagataa 2100 atgtgatgat tggtcaagaa attatccagt tatttacaag gccactgata ttttaaacgt 2160 ccaaaagttt qtttaaatqq qctqttaccg ctgagaatga tgaggatgag aatgatggtt 2220 gaaggttaca ttttaggaaa tgaagaaact tagaaaatta atataaagac agtgatgaat 2280 acaaagaaga tttttataac aatgtgtaaa atttttggcc agggaaagga atattgaagt 2340 tagatacaat tacttacctt tgagggaaat aattgttggt aatgagatgt gatgtttctc 2400 ctgccacctg gaaacaaagc attgaagtct gcagttgaaa agcccaacgt ctgtgagatc 2460 tgacttgctt attggtcatt gctagtatta tcgactcaga acctctttac taatggctag 2580 taaatcataa ttgagaaatt ctgaattttg acaaggtctc tgctgttgaa atggtaaatt 2640 tattattttt tttgtcatga taaattctgg ttcaaggtat gctatccatg aaataatttc 2700 tgaccaaaac taaattgatg caatttgatt atccatctta gcctacagat ggcatctggt 2760 aacttttgac tgttttaaaa aataaatcca ctatcagagt agatttgatg ttggcttcag 2820 aaacatttag aaaacaaaa gttcaaaaat gttttcagga ggtgataagt tgaataactc 2880 tacaatgtta gttctttgag ggggacaaaa aatttaaaat ctttgaaagg tcttatttta 2940 cagccatate taaattatet taagaaaatt tttaacaaag ggaatgaaat atatateatg 3000 attctgtttt tccaaaagta acctgaatat agcaatgaag ttcagttttg ttattggtag 3060 tttgggcaga gtctcttttt gcagcacctg ttgtctacca taattacaga ggacatttcc 3120 atgttctagc caagtatact attagaataa aaaaacttaa cattgagttg cttcaacagc 3180 atgaaactga gtccaaaaga ccaaatgaac aaacacatta atctctgatt atttatttta 3240 aatagaatat ttaattgtgt aagatctaat agtatcatta tacttaagca atcatattcc 3300 tgatgatcta tgggaaataa ctattattta attaatattg aaaccaggtt ttaagatgtg 3360 ttagccagtc ctgttactag taaatctctt tatttggaga gaaattttag attgttttgt 3420 tctccttatt agaaqgattg tagaaagaaa aaaatgacta attggagaaa aattggggat 3480 atatcatatt tcactgaatt caaaatgtct tcagttgtaa atcttaccat tattttacgt 3540 acctctaaga aataaaagtg cttctaatta aaatatgatg tcattaatta tgaaatactt 3600 cttgataaca gaagttttaa aatagccatc ttagaatcag tgaaatatgg taatgtatta 3660 ttttcctcct ttgagtnagg tcttgtgctt tttnttcctg gccactaaat ntcaccatnt 3720 ccaanaagca aantaaacct attctgaata tttttgctgt gaaacacttg ncagcagagc 3780 tttcccncca tqnnagaaqc ttcatqaqtc acacattaca tctttgggtt gattgaatgc 3840 cactgaaaca tttctagtag cctggagnag ttgacctacc tgtggagatg cctgccatta 3900 aatggcatcc tgatggctta atacacatca ctcttctgtg nagggtttta attttcaaca 3960 cagettaete tgtageatea tgtttaeatt gtatgtataa agattataen aaggtgeaat 4020 tgtgtatttc ttccttaaaa tgtatcagta taggatttag aatctccatg ttgaaactct 4080 aaatgcatag aaataaaaat aataaaaaat ttttcatttt ggcttttcag cctagtatta 4140 aaactgataa aagcaaagcc atgcacaaaa ctacctccct agagaaaggc tagtcccttt 4200 tetteeceat teattteatt atgaacatag tagaaaacag catattetta teaaatttga 4260 tgaaaagcgc caacacgttt gaactgaaat acgacttgtc atgtgaactg taccgaatgt 4320 ctacgtattc cacttttcct gctggggttc ctgtctcaga aaggagtctt gctcgtgctg 4380 gtttctatta cactggtgtg aatgacaagg tcaaatgctt ctgttgtggc ctgatgctgg 4440 ataactggaa aagaggagac agtcctactg aaaagcataa aaagttgtat cctagctgca 4500 gattegitea gagtetaaat teegttaaca aettggaage taceteteag cetaetttte 4560 cttcttcagt aacacattcc acacactcat tacttccggg tacagaaaac agtggatatt 4620 teegtggete ttattcaaac tetecateaa ateetgtaaa eteeagagea aateaagaat 4680 tttctgcctt gatgagaagt tcctacccct gtccaatgaa taacgaaaat gccagattac 4740 ttacttttca gacatggcca ttgacttttc tgtcgccaac agatctggca cgagcaggct 4800 tttactacat aggacctgga gacagagtgg cttgctttgc ctgtggtgga aaattgagca 4860

```
attgggaacc gaaggataat gctatgtcag aacacctgag acattttccc aaatgcccat 4920
ttatagaaaa tcagcttcaa gacacttcaa gatacacagt ttctaatctg agcatgcaga 4980
cacatgcage cegetttaaa acattettta actggeeete tagtgtteta gttaateetg 5040
agcagettge aagtgegggt tittattatg tgggtaacag tgatgatgte aaatgettit 5100
gctgtgatgg tggactcagg tgttgggaat ctggagatga tccatgggtt caacatgcca 5160
agtggtttcc aaggtgtgag tacttgataa gaattaaagg acaggagttc atccgtcaag 5220
ttcaagccag ttaccetcat ctacttgaac agetgetate cacatcagae ageccaggag 5280
atgaaaatgc agagtcatca attatccatt ttgaacctgg agaagaccat tcagaagatg 5340
caatcatgat gaatactcct gtgattaatg ctgccgtgga aatgggcttt agtagaagcc 5400
tggtaaaaca gacagttcag agaaaaatcc tagcaactgg agagaattat agactagtca 5460
atgatettgt gttagaetta etcaatgeag aagatgaaat aagggaagag gagagagaaa 5520
gagcaactga ggaaaaagaa tcaaatgatt tattattaat ccggaagaat agaatggcac 5580
tttttcaaca tttgacttgt gtaattccaa tcctggatag tctactaact gccggaatta 5640
ttaatgaaca agaacatgat gttattaaac agaagacaca gacgtcttta caagcaagag 5700
aactgattga tacgatttta gtaaaaggaa atattgcagc cactgtattc agaaactctc 5760
tgcaagaagc tgaagctgtg ttatatgagc atttatttgt gcaacaggac ataaaatata 5820
ttcccacaga agatgtttca gatctaccag tggaagaaca attgcggaga ctacaagaag 5880
aaagaacatg taaagtgtgt atggacaaag aagtgtccat agtgtttatt ccttgtggtc 5940
atctagtagt atgcaaagat tgtgctcctt ctttaagaaa gtgtcctatt tgtaggagta 6000
caatcaaggg tacagttcgt acatttcttt catgaagaag aaccaaaaca tcgtctaaac 6060
tttagaatta atttattaaa tgtattataa ctttaacttt tatcctaatt tggtttcctt 6120
aaaaltttta tttatttaca actcaaaaaa cattgttttg tgtaacatat ttatatatgt 6180
atctaaacca tatqaacata tattttttag aaactaagag aatgataggc ttttgttctt 6240
atgaacgaaa aagaggtagc actacaaaca caatattcaa tcaaaatttc agcattattg 6300
aaattgtaag tgaagtaaaa cttaagatat ttgagttaac ctttaagaat tttaaatatt 6360
ttggcattgt actaataccg ggaacatgaa gccaggtgtg gtggtatgtg cctgtagtcc 6420
caggetgagg caagagaatt acttgageee aggagtttga atecateetg ggeageatae 6480
tgagaccctg cctttaaaaa caaacagaac aaaaacaaaa caccagggac acatttctct 6540
gtcttttttg atcagtgtcc tatacatcga aggtgtgcat atatgttgaa tcacatttta 6600
qqqacatqqt qtttttataa agaattctgt gagaaaaaat ttaataaagc aaccaaaaaa 6660
aaaaaaaa
```

<210> 231 <211> 3000 <212> DNA <213> Homo sapiens

<400> 231

ttgcaggtac ttagaatttt tcctgagcca ccctctagag ggcagtgtta catatatat 60 tqtaattatc caqttacaac aaaaaaggg ctctcattca tgcatgaaaa tcagaaatat 120 ttcatactct taaagaacac attggaacca atattatgat taaaacatat tttgctaagc 180 aaagagatat taaaaattaa ttcattaaca ttctgaacat tttttaactt gtaaaaacaa 240 ctttgatgcc ttgaatatat aatgattcat tataacaatt atgcatagat tttaataatc 300 tgcatatttt atgctttcat gtttttccta attaatgatt tgacatggtt aataattata 360 atatattctq catcacaqtt tacatattta tgtaaaataa gcatttaaaa attattagtt 420 ttattctqcc tqcttaaata ttactttcct caaaaagaga aaacaaaaat gctagatttt 480 actttatgac ttgaatgatg tggtaatgtc gaactctagt atttagaatt agaatgtttc 540 ttagcggtcg tgtagttatt tttatgtcat aagtggataa tttgttagct cctataacaa 600 aagtetgttg cttgtgttte acattttgga tttcctaata taatgttete tttttagaaa 660 aggtggacaa gtcctatttt caagagaaga tgacttttaa cagttttgaa ggatctaaaa 720 cttgtgtacc tgcagacatc aataaggaag aagaatttgt agaagagttt aatagattaa 780 aaacttttgc taattttcca agtggtagtc ctgtttcagc atcaacactg gcacgagcag 840 ggtttcttta tactggtgaa ggagataccg tgcggtgctt tagttgtcat gcagctgtag 900 atagatggca atatggagac tcagcagttg gaagacacag gaaagtatcc ccaaattgca 960 gatttatcaa cggcttttat cttgaaaata gtgccacgca gtctacaaat tctggtatcc 1020 agaatggtca gtacaaagtt gaaaactatc tgggaagcag agatcatttt gccttagaca 1080

ggccatctga g	gacacatgca	gactatcttt	tgagaactgg	gcaggttgta	gatatatcag	1140
acaccatata	cccgaggaac	cctgccatgt	attgtgaaga	agctagatta	aagtcctttc	1200
agaactggcc	agactatgct	cacctaaccc	caagagagtt	agcaagtgct	ggactctact	1260
acacaggtat	tggtgaccaa	gtgcagtgct	tttgttgtgg	tggaaaactg	aaaaattggg	1320
aaccttgtga	tegtgeetgg	tcagaacaca	ggcgacactt	tcctaattgc	ttctttgttt	1380
tgggccggaa	tcttaatatt	cgaagtgaat	ctgatgctgt	gagttctgat	aggaatttcc	1440
caaattcaac	aaatcttcca	agaaatccat	ccatggcaga	ttatgaagca	cggatcttta	1500
cttttgggac	atggatatac	tcagttaaca	aggagcagct	tgcaagagct	ggattttatg	1560
ctttaggtga .	aggtgataaa	gtaaagtgct	ttcactgtgg	aggagggcta	actgattgga	1620
agcccagtga	agacccttgg	gaacaacatg	ctaaatggta	tccagggtgc	aaatatctgt	1680
tagaacagaa	gggacaagaa	tatataaaca	atattcattt	aactcattca	cttgaggagt	1740
gtctggtaag	aactactgag	aaaacaccat	cactaactag	aagaattgat	gataccatct	1800
tccaaaatcc	tatggtacaa	gaagctatac	gaatggggtt	cagtttcaag	gacattaaga	1860
aaataatgga	ggaaaaaatt	cagatatctg	ggagcaacta	taaatcactt	gaggttctgg	1920
ttgcagatct	agtgaatgct	cagaaagaca	gtatgcaaga	tgagtcaagt	cagacttcat	1980
tacagaaaga	gattagtact	gaagagcagc	taaggcgcct	gcaagaggag	aagctttgca	2040
aaatctgtat	ggatagaaat	attgctatcg	tttttgttcc	ttgtggacat	ctagtcactt	2100
gtaaacaatg	tgctgaagca	gttgacaagt	gtcccatgtg	ctacacagtc	attactttca	2160
agcaaaaaat	ttttatgtct	taatctaact	ctatagtagg	catgttatgt	tgttcttatt	2220
accctgattg	aatgtgtgat	gtgaactgac	tttaagtaat	caggattgaa	ttccattagc	2280
atttgctacc	aagtaggaaa,	aaaaatgtac	atggcagtgt	tttagttggc	aatataatct	2340
ttgaatttct	tgatttttca	gggtattagc	tgtattatcc	attttttta	ctgttattta	2400
attgaaacca	tagactaaga	ataagaagca	tcatactata	actgaacaca	atgtgtattc	2460
atagtatact	gatttaattt	ctaagtgtaa	gtgaattaat	catctggatt	ttttattctt	2520
ttcagatagg	cttaacaaat	ggagctttct	gtatataaat	gtggagatta	gagttaatct	2580
ccccaatcac	ataatttgtt	ttgtgtgaaa	aaggaataaa	ttgttccatg	ctggtggaaa	2640
gatagagatt	gtttttagag	gttggttgtt	gtgttttagg	attctgtcca	ttttctttta	2700
aagttataaa	cacgtacttg	tgcgaattat	ttttttaaag	tgatttgcca	tttttgaaag	2760
cgtatttaat	gatagaatac	tatcgagcca	acatgtactg	acatggaaag	atgtcaaaga	2820
tatgttaagt	gtaaaatgca	agtggcaaaa	cactatgtat	agtctgagcc	agatcaaagt	2880
atgtatgttt	ttaatatgca	tagaacaaaa	gatttggaaa	gatatacacc	aaactgttaa	2940
atgtggtttc	L	~~~~~~++	aaaaaaaaaa	caccataggg	attttataaa	3000